“
Mutation of
POT1 results in
increased levels
of telomere-
based genomic
rearrangements
”

Maria Blasco, Elias Campo, Carlos López-Otín and colleagues have found that protection of telomeres 1 (POT1), a component of the shelterin complex that caps the ends of telomeres, is mutated in a subset of patients with chronic lymphocytic leukaemia (CLL). Mutation of POT1 results in increased levels of telomere-based genomic rearrangements.

Initially, the authors examined exome sequence data from 127 matched tumour and normal samples from patients who were diagnosed with CLL but who had not started treatment. In addition to the known driver mutations, such as *SF3B1*, *NOTCH1* and *MYD88*, they found that approximately 5% of the samples had a mutation in *POT1* and that these mutations were point mutations. Sanger sequencing of *POT1* in an additional 214 samples from patients with CLL supported the initial findings and, overall, 12 somatic

point mutations in *POT1* were found in 341 samples. Interestingly, *POT1* mutations occurred exclusively in patients with a wild-type immunoglobulin heavy chain variable gene cluster (*IGHV@*), a biological status that correlates with a poorer prognosis. Indeed, the authors found that *POT1* mutations also correlate with a more advanced disease at diagnosis.

Three of the 12 somatic mutations resulted in a truncated protein, and the remaining nonsynonymous mutations were all predicted to have deleterious effects on the function of POT1. The majority of the mutations reside in the two oligonucleotide–oligosaccharide binding (OB) folds present in the amino terminus of POT1. These folds enable POT1 to bind to the telomeric single-stranded (ss) DNA sequence TTAGGG, suggesting that this crucial function is compromised in *POT1* mutants. Indeed, *POT1*

mutations found in other types of tumours also predominately occur in the OB folds. Immunofluorescence, chromatin immunoprecipitation and electrophoretic mobility shift assays indicated that mutations that disrupt the OB folds do not inhibit the localization of POT1 to telomere ends, which is perhaps consistent with the fact that POT1 binds to TPP1, another component of shelterin, through its carboxyl terminus, but that mutant POT1 is unable to bind telomeric ssDNA.

The authors also found that only one allele of *POT1* is mutated in CLL, suggesting that the mutations in *POT1* result in a dominant-negative protein. In agreement with this hypothesis, a wild-type *POT1* human fibrosarcoma cell line (HT1080) expressing exogenous mutant *POT1* proteins had longer telomeres, but there was no change in the activity or level of telomerase. Moreover, these cells also had an increased incidence of chromosomal aberrations, including the fusion of sister chromatids at metaphase, increased telomere fragility and significantly increased levels of chromosomal breaks and fusions. Similar results were evident in CLL cells from patients with *POT1* mutations, although there were no obvious differences in telomere lengths compared with patient samples with wild-type *POT1*. This probably reflects the fact that telomere lengths are highly variable in humans.

Collectively, these data indicate that *POT1* mutations result in telomere uncapping, leading to an increased incidence of telomere-related chromosomal fusions. It will be interesting to determine why *POT1* mutations only seem to occur in patients with wild-type *IGHV@*.

Nicola McCarthy

ORIGINAL RESEARCH PAPER Ramsay, A. J. et al.
POT1 mutations cause telomere dysfunction in
chronic lymphocytic leukemia. *Nature Genet.*
17 Mar 2013 (doi:10.1038/ng.2584)

CORRECTION**Leukaemia: ‘Gimme shelter’***Nicola McCarthy**Nature Reviews Cancer* **13**, 297 (2013)

In the original version of this article the DOI number was incorrect. This error has been corrected in the online version of the article.