PROGNOSTICS

Profiles of lung cancer

The outlook for patients presenting with advanced non-small-cell lung cancer is grim — the overall 10-year survival rate is 8–10%. Of the 25–30% who present with stage I disease, up to 50% will relapse within 5 years of surgical intervention. But now, David Beer and colleagues report in the August issue of *Nature Medicine* that the patients with stage I disease who are at high risk of relapse can be identified by gene-expression profiling. This information could be valuable for treatment planning.

The authors correlated gene-expression profiles with clinical outcome in patients with lung adenocarcinoma and identified specific gene patterns that predict survival in patients with stage I disease.

Gene-expression profiles were obtained for 86 primary lung adenocarcinomas and 10 non-tumour

lung samples, and 4,966 genes were identified. Use of hierarchical clustering yielded three clusters of tumours, and these groups showed significant differences with respect to tumour stage and tumour classification. This indicates that, as expected, tumours with similar differentiation characteristics show similar gene expression, and that many higher-stage tumours are often poorly differentiated and are associated with poor survival — most of these tumours were located in cluster three. Interestingly, eleven stage I tumours were also located in cluster three.

Of the 4,966 genes examined, 967 differed significantly between stage I and III adenocarcinomas. The authors verified the reliability of the expression data from the microarray analysis by northern blotting and immunohistochemistry of three genes that were selected arbitrarily. But the most important question was whether this expression data could be correlated with survival.

Survival curves showed significantly poorer survival among stage III compared with stage I adenocar-

DRUG TARGETING

Right on target

Although many drugs have been developed to interfere with tumour angiogenesis, it has been difficult to find ones that specifically target tumour blood vessels without affecting normal tissues. David Cheresh and colleagues have developed a gene-delivery strategy to block endothelial signalling pathways that are required for new blood-vessel formation, and report that this approach can be used to destroy established and metastatic tumours.

The angiogenic epithelium expresses several surface molecules that are potential therapeutic targets. One of these, the integrin $\alpha v \beta 3$, is a particularly good candidate because it is involved in cancer-cell invasion and has also been shown to mediate virus internalization — meaning it could mediate entry of other ligands into vascular endothelial cells. Cheresh and colleagues report in *Science* that a cationic polymerized lipid-based nanoparticle (NP) coupled to a small, organic integrin $\alpha v \beta 3$ ligand can be used to deliver therapeutic genes to

angiogenic blood vessels. But what genes are best for stopping blood-vessel growth?

The angiogenic factors basic fibroblast growth factor (bFGF) and vascular endothelial growth factor (VEGF) signal through a RAF1-mediated pathway, and blocking this pathway induces endothelial-cell apoptosis. A mutant form of Raf1 that fails to bind ATP (ATP^u-Raf) has been shown to block endothelial-cell Raf1 activity and inhibit angiogenesis *in vivo*. The authors coupled this mutant gene to $\alpha v \beta 3$ -NP, creating $\alpha v \beta 3$ -NP/Raf(-). When mice bearing established M21-L melanomas received a single injection of $\alpha v \beta 3$ -NP/Raf(-), the vector specifically targeted tumour blood vessels, where it induced apoptosis within 24 hours. By 72 hours, concentric rings of apoptotic tumour cells were observed around each apoptotic vessel, and the tumours began to necrose.

Consequently, treated mice displayed rapid tumour regression — in four out of six mice the tumours completely disappeared, and the other two showed a 95% reduction in tumour mass and a 75% suppression of blood-vessel density. Tumour regression was sustained for over 250 days. M21-L cells do

not express $\alpha v \beta 3$, indicating that the tumour regressed as a result of anti-angiogenic effects, rather than due to a direct effect on the cancer cells themselves.

But is this therapy effective against metastases? The authors injected mice with CT-26 colon adenocarcinoma cells, which led to the formation of lung or liver metastases, and 10 days later injected $\alpha v \beta 3$ -NP/Raf(-). Mice treated with control vector developed extensive lung and liver metastases, whereas treatment with $\alpha v \beta 3$ -NP/Raf(-) caused a regression and disappearance of these tumours. NPs are less immunogenic than viral vectors, so this molecule should be safer than other cancer gene-therapy approaches, and can also be used to deliver other types of therapeutic genes to blood vessels.

Kristine Novak

References and links

ORIGINAL RESEARCH PAPER Hood, J. D. *et al.* Tumour regression by targeted gene delivery to the neovasculature. *Science* **296**, 2404–2407 (2002)

FURTHER READING Hood, J. D. & Cheresh, D. A. Role of integrins in cell invasion and migration. *Nature Rev. Cancer* **2**, 91–100 (2002)

WEB SITE
David Cheresh's lab:
<http://www.scripps.edu/imm/cheresh/>