CORONARY ARTERY DISEASE ISCHEMIA AND TREATMENT CHOICE

A large, long-term observational series from the USA has demonstrated that patients with stable coronary artery disease (CAD) and minimal ischemia are best treated with medical therapy, whereas early revascularization is the best treatment strategy for those with substantial ischemia without extensive scarring.

Although guidelines for patients with CAD recommend ischemia testing and tailoring of management strategies on the basis of the findings, the efficacy of this approach was unproven.

Rory Hachamovitch and colleagues, therefore, set out to determine whether the extent of ischemia, as visualized by myocardial perfusion scintigraphy, should influence treatment choice in these patients.

Long-term follow-up (8.7 ± 3.3 years) was completed in 13,555 patients with known or suspected stable CAD. Of these patients, 12,329 received medical therapy and 1,226 underwent early revascularization (of which 41% and 59% underwent CABG surgery and PCI, respectively). Mortality for the whole patient population was 3.3% per year.

The extent and severity of ischemia was found to be the predominant factor predictive of early referral for revascularization in the patient population. In patients without prior CAD, in individuals with prior revascularization but no prior myocardial infarction, and in patients without extensive scarring (less than 10% myocardium), early revascularization was associated with better survival than medical therapy in those with more than ~10% myocardial ischemia, whereas optimal medical therapy was associated with better survival in those with less ischemia. No such relationship between extent of ischemia, therapy choice, and survival was observed in patients with extensive scarring from prior myocardial infarction.

Bryony M. Mearns

Original article Hachamovitch, R. *et al.* Impact of ischaemia and scar on the therapeutic benefit derived from myocardial revascularization vs. medical therapy among patients undergoing stress-rest myocardial perfusion scintigraphy. *Eur. Heart J.* doi:10.1093/eurheartj/ehq500