ANTICOAGULATION THERAPY

Enoxaparin dosing for VTE during pregnancy

Once-daily injection of enoxaparin (60–230 mg depending on body weight) is the optimum dosing strategy for the management of venous thromboembolism during pregnancy, according to the findings of a population pharmacokinetic modelling study by Patel and colleagues from King's College London, UK. The investigators propose that “current national and international guidelines should be reconsidered in light of our findings”.

Venous thromboembolism is one of several conditions for which pharmacological management is required during pregnancy. Low-molecular-weight heparins (LMWH), such as enoxaparin, have been shown to be safe for use in the antenatal period, but questions remain about optimum dosing, given that pregnancy can change the pharmacokinetic profile of the drug. An increase in glomerular filtration rate speeds up clearance, and rising plasma volume alters the volume of distribution of the drug.

To develop their model, Patel *et al.* enrolled 123 pregnant women aged ≥ 18 years who were prescribed enoxaparin and referred to the thrombophilia clinic at King's College Hospital. At each monthly clinic visit, participants provided blood samples for assessment of anti-factor Xa activity in plasma at three time points—trough (the lowest concentration before the next dose), and 1 h and 3 h after enoxaparin injection. A total

of 795 anti-factor Xa activity measurements (712 antenatal) were incorporated into the nonlinear, mixed effects model (an approach that allows the use of small samples). The model was adjusted for the covariates of body weight, prepregnancy lean body weight, creatinine clearance, and gestational stage. Pharmacokinetic simulations revealed that the target peak plasma concentrations of enoxaparin would be reached with either once-daily or twice-daily dosing. However, anti-factor Xa trough activity was shown to increase during gestation, effectively prolonging the half-life of enoxaparin as pregnancy progresses. Therefore, once-daily dosing was deemed to be the optimum strategy. “We acknowledge that some clinicians may feel apprehensive about employing a once-daily LMWH regimen during the first few weeks following diagnosis, particularly in those women who suffer from extensive thrombosis or who suffer their event early on during pregnancy,” write the investigators. “In such situations, an initial twice-daily strategy, followed by once[-daily] LMWH might be justifiable.”

Commenting on this study, Professor Jeff Ginsberg from McMaster University (Hamilton, ON, Canada) raises a point about the use of anti-factor Xa levels as a surrogate for clinical events: “a therapeutic range for this laboratory test has never been validated,” he cautions. “It concerns me that clinicians might extrapolate these results to pregnant women with mechanical heart valves,” he continues, “a group that might be more susceptible to treatment failure with LMWH.” Ultimately, Professor Ginsberg believes that a change in the guidelines should not be considered before an adequately powered randomized trial of once-daily versus twice-daily dosing of enoxaparin, combined with substudies to validate the link between anti-factor Xa levels and bleeding or thrombosis, have been conducted.

Alexandra Roberts

“...once-daily dosing was deemed to be the optimum strategy...”

Original article Patel, J. P. *et al.* Population pharmacokinetics of enoxaparin during the antenatal period. *Circulation* doi:10.1161/CIRCULATIONAHA.113.003198

© PhotoDisc/Getty Images