

IN BRIEF

EPILEPSY**HSP90 inhibition suppresses seizures**

Loss of the glutamate transporter GLT1 (also known as SLC1A2) occurs in temporal lobe epilepsy (TLE), but the mechanisms mediating GLT1 degradation are not understood. Here, Sha *et al.* report upregulated expression of HSP90 β in reactive astrocytes of human epileptogenic tissue and in mouse models of TLE and febrile seizures. In cultured astrocytes, HSP90 β negatively regulated GLT1 protein levels through proteasome-dependent GLT1 degradation, which was prevented by the HSP90 inhibitor, 17AAG. In mouse models of TLE, long-term systemic administration of 17AAG suppressed spontaneous recurrent seizures and ameliorated astroglialosis.

ORIGINAL ARTICLE Sha, L. *et al.* Pharmacologic inhibition of Hsp90 to prevent GLT-1 degradation as an effective therapy for epilepsy. *J. Exp. Med.* <http://dx.doi.org/10.1084/jem.20160667> (2016)

ALZHEIMER DISEASE**Identification of novel A β inhibitors**

Targeting amyloid- β (A β) aggregation and accumulation has been pursued as a major potential therapeutic strategy against AD, but no compound has yet gained regulatory approval. Habchi *et al.* now describe the use of a quasi-structure-based and kinetics-based drug discovery approach to identify a pool of ligands for retinoid acid receptors and retinoid X receptors that inhibit aggregation of the 42-residue form of A β (A β ₄₂). The compounds inhibited A β ₄₂ aggregation in human cerebrospinal fluid solutions and rescued A β ₄₂-mediated dysfunction in a *Caenorhabditis elegans* model of A β ₄₂-mediated cytotoxicity.

ORIGINAL ARTICLE Habchi, J. *et al.* Systematic development of small molecules to inhibit specific microscopic steps of A β 42 aggregation in Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **114**, E200–E208 (2016)

NEURODEGENERATIVE DISEASE**Pituitary adenylate cyclase activator ameliorates SBMA**

Post-translational modifications, such as phosphorylation, modify the toxicity of the polyglutamine (polyQ) expansion in the androgen receptor (AR) (polyQ-AR), which occurs in spinobulbar muscular atrophy (SBMA). Here, Polanco *et al.* show in cell models that cyclin-dependent kinase 2 phosphorylates polyQ-AR specifically at Ser96 and this is negatively regulated by the adenylate cyclase–protein kinase A signalling pathway. Chronic intranasal administration of an analogue of pituitary adenylate cyclase-activating polypeptide to a mouse model of SBMA reduced Ser96 phosphorylation, promoted polyQ-AR degradation and ameliorated the disease outcome.

ORIGINAL ARTICLE Polanco, M. J. *et al.* Adenylyl cyclase activating polypeptide reduces phosphorylation and toxicity of the polyglutamine-expanded androgen receptor in spinobulbar muscular atrophy. *Sci. Transl Med.* **8**, 370ra181 (2016)

HIV**CRISPR screen identifies novel therapeutic targets**

Host proteins, termed host dependency factors (HDFs), are crucial for productive HIV infection but dispensable for cellular viability, thereby representing promising therapeutic targets. Here, Park *et al.* conduct a CRISPR-based genetic screen in a CD4⁺ T cell line to identify five HDFs (the HIV co-receptors CD4 and CCR5, as well as TPST2, SLC35B2, and ALCAM) that, when inactivated, conferred protection from HIV infection. Using the CD4⁺ T cell line and primary CD4⁺ T cells, TPST2 and SLC35B2 were shown to facilitate CCR5 recognition by the HIV envelope, whereas ALCAM mediated cell aggregation, which is required for cell-to-cell HIV transmission.

ORIGINAL ARTICLE Park, R. J. *et al.* A genome-wide CRISPR screen identifies a restricted set of HIV host dependency factors. *Nat. Genet.* <http://dx.doi.org/10.1038/ng.3741> (2016)