HIGHLIGHTS

IN BRIEF

HYPERLIPIDAEMIC DISEASE

Angptl3 regulates lipid metabolism in mice

Koishi, R. *et al. Nature Genet.* 2002 Jan 14 (DOI 10.1038/ng814)

Several lipoprotein deficiencies are known to arise from genetic disorders of lipid metabolism. KK obese mice show signs of hyperinsulinaemia and hyperglycaemia, and have low levels of lipid in their plasma due to a recessively inherited locus, named *hypl*. Koishi and colleagues identified a mutation in the gene *Angptl3* as being the cause of the *hypl* trait. Overexpression of the *Angptl3* protein resulted in a marked increase in plasma lipid levels. As *Angptl3* is secreted, it might prove to be a more accessible therapeutic target than many other proteins that have been implicated in lipid disorders, which tend to be cloistered in lysosomes or other vesicles.

NEURODEGENERATIVE DISEASE

Recruitment and activation of caspase-8 by the huntingtin-interacting protein Hip-1 and a novel partner Hippi

Gervais, F. G. *et al. Nature Cell Biol.* 2002 Jan 14 (DOI 10.1038/ncb735)

Huntington's disease is caused by the expression of a mutant huntingtin protein (HTT). Normally, HTT interacts with the HIP1 protein. Nicholson and colleagues show that the affinity of HIP1 for mutant HTT is much lower than for wild-type HTT. They also identify another partner for HIP1, HIPPI. In diseased brains, levels of the HIP1–HIPPI complex are increased relative to levels of the HIP1–HTT complex. The HIP1–HIPPI complex initiates the apoptotic cascade by recruiting procaspase-8. Elucidation of this pathway could pave the way for new strategies to interfere with neurodegeneration.

STRUCTURE-BASED DRUG DESIGN

Design and synthesis of non-peptidic inhibitors for the Syk C-terminal SH2 domain based on structure-based *in-silico* screening

Niimi, T. *et al. J. Med. Chem.* 44, 4737–4740 (2001)

The tyrosine kinase SYK is important in the cellular response to antigen binding to antibody receptors, and is thus a potential target for immune suppression. SYK-kinase activity is stimulated by the interaction of its SH2 domains with phosphorylated tyrosine residues on an activated receptor, but the phosphotyrosyl group has limited use in inhibitor design owing to its hydrolytic lability and low membrane permeability.

Niimi *et al.* used a structure-based approach to assemble fragments into a non-peptidic non-phosphorous-containing ligand for the SH2 domain, with an affinity comparable to that of a natural ligand mimic.

PROTEIN–PROTEIN INTERACTIONS

Preventing tangles

A tangled chain is not particularly useful for carrying out its usual tasks. Similarly, when proteins get tangled up, interacting with each other to form oligomers, some of them might function incorrectly, or even cause harm. Oligomerization is essential for the oncogenicity of the BCR–ABL oncoprotein, which is responsible for a wide range of human leukaemias. In the February issue of *Nature Structural Biology*, Peter Kim and colleagues describe the crystal structure for the N-terminal oligomerization domain of BCR–ABL, together with a new mode of oligomer formation. This N-terminal structure is a promising target for the design of oligomerization inhibitors that could disrupt the transforming activity of BCR–ABL.

The BCR–ABL oncogene is formed by the fusion of the BCR (breakpoint cluster region) gene on chromosome 22 with the proto-oncogene ABL (Abelson murine leukaemia viral oncogene homologue) on chromosome 9. The activity of cytoplasmic ABL (c-ABL), a tyrosine kinase, is normally under tight control. However, fusion of the BCR sequences constitutively activates ABL, which is an essential step for BCR–ABL transformation. The first specific kinase inhibitor to be developed, *Gleevec*, targets the ABL kinase domain and has been shown to be clinically effective in the treatment of chronic myelogenous leukaemia. The N-terminal BCR oligomerization domain is also essential for ABL activation.

The structure showed that BCR–ABL first forms dimers, which then dimerize to form tetramers. Initially, two monomers form a coiled–coil motif between C-terminal α -helices and swap N-terminal α -helices. Subsequently, dimers stack onto each other to form tetramers, the four α -helices lying approximately in the same plane. The use of both coiled–coil packing and domain swapping, each recognized as being important mechanisms of oligomerization, leads to a very stable structure. Clustering of kinase domains is thought to be sufficient to induce kinase activation. The crystal structure indicates that a tetramer might be even more efficient than a dimer at bringing domains together.

Protein–protein interactions are ubiquitous in many diseases and are likely to deliver a wealth of new sites for drug targeting. The few examples of drugs that inhibit such interactions show that this can be a successful strategy. Dimerized domains of anti- and pro-apoptotic proteins block apoptosis, but small-molecule inhibitors of dimerization have been shown to induce cell death. Inhibition of the human immunodeficiency virus 1 (HIV-1) gp41 fusion protein with various peptides provides a further proof of principle. Coiled–coils are well-studied protein–protein interface motifs, and the solved structure of the oligomerization domain of BCR–ABL provides a template for the rational design of inhibitors to disrupt oligomerization.

Melanie Brazil

References and links

ORIGINAL RESEARCH PAPER Zhao, X. *et al.* Structure of the Bcr–Abl oncoprotein oligomerization domain. *Nature Struct. Biol.* 9, 117–120 (2002)

FURTHER READING Eckert, D. M. & Kim, P. S. Mechanisms of viral membrane fusion and its inhibition. *Annu. Rev. Biochem.* 70, 777–810 (2001) | Degterev, A. *et al.* Identification of small-molecule inhibitors of interaction between the BH3 domain and BCL-X_L. *Nature Cell Biol.* 3, 173–182 (2001)

WEB SITE Kim's lab: <http://web.wi.mit.edu/kim/>