PEDIATRICS

Congenital hypothyroidism: are cases being missed?

The incidence of congenital hypothyroidism is twice as high as currently estimated, according to results published by Corbetta and coworkers. The Italian study also shows, unexpectedly, that inborn functional defects are more prevalent than dysgenetic forms of the disease.

Congenital hypothyroidism can lead to severe mental retardation and growth failure, unless treated promptly by levothyroxine. The National Academy of Clinical Biochemistry recommends further investigations for newborn babies with dried-blood-spot TSH levels >10 mU/l—a lower cutoff value than previously used, which reflects the increased sensitivity of today's TSH assays. In line with these recommendations, a screening center in Lombardy, Italy, shifted their dried-blood-spot TSH screening cutoff level from 20 mU/l to 12 mU/l in 1999 and to 10 mU/l in 2002. Now, Corbetta and co-investigators reveal the impact of this new screening strategy.

The retrospective study involved 629,042 newborn babies screened

between 1999 and 2002. The screening program identified 435 babies who were confirmed to have congenital hypothyroidism—an incidence of 1:1,446 live births, double previously reported estimates. The researchers were concerned to find that 45% of newborn babies with congenital hypothyroidism would have been missed had the previous cutoff of 20 mU/l been used; furthermore, a number of these babies had thyroid ectopy or hemiagenesis. Among the confirmed cases of congenital hypothyroidism, 68% of babies had a gland *in situ* form and 32% had thyroid dysgenesis, a surprising result given current understanding that about 75% of affected individuals have thyroid dysgenesis. Clinical re-evaluation of a representative group of the children with congenital hypothyroidism revealed permanent thyroid dysfunction in 78% of 59 children aged 3–5 years.

Premature birth was a highly prevalent factor among the cases (20%). The 3–5-fold increased risk of gland *in situ* congenital hypothyroidism found among prematurely born babies in the study

© Dreamstime®

prompts the investigators to recommend additional testing of TSH in such babies. The investigators highlight the rationale for the use of low TSH cutoffs for screening of congenital hypothyroidism. Encouragingly, this new screening strategy increased the number of tests per neonate per year only by an estimated 22% versus the 20 mU/l cutoff during the period studied.

Carol Wilson

Original article Corbetta, C. *et al.* A 7-year experience with low blood TSH cutoff levels for neonatal screening reveals an unsuspected frequency of congenital hypothyroidism. *Clin. Endocrinol. (Oxf.)* doi:10.1111/j.1365-2265.2009.03568.x