REPRODUCTIVE ENDOCRINOLOGY
**BISPHENOL A
AND PCOS**

Levels of bisphenol A (BPA) are significantly higher in women with polycystic ovary syndrome (PCOS) than in normally ovulating, nonhyperandrogenemic women, independent of the degree of obesity, new data from a cross-sectional study reveal.

“BPA is a potent endocrine disrupting chemical, and it has been shown that the ovary is sensitive to increased BPA exposure,” says Evanthia Diamanti-Kandarakis of the Medical School University of Athens. “Furthermore, there is a bidirectional relationship between levels of BPA and androgens, since BPA increases concentrations of circulating free androgens, whereas androgens increase BPA levels.”

The researchers determined levels of BPA and hormones and metabolic parameters in 71 women with PCOS and 100 age-matched and BMI-matched normally ovulating, nonhyperandrogenemic women.

Serum levels of BPA were significantly higher in the PCOS group than the control group. Moreover, lean (BMI <25 kg/m²) or overweight (BMI ≥25 kg/m²) women with PCOS had significantly higher levels of BPA, respectively, than lean or overweight women in the control group. Testosterone levels and insulin resistance were also positively correlated with levels of BPA in the entire study cohort and PCOS groups, respectively. Lastly, multiple regression analysis revealed a strong association of BPA levels with the presence of PCOS. “In susceptible, dysfunctional gonads, BPA exposure may lead to subsequent PCOS development, as experiments in rats have already shown,” comments Diamanti-Kandarakis.

The findings emphasize the impact of everyday exposure to BPA on reproductive function in women with pre-existing anovulatory or hyperandrogenic derangements, such as PCOS, Diamanti-Kandarakis concludes.

Carol Wilson

Original article Kandaraki, E. *et al.* Endocrine disruptors and polycystic ovary syndrome (PCOS): elevated serum levels of bisphenol A in women with PCOS. *J. Clin. Endocrinol. Metab.* doi:10.1210/jc.2010-1658