Nature Reviews Endocrinology **8**, 628 (2012); published online 28 August 2012;
doi:10.1038/nrendo.2012.159;
doi:10.1038/nrendo.2012.161;
doi:10.1038/nrendo.2012.160;
doi:10.1038/nrendo.2012.162

IN BRIEF

METABOLISM

A role for neutrophils in inflammation-induced metabolic disease in mice

Chronic, low-grade inflammation is a feature of obesity and type 2 diabetes mellitus. Neutrophils secrete neutrophil elastase in response to inflammation, which can exacerbate inflammatory responses. Talukdar *et al.* found that deletion of neutrophil elastase in mice with obesity induced by a high-fat diet reduced tissue inflammation, improved glucose tolerance and increased insulin sensitivity. The researchers conclude that neutrophils participate in inflammation-induced metabolic disease.

Original article Talukdar, S. *et al.* Neutrophils mediate insulin resistance in mice fed a high-fat diet through secreted elastase. *Nat. Med.* doi:10.1038/nm.2885

CARDIOVASCULAR ENDOCRINOLOGY

Serum lipid levels have improved in children in the USA

Kit and co-workers examined trends in lipid concentrations in 16,116 US children aged 6–19 years in the time periods 1988–1994 and 2007–2010. The researchers found that mean levels of serum total cholesterol and non-HDL cholesterol had decreased, whereas mean levels of HDL cholesterol had increased from 1988–1994 to 2007–2010. In children aged 12–19 years, mean levels of LDL cholesterol and triglycerides had decreased from 1988–1994 to 2007–2010. Although lipid levels have improved in the studied time frame, the researchers stress that adverse lipid profiles are still observed; for example, almost 1 in 10 children had elevated levels of total cholesterol in 2007–2010.

Original article Kit, B. K. *et al.* Trends in serum lipids among US youths aged 6 to 19 years, 1988–2010. *JAMA* **308**, 591–600 (2012)

METABOLISM

Mice lacking gp78 in the liver are protected from diet-induced obesity

Liu *et al.* produced liver-specific gp78 knockout mice. Production of FGF-21 in the liver of these mice was increased, which activated thermogenesis in brown adipocytes. The mice had reduced lipid levels, improved insulin resistance and protection from diet-induced or age-induced obesity. The researchers conclude that gp78, a ubiquitin ligase, has an important role in regulating lipid biosynthesis in the liver and is a potential target for the treatment of metabolic diseases.

Original article Liu, T.-F. *et al.* Ablation of gp78 in liver improves hyperlipidemia and insulin resistance by inhibiting SREBP to decrease lipid biosynthesis. *Cell Metab.* doi:10.1016/j.cmet.2012.06.014

DIABETES

Effects of obesity and ageing on β -cell mass

Saisho *et al.* investigated the effects of ageing and obesity on β -cell mass and turnover by examining pancreatic samples obtained from human autopsies of individuals without diabetes mellitus who were aged 20–102 years. The β -cell mass in samples from individuals with obesity ($n=61$) was ~50% increased in comparison with that in samples from lean individuals ($n=53$). By examination of samples from 106 lean individuals, the researchers also showed that β -cell mass is preserved in old age.

Original article Saisho, Y. *et al.* β -cell mass and turnover in humans: effects of obesity and aging. *Diabetes Care* doi:10.2337/dc12-0421