Vitamin D and human health: more than just bone

John H. White

In their Perspective article ([Common misconceptions about vitamin D—implications for clinicians. *Nat Rev. Endocrinol.* doi:10.1038/nrendo.2013.75](#)),¹ Rosen and Taylor have provided a primer on the state of vitamin D research. However, the analyses presented of the current literature are not balanced. The conclusions of the authors largely reflect those of the 2011 Institute of Medicine (IOM) report,² which immediately created a controversy, leading to published rebuttals.³ The IOM report was controversial in part because of its very conservative recommendations for dietary intakes, its conclusions of insufficient evidence for any role of vitamin D in nonbone health, and the way it presented evidence for the potential harm associated with circulating 25-hydroxyvitamin D levels.

Notably, considerable debate surrounds potential roles of vitamin D in nonbone indications including cancer prevention and control of immune system function. The authors conclude that “effects of vitamin D on nonbone disorders is currently best described as consisting of hypotheses of emerging interest”.¹ They also claim that vitamin D supplementation has not been shown to prevent infections. In fact, several randomized placebo-controlled trials have been published providing evidence for vitamin D supplementation of deficient populations in preventing a variety of infections. In addition to the (highly cited) results of a trial published in 2010 concluding that

vitamin D supplementation reduced the risk of seasonal influenza infections in a paediatric population,⁴ recent studies have provided evidence for a beneficial role of vitamin D supplementation of populations at risk for upper respiratory tract or ear infections, either because of severe vitamin D deficiency or a history of recurrent infections.^{5,6} By contrast, one study that produced a negative result involved a healthy population with baseline 25-hydroxyvitamin D levels of 73 nmol/l.⁷ The authors also discuss the emerging evidence for a U-shaped curve of risk associated with 25-hydroxyvitamin D levels and focus largely on the potential risks associated with excessive vitamin D intake, an area that should not be ignored. They cite a recent review⁸ suggesting that there may be an increase in prevalence of certain cancers associated with high serum 25-hydroxyvitamin D levels (>75 nmol/l). However, U-shaped curves have two sides, and the IOM recommends that the vitamin D needs of ~50% of the population can be met with 25-hydroxyvitamin D levels of 30–40 nmol/l, whereas those of most of the remainder require 40–50 nmol/l. If followed, these recommendations would place the entire population solidly within the left-hand side of the U-shaped curve of cancer risk. It can therefore be argued that one of the misconceptions arising from conservative estimates of recommended vitamin D intake in the IOM report is that less vitamin D does no harm.

Departments of Physiology and Medicine,
McGill University, 3655 Drummond Street,
Room 1112, H3G 1Y6 Montreal, QC, Canada.
john.white@mcgill.ca

Competing interests

The author declares no competing interests.

1. Rosen, C. J. & Taylor, C. L. Common misconceptions about vitamin D—implications for clinicians. *Nat. Rev. Endocrinol.* **9**, 434–438 (2013).
2. Institute of Medicine. *Dietary Reference Intakes for Calcium and Vitamin D* (eds Ross, A. C., Taylor, C. L., Yaktine, A. L. & Del Valle, H. B.) (National Academies Press, Washington DC, 2011).
3. Holick, M. F. et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J. Clin. Endocrinol. Metab.* **96**, 1911–1930 (2011).
4. Camargo, C. A. Jr. et al. Randomized trial of vitamin D supplementation and risk of acute respiratory infection in Mongolia. *Pediatrics* **130**, 561–567 (2012).
5. Bergman, P. et al. Vitamin D₃ supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study. *BMJ Open* **2**, e001663 (2012).
6. Marchisio, P. et al. Vitamin D supplementation reduces the risk of acute otitis media in otitis-prone children. *Pediatr. Infect. Dis. J.* <http://dx.doi.org/10.1097/INF.0b013e31829be0b0>.
7. Murdoch, D. R. et al. Effect of vitamin D3 supplementation on upper respiratory tract infections in healthy adults: the VIDARIS randomized controlled trial. *JAMA* **308**, 1333–1339 (2012).
8. Mayne, S. T., Ferrucci, L. M. & Cartmel, B. Lessons learned from randomized clinical trials of micronutrient supplementation for cancer prevention. *Annu. Rev. Nutr.* **32**, 369–390 (2012).