BONE

KEY ROLE FOR S1P IN BONE REMODELLING

New research published in *Nature Communications* shows that the hormone calcitonin negatively regulates bone formation by inhibiting the release of the anabolic bone factor sphingosine 1-phosphate (S1P) from osteoclasts. The findings highlight the existence of a novel mechanism involved in bone remodelling.

Although calcitonin pharmacologically inhibits bone resorption, its physiological role is uncertain as calcitonin-deficient mice have increased bone formation even though the calcitonin receptor (CT-R) is not expressed on osteoblasts. “To clarify the cellular and molecular mechanisms of calcitonin action in bone remodelling, we generated a mouse model that enabled cell-type-specific inactivation of CT-R,” explains lead investigator Michael Amling.

Transgenic mice with ubiquitous or osteoclast-specific inactivation of CT-R had both increased trabecular bone mass and bone formation, in comparison with wild-type littermates. Mechanistic insight into how CT-R in osteoclasts regulates bone formation was provided by transcriptional profiling of calcitonin-treated osteoclasts isolated from CT-R-deficient and wild-type mice. Of the differentially expressed genes identified, *Spns2* (which encodes a transporter for S1P) was found to be negatively regulated by calcitonin in wild-type osteoclasts. This finding, together with the increased levels of S1P present in CT-R-deficient mice, suggests that calcitonin regulation of bone formation is mediated via S1P release from osteoclasts.

Further observations showed that mice deficient in both CT-R and the S1P receptor 3 (S1PR3) had a normal skeletal phenotype. The potential clinical implications of this finding were further supported by the observation that a nonselective S1PR agonist induced bone formation in wild-type mice but not in S1PR3-deficient mice.

“S1PR3 agonists could potentially be used to treat patients with bone loss, such as those with osteoporosis,” speculates Amling. “It is important to note that S1PR3-deficient mice displayed no obvious abnormalities in other organs, which indicates that drugs targeting S1PR3 could potentially have minimal adverse effects,” he adds.

David Holmes

Original article Keller, J. *et al.* Calcitonin controls bone formation by inhibiting the release of sphingosine 1-phosphate from osteoclasts. *Nat. Commun.* doi:10.1038/ncomms6215