RESEARCH HIGHLIGHTS

HIGHLIGHT ADVISORS

SEAN B. CARROLL

UNIVERSITY OF WISCONSIN,
USA

NANCY J. COX

UNIVERSITY OF CHICAGO, USA

RALPH J. GREENSPAN

THE NEUROSCIENCES
INSTITUTE, CALIFORNIA, USA

YOSHIHIDE HAYASHIZAKI

RIKEN GENOMIC SCIENCES
CENTER, JAPAN

PETER KOOPMAN

UNIVERSITY OF QUEENSLAND,
AUSTRALIA

LEONID KRUGLYAK

FRED HUTCHINSON CANCER
RESEARCH CENTER, USA

BARBARA MEYER

UNIVERSITY OF CALIFORNIA,
BERKELEY, USA

JOHN QUAKENBUSH

THE INSTITUTE FOR GENOMIC
RESEARCH, USA

NADIA ROSENTHAL

EMBL MONTEROTONDO, ITALY

NORIYUKI SATOH

KYOTO UNIVERSITY, JAPAN

MARC VIDAL

DANA-FARBER CANCER
INSTITUTE, BOSTON, USA

VIRGINIA WALBOT

STANFORD UNIVERSITY, USA

DETLEF WEIGEL

MAX PLANCK INSTITUTE FOR
DEVELOPMENTAL BIOLOGY,
GERMANY

PHIL ZAMORE

UNIVERSITY OF
MASSACHUSETTS, USA

LEONARD I. ZON

CHILDREN'S HOSPITAL,
BOSTON, USA

GENOMICS

Closing the gap

A new systematic approach to validating computational gene predictions in the *Drosophila* genome indicates that the number of unrecognized genes in sequenced eukaryotic genomes is probably a lot smaller than expected.

It was thought that the availability of full genome sequences would largely resolve the heated debates about just how many genes were required to make an organism. However, disagreements remained over how good our computational methods for predicting genes from sequences were. The *Drosophila* genome was a case in point: with large numbers of gene predictions falling in intergenic regions and no agreed standard way of finding out whether these were real, the gene-number topic remained a fertile breeding ground for dissent.

In their well-thought-out study that was designed to finally settle the arguments, Mark Yandell, Adina Bailey, Sima Misra and colleagues assembled a large collection of *Drosophila melanogaster* multiexon gene predictions in intergenic regions, which were identified from either computer predictions (9,811 from GENSCAN and 375 from FGENESH) or microarray-based validation experiments (1,266).

The authors then tested a subset of these predictions, subdivided into groups that were based on different criteria, to see what proportion of them corresponded to real gene transcripts. A gene prediction was

considered to be valid if sequences — obtained from an RT-PCR product which was generated from primers designed for sites in a pair of predicted exons — reflected transcript splicing.

In general, the gene predictions proved somewhat less than reliable: 16.4% of those tested were valid (122 valid/744 tested), and splice-site conservation proved to be by far the best criterion for prioritizing predictions for validation (64/174; 37.4%). Extrapolating their result to the entire collection of *Drosophila* genome gene predictions, the authors predict approximately 700 additional

protein-coding genes: a number that settles the *Drosophila* gene number argument at the low end of the spectrum: at ~14,000 genes.

The lessons for those looking for unrecognized protein-coding genes in other genome sequences are twofold: take those computer predictions with a large grain of salt and take a systematic experimental approach to a subset of these predictions if you want to get a true picture of what you are missing.

Nick Campbell

NPG Executive Editor, Heredity

References and links

ORIGINAL RESEARCH PAPER

Yandell, M. *et al.*
A computational and experimental approach to validating annotations and gene predictions in the *Drosophila melanogaster* genome. *Proc. Natl Acad. Sci. USA* **102**, 1566–1571 (2005)

FURTHER READING Zhang, M. Q. Computational prediction of eukaryotic protein-coding genes. *Nature Rev. Genet.* **3**, 698–709 (2002)

WEB SITE

Gerald Rubin's laboratory:

<http://mcb.berkeley.edu/faculty/GEN/rubing.html>