EVO-DEVO

Doing the time warp

This study describes a clever way of using gene expression data to study evolutionary novelties ...

Evolutionary developmental biology is being revolutionized by increasingly extensive and detailed gene expression data for a growing range of species. A recent study demonstrates a new way of investigating evolutionary novelties by looking for clusters of genes that are expressed differently between fruitflies and mosquitoes at particular times during development.

An important challenge for comparing temporal patterns of gene expression is that the time course of development can vary widely between species. Indeed, embryonic development in the mosquito *Anopheles gambiae* takes about twice

as long as in the fruitfly *Drosophila melanogaster*. Papatsenko and colleagues tackled this challenge using a computational method known as 'time warping' to align embryonic gene expression profiles from the two organisms. This alignment revealed that orthologous mosquito and fruitfly developmental genes are generally expressed at very similar times, despite the difference in the duration of development.

Might groups of genes that deviate from this temporal concordance underlie phenotypic differences between the two species? Using a computational approach, the authors built a comparative map of

development in which clusters of discordant genes are mapped according to the time and the duration of developmental processes. One known difference between mosquito and fruitfly embryos is that mosquitoes have a distinct amnion and serosa, whereas fruitflies have a single amnioserosa. Indeed, serosa genes stood out as a unique cluster on the authors' discordance map at a specific time during development. After optimizing their approach using the serosa genes, the authors identified several other groups of discordant genes. For example, one such cluster was enriched for metabolic genes that are expressed in the fruitfly yolk but are maternally expressed in mosquitoes, which might indicate an important difference in yolk metabolism.

This study describes a clever way of putting gene expression data to use to study evolutionary novelties, which should be applicable to other species with overall similar patterns of development.

Louisa Flintoft

ORIGINAL RESEARCH PAPER Papatsenko, D., Levine, M. & Goltsev, Y. Clusters of temporal discordances reveal distinct embryonic patterning mechanisms in *Drosophila* and *Anopheles*. *PLoS Biol.* **9**, e1000584 (2011)