GENE EXPRESSION

Transposon terminators

This study shows that intronic retrotransposon insertions can have substantial effects on gene expression over several kilobases and even *in trans*.

In mice, endogenous retrovirus (ERV) insertions are thought to account for several of the phenotypic differences among strains. Li *et al.* mapped polymorphic ERV insertions in diverse mouse strains by sequencing. They found that ERV polymorphisms are depleted from introns and from genes with roles in early development.

Their relative depletion from introns suggested that intronic ERV insertions might have deleterious effects, so the authors analysed the effects of an intronic insertion in *Slc15a2*, which encodes a peptide transporter. They found that this intronic ERV results in a ~49-fold increase in prematurely terminated *Slc15a2* transcripts. The termination can occur >12.5 kb upstream of the insertion site, indicating that the ERV element has long-range effects. Surprisingly, in heterozygous mice, the presence of the insertion on one allele can increase premature termination on the other allele.

The intronic ERV insertion also decreases expression of the full-length transcript and reduces protein levels

and activity *in vivo*. This disruption can occur *in trans* in heterozygotes, but the decrease was only seen when the insertion was on the paternal allele.

The authors suggest that this parent-of-origin effect might be linked to differential DNA methylation, as the 5' long terminal repeat of the ERV is only densely methylated when it is paternally inherited.

The authors found similar disrupted gene expression associated with other intronic ERV insertions, emphasizing the need to consider such variants when studying gene expression differences among individuals.

Mary Muers

ORIGINAL RESEARCH PAPER Li, J. *et al.* Mouse endogenous retroviruses can trigger premature transcriptional termination at a distance. *Genome Res.* 23 Feb 2012 (doi:10.1101/gr.130740.111)