HIGHLIGHTS

HIGHLIGHT ADVISORS

KONRAD BASLER

UNIVERSITY OF ZURICH,
SWITZERLAND

WENDY BICKMORE

MRC HUMAN GENETICS UNIT,
UK

PEER BORK

EMBL, GERMANY

SEAN B. CARROLL

UNIVERSITY OF WISCONSIN,
USA

ADAM EYRE-WALKER

UNIVERSITY OF SUSSEX, UK

JANE GITSCHIER

UNIVERSITY OF CALIFORNIA,
SAN FRANCISCO, USA

SARAH HAKE

UNIVERSITY OF CALIFORNIA,
BERKELEY, USA

PETER KOOPMAN

UNIVERSITY OF QUEENSLAND,
AUSTRALIA

LEONID KRUGLYAK

FRED HUTCHINSON CANCER
RESEARCH CENTER, USA

GAIL MARTIN

UNIVERSITY OF CALIFORNIA,
SAN FRANCISCO, USA

BARBARA MEYER

UNIVERSITY OF CALIFORNIA,
BERKELEY, USA

TOSHIHIKO SHIROISHI

NATIONAL INSTITUTE OF
GENETICS, JAPAN

KEN WOLFE

UNIVERSITY OF DUBLIN,
IRELAND

RICHARD YOUNG

MASSACHUSETTS INSTITUTE
OF TECHNOLOGY, USA

GENE REGULATION

New partner — new potential

The complex gene expression patterns that occur during development are often established through gene repression. The *Drosophila* Hairless protein, a bHLH-type transcription factor, silences genes by modifying their chromatin environment — it does so by interacting with its co-repressors Groucho and Rpd3, a histone deacetylase (HDAC). Rosenberg and Parkhurst have now identified a new HDAC partner for Hairless — a *Drosophila* homologue of yeast SIR2. They show that, in addition to its conserved association with heterochromatin, in flies Sir2 has a newly identified role — it helps to repress genes located in transcriptionally active chromatin (euchromatin), a function that is required for normal fly development.

Drosophila Sir2 had been identified by its sequence homology to the yeast protein by the Berkeley *Drosophila* Genome Project, and Rosenberg and Parkhurst sought to establish whether the two proteins shared the conserved role of SIR2 in silencing heterochromatin at different chromosomal locations. They found that in both an *in vitro* assay and *in vivo* — by monitoring the spreading of silencing to a marker gene placed near a heterochromatic region — fly Sir2 behaved as an HDAC. However, unlike its yeast counterpart, Sir2 is not involved in telomeric silencing.

The analysis of *Sir2* mutants made clear this gene's importance for fly embryogenesis. Segmentation is

disrupted in mutant embryos, and the abnormal expression of segmentation genes in *Sir2* mutants was reminiscent of the phenotype of weak loss-of-function *Hairless* mutations. Double-mutant analysis showed that Sir2 and Hairless interact genetically, and *in vitro* assays confirmed that Sir2 binds directly to Hairless's basic domain — the domain through which Hairless interacts with DNA. As the authors found that this interaction does not affect Hairless's ability to bind to DNA, they propose that Sir2 might instead affect Hairless's ability to recruit other factors or to alter the local chromatin state.

Interestingly, although this domain is highly conserved among HES family proteins, of which Hairless is a defining member, Sir2 binds to only one other HES protein, Deadpan, indicating that Sir2 might recognize sites other than the basic domain. The fact that Sir2 is required for embryonic patterning and that it interacts genetically and physically with Hairless suggests that, unlike its yeast homologue, which is exclusively associated with heterochromatic silencing, Sir2 is also involved in silencing genes located in euchromatin.

It remains to be confirmed whether Hairless's ability to recruit two different HDACs reflects its capability to independently regulate distinct developmental processes. The work of Rosenberg and Parkhurst poses many questions about the details of gene repression and the role that HDACs have in

this process. It does, however, show that euchromatic and heterochromatic silencing share common mechanisms.

Magdalena Skipper

References and links

ORIGINAL RESEARCH PAPER Rosenberg, M. I. & Parkhurst, S. M. *Drosophila* Sir2 is required for heterochromatic silencing and by euchromatic Hairless/E(Spl) bHLH repressors in segmentation and sex determination. *Cell* **109**, 447–458 (2002)