PNPLA3 AND LIVER DISEASE

A gene polymorphism could make children susceptible to liver damage from a young age, according to new research published in the *Journal of Hepatology*.

“Our research group had previously demonstrated, through the work of Stefano Romeo, that the *PNPLA3* gene is associated with increased hepatic fat content in genome-wide association studies in obese adult populations,” explains senior investigator Marco Baroni. “We decided to investigate if the genetic susceptibility for hepatic steatosis conferred by ... the I148M variant of *PNPLA3* could be present already in childhood or if it was exposed only in adults.”

The researchers genotyped 475 obese or overweight children and adolescents in Italy. They also recorded clinical and metabolic parameters (including indices of hepatic injury) for each individual and used ultrasonography to measure levels of steatosis in the liver.

Children with the 148M allele had increased levels of alanine transaminase (a 52% increase in individuals homozygous for 148M and a 9% increase in those heterozygous for 148M) than children with the wild-type 148I allele. Moreover, children who had two 148M alleles had higher levels of aspartate transaminase and steatosis than carriers of the wild-type alleles.

“Our data strongly suggest that overweight or obese carriers of the 148M allele experience chronic elevation of transaminase levels, starting during childhood and persisting in adult life,” adds Baroni. He proposes that the *PNPLA3* 148M allele increases the risk of hepatic injury from an early age, which could be further precipitated by ‘stressors’ such as obesity or excessive alcohol intake. The researchers will now investigate this ‘environmental influence’ on liver disease by determining the prevalence of the 148M allele in people affected by liver cirrhosis owing to alcohol abuse or viral infection.

Katrina Ray

Original article Romeo, S. et al. The 148M allele of the *PNPLA3* gene is associated with indices of liver damage early in life. *J. Hepatol.* 53, 335–338 (2010)

RESEARCH HIGHLIGHTS