NEONATAL IMMUNITY

Neonatal lymphocyte competence

Babies have a hard time of it — not only do they have a bit of a struggle coming into the world, but also they are then particularly susceptible to various infections with viruses and bacteria. It is generally believed that neonatal susceptibility to infection is due to a developmental defect in immune function, particularly a delay in the maturation of effective T helper 1 (T_H1)-cell responses. But, little is known about how neonatal CD8⁺ T cells respond to virus infection. Now, Marchant and colleagues show that neonates can generate a mature and functional CD8⁺ T-cell response to infection with human cytomegalovirus (HCMV), which might have implications for the development of neonatal vaccines.

Marchant and colleagues studied the functional capability of CD8⁺ T cells in congenitally infected neonates. Fetal T cells at the 28-week stage showed characteristics of early differentiated cells, with numerous highly activated cells undergoing proliferation. Infected newborns had higher proportions of activated CD8⁺ T cells than uninfected neonates, and repertoire analysis showed that HCMV infection had triggered the oligoclonal expansion of CD8⁺ T cells. Analysis of expression markers showed that the T cells of newborns had the characteristics of antigen-experienced cells with a phenotype resembling that of memory cells. Intracellular staining of these T cells showed high levels of perforin and

granzyme A, and the T cells had perforin-dependent cytolytic activity. The T cells were also able to produce antiviral cytokines, such as interferon- γ .

So, fetal and neonatal CD8⁺ T cells do seem to be able to respond to virus infections. These results indicate that there might be fewer developmental constraints on the CD8⁺ T-cell compartment than on CD4⁺ T cells, and although further work on the mechanisms of T-cell priming in neonates is necessary, it might be possible to develop neonatal vaccination for virus infections.

Elaine Bell

References and links

ORIGINAL RESEARCH PAPER Marchant, A. *et al.* Mature CD8⁺ T lymphocyte response to viral infection during fetal life. *J. Clin. Invest.* **111**, 1747–1755 (2003)

HIV

Under pressure

Mutation of HIV-1 in epitopes that are recognized by cytotoxic T lymphocytes (CTLs) is thought to be an important factor in the failure of CTLs to control infection. Yet, this phenomenon has not been consistently observed *in vivo*. Now, Otto Yang *et al.*, reporting in the *Journal of Experimental Medicine*, convincingly show that virus replicating *in vitro* under selective pressure can rapidly develop resistance to CTL-mediated antiviral activity as a result of genotypic changes.

Although analysis of virus sequences from infected individuals has previously shown that epitope-escape mutations can occur, it has remained controversial as to whether it is the immune pressure exerted by CTLs that leads to virus escape. The emergence of HIV-1 escape mutations depends on the balance of selective pressure — exerted by numerous factors, such as CTL frequency, epitope specificity, properties of the T-cell receptor and the functional state of the CTL — and the cost of the mutation to virus fitness.

The *in vitro* system reported here allowed the study of HIV-1 escape from CTL clones in a controlled manner, which cannot be readily achieved *in vivo*. The authors used three well-characterized CTL clones that are specific for an

epitope (SL9) in HIV-1-encoded structural protein Gag p17. However, these clones varied in their ability to recognize mutations in the SL9 epitope. Accordingly, when virus was co-cultured for two weeks with the SL9-specific and control clones, under conditions that favoured vigorous virus replication, a single point mutation in the SL9 epitope occurred only when it afforded resistance to suppression by the SL9-specific clones. The ability of the virus variant to arise in the face of immune pressure indicates that the mutation did not carry a significant fitness cost. Remarkably, although there are many potential point mutations in the epitope, as well as flanking sequences, that can prevent recognition for any given CTL clone, only one or no escape mutation arose under pressure of the CTL clones tested. By contrast, CTLs specific for HIV-1 protein Nef rapidly — after only 7 days of co-culture — selected for virus with polyclonal mutations in both the epitope and upstream sequences. Nef is dispensable for virus growth in this system, indicating that fitness constraints have a crucial role in determining escape.

So, CTLs can select for virus escape mutations, the extent of which depends on the fitness cost to the virus and the particular selecting CTL clone.

Lucy Bird

References and links

ORIGINAL RESEARCH PAPER Yang, O. O. *et al.* Determinants of HIV-1 mutational escape from cytotoxic T lymphocytes. *J. Exp. Med.* **10**, 1365–1375 (2003)