In the news

AGGRAVATING ASTHMA

New research, published in *The New England Journal of Medicine* (6 December 2007), has confirmed what many asthmatics have known for years — that the urban environment aggravates asthma.

In this study, 60 individuals with mild or moderate asthma took two-hour walks on Oxford Street, London, UK and in Hyde Park, London, UK, and their airway function was monitored before, during and after each walk. Only diesel-powered buses and taxis are allowed on Oxford Street, whereas Hyde Park is traffic free, providing the researchers with unique, 'real-life' locations to test whether traffic fumes worsen asthma symptoms.

"For the first time we are able to measure exactly what's happening inside the lungs of people with asthma when they spend only a couple of hours strolling in a real-life polluted area," said Professor Fan Chung, one of the authors of the study (*ScienceDaily*, 6 December 2007).

They found that diesel-exhaust fumes have a measurable effect on asthma symptoms, lung function and lung inflammation. Air-quality tests showed that there were three times as many ultra-fine particles in the air in Oxford Street compared with the air in Hyde Park, and the observed effects were associated with the higher concentrations of nitrogen dioxide and elemental carbon that were detected in the air in Oxford Street. "Our study illustrates the need to reduce pollution in order to protect people's health," said Fan Chung (*AFP*, 6 December 2007).

The assistant director of research at Asthma UK, Leanne Male, predicts that "Further research like this will provide the answers to help us improve the lives of the millions of people with asthma in the UK who are affected by traffic fumes every day." (*BBC News*, 6 December 2007.)

Olive Leavy