The relationship between a host and its intestinal microbiota may be even more complex and extensive than we previously anticipated, as suggested by a recent study in *Nature Medicine*. Clarke *et al.* show that peptidoglycan derived from the microbiota can leak out of the normal gut and enhance the function of neutrophils residing far away in the bone marrow.

Previous studies showed that recognition of peptidoglycan by the pattern recognition receptor NOD1 (nucleotide-binding oligomerization domain protein 1) enhanced the ability of neutrophils to kill *Streptococcus pneumoniae*. Clarke *et al.* now extend this finding by showing that products from the intestinal microbiota function on a systemic level to prime neutrophil function. Neutrophils isolated from the bone marrow of mice given broad-spectrum antibiotics to deplete the microbiota or raised in germ-free conditions were less efficient at killing *S. pneumoniae* and *Staphylococcus aureus* than those from mice with a normal microbiota. This effect was dependent on NOD1 but not NOD2 or Toll-like receptor 4.

After colonizing germ-free mice with *Escherichia coli* expressing a radioactively labelled peptidoglycan NOD1 ligand, the authors detected radioactivity in the serum, bone

marrow and neutrophils, which is consistent with the hypothesis that gut-derived peptidoglycan can enter the circulation and have far-reaching systemic effects. Indeed, sera from mice with a normal microbiota elicited greater NOD1 signalling in an *in vitro* reporter assay than sera from germ-free or antibiotic-treated mice.

Importantly, the authors showed that intraperitoneal administration of a NOD1 ligand is sufficient to restore neutrophil function and killing of bacterial pathogens in microbiota-deficient mice. Finally, the finding that *Nod1*^{-/-} mice were more susceptible to early pneumococcal sepsis than wild-type mice suggested that NOD1 signalling induced by microbiota-derived peptidoglycan primes neutrophils for rapid responses to subsequent infections.

So, these data provide a mechanism for the beneficial (or probiotic) activity of the microbiota on systemic innate immune function and point to possible adverse effects on health of extensive antibiotic use.

Lucy Bird

ORIGINAL RESEARCH PAPER Clarke, T. B. *et al.* Recognition of peptidoglycan from the microbiota by Nod1 enhances systemic innate immunity. *Nature Med.* **16**, 228–231 (2010)

FURTHER READING Philpott, D. J. & Girardin, S. E. Gut microbes extend reach to systemic innate immunity. *Nature Med.* **16**, 160–161 (2010)