High-dose intravenous immunoglobulin (IVIG) has been used for the treatment of autoantibody-mediated autoimmune diseases, but the molecular mechanism behind its anti-inflammatory effect was largely unknown. Previous studies have shown that sialylated Fc (sFc) fragments of IVIG (the active component of IVIG) bind to DC-specific ICAM3-grabbing non-integrin (DC-SIGN) on myeloid cells, thereby suppressing inflammation.

Ravetch and colleagues, writing in *Nature*, propose a detailed mechanism for the IVIG-mediated anti-inflammatory response. The authors generated transgenic mice expressing human DC-SIGN on dendritic cells, macrophages and monocytes, thus recapitulating the human expression pattern. DC-SIGN expression on mouse macrophages treated *in vitro* with IVIG or sFc was sufficient

to trigger their suppressive activity, as assessed following their adoptive transfer into mice with autoantibody-induced arthritis. Interestingly, the expression of both the inhibitory Fc receptor FcγRIIB and interleukin-4 (IL-4) by cells of the recipient mice was essential for this sFc–DC-SIGN-mediated suppression. Thus, the authors hypothesized that the binding of sFc to DC-SIGN may induce the IL-4-dependent upregulation of FcγRIIB on cells of the recipient mice. Indeed, administration of IL-4 had the same protective effect as IVIG in mice with arthritis but failed to suppress the disease in FcγRIIB-deficient mice.

Moreover, IVIG administration resulted in enhanced expression of IL-33 in wild-type mice but not in mice deficient for the mouse DC-SIGN orthologue SIGN-R1. Experiments using blocking

antibodies specific for the IL-33 receptor demonstrated that IL-33 is responsible for the sFc–DC-SIGN-induced anti-inflammatory response.

But, what is the link between the IL-33 response induced by the binding of sFc to DC-SIGN and the IL-4-mediated upregulation of FcγRIIB? Further analysis showed that IL-33 induces the expansion of IL-4-secreting basophil populations and, thus, increases the serum levels of IL-4. Adoptively transferred IL-33-treated basophils increased the levels of basophil-derived IL-4 in the recipient mice and protected these mice from arthritic inflammation. Protection was mediated by both wild-type and FcγRIIB-deficient basophils, and was associated with the induction of FcγRIIB expression on inflammatory macrophages.

So, taken together, these findings indicate that the anti-inflammatory effect of IVIG treatment is mediated by the upregulation of IL-33 production by myeloid cells, following the binding of sFc fragments to DC-SIGN. IL-33, in turn, expands IL-4-expressing basophil populations, and the resulting increase in IL-4 induces upregulation of FcγRIIB on the surface of inflammatory macrophages and suppression of autoantibody-mediated inflammation. The understanding of the IVIG-mediated anti-inflammatory mechanism reveals further targets for therapeutic intervention in autoimmunity.

Maria Papatriantafyllou

ORIGINAL RESEARCH PAPER Anthony, R. M. et al. Intravenous gammaglobulin suppresses inflammation through a novel T_H2 pathway. *Nature* 19 Jun 2011 (doi:10.1038/nature10134)

“the anti-inflammatory effect of IVIG treatment is mediated by the upregulation of IL-33 production by myeloid cells”