In the news

FIRST EARLY EMBRYOS IN VITRO

A team led by Magdalena Zernicka-Goetz at the University of Cambridge, UK, has reported in *Science* that it is possible to recreate the first stages of mouse embryogenesis *in vitro*, by combining embryonic stem (ES) cells and extra-embryonic trophoblast stem (TS) cells.

Recently, pluripotent ES cells were induced to form embryoid bodies and organoids, which are valuable tools for the study of development, as they can recapitulate several aspects of organ formation *in vitro*. However, these approaches are limited by their inability to mimic the spatiotemporal events of early embryogenesis.

In the current study, the authors cultured single ES cells with small clumps of TS cells (which normally give rise to the placenta) in a 3D matrix, and showed that the interaction between these two cell types — which involves the Nodal signalling pathway — enables them to self-assemble into structures that are very similar to natural mouse embryos. “We saw that the cells could self-organise themselves without our help,” says Zernicka-Goetz (*New Scientist*, 2 Mar 2017).

These findings open new avenues for the study of early development, although, whereas “the authors show mesoderm development, there is no data on the other two germ layers, ectoderm and endoderm,” comments Robin Lovell-Badge, a professor at The Francis Crick Institute, UK (*Science Media Centre*, 2 Mar 2017) — this warrants further investigation.

It is unclear whether these results can be replicated using human cells, especially as currently trophoblast cells cannot be isolated from human embryos. “We are very optimistic that this will allow us to study key events of this critical stage of human development without actually having to work on embryos,” comments Zernicka-Goetz (*The Guardian*, 2 Mar 2017).

These studies will undoubtedly stir both scientific and ethical debates.

Kim Baumann