HIGHLIGHTS

HIGHLIGHTS ADVISORS

JOAN S. BRUGGE

HARVARD MEDICAL SCHOOL,
BOSTON, MA, USA

PASCALE COSSART

INSTITUT PASTEUR, PARIS,
FRANCE

GIDEON DREYFUSS

UNIVERSITY OF PENNSYLVANIA,
PHILADELPHIA, PA, USA

PAMELA GANNON

CELL AND MOLECULAR
BIOLOGY ONLINE

JEAN GRUENBERG

UNIVERSITY OF GENEVA,
SWITZERLAND

ULRICH HARTL

MAX-PLANCK-INSTITUTE,
MARTINSRIED, GERMANY

NOBUTAKA HIROKAWA

UNIVERSITY OF TOKYO, JAPAN

STEPHEN P. JACKSON

WELLCOME/CRC INSTITUTE,
CAMBRIDGE, UK

VICKI LUNDBLAD

BAYLOR COLLEGE OF
MEDICINE, HOUSTON, TX, USA

TONY PAWSON

SAMUEL LUNENFELD RESEARCH
INSTITUTE, TORONTO, CANADA

NORBERT PERRIMON

HARVARD MEDICAL SCHOOL,
BOSTON, MA, USA

THOMAS D. POLLARD

THE SALK INSTITUTE,
LA JOLLA, CA, USA

JOHN C. REED

THE BURNHAM INSTITUTE,
LA JOLLA, CA, USA

KAREN VOUSDEN

NATIONAL CANCER INSTITUTE,
FREDERICK, MD, USA

JOHN WALKER

MRC DUNN HUMAN NUTRITION
UNIT, CAMBRIDGE, UK

GENE EXPRESSION

Be specific

How do you begin to comprehend the complex relationship between chromatin remodelling and transcription? Although many ATP-dependent chromatin-remodelling proteins have been identified, their function in transcription remains poorly defined due to the inherent difficulties in assaying this process. In a recent report in *Nature*, however, Tjian and colleagues describe the identification of a chromatin-remodelling protein complex that is specifically required for ligand-activated transcription from chromatin templates.

To search for transcription cofactors, the authors set up an *in vitro* transcription assay containing chromatin templates that were under the control of a nuclear hormone receptor. Starting with a set of purified proteins that are sufficient to drive transcription from 'naked' DNA, the authors asked which additional factors were required to drive transcription from a chromatin template.

They found an activity in human nuclear fractions that strongly potentiated ligand-dependent transcription from chromatin templates. Purification of this cofactor showed that it was a large multi-protein complex of the SWI/SNF family of chromatin-remodelling proteins, called SWI/SNF-B or polybromo- and BRG1-associated factor-containing complex (PBAF). The members of the SWI/SNF protein family use BRG1 or hBRM as their central ATPase, and this family is the largest

of the four families of chromatin-remodelling proteins.

SWI/SNF complexes share many common subunits, but BAF180, which contains six bromodomains, is unique to PBAF. The bromodomain motif has previously been shown to selectively bind to acetylated histone tails, and this binding could be important in targeting the cofactor complex to chromatin.

Tjian and colleagues wanted to know if other ATP-dependent chromatin-remodelling complexes could mimic the transcriptional activation activity shown by PBAF in the *in vitro* assay. They compared the effects of PBAF, SWI/SNF-A (which shares considerable similarity to PBAF) and ACF (a mechanistically different chromatin-remodelling protein complex) in their assay, and showed that although these proteins could all remodel the chromatin template, only PBAF efficiently activated transcription.

The authors also observed this effect using a class of transcription activators other than nuclear receptors, which showed that chromatin-remodelling proteins are not functionally interchangeable and that PBAF may have a function in transcription that is not limited to its chromatin-remodelling activity.

The finding that PBAF is required for nuclear-receptor-mediated transcription from chromatin templates is the first documented example of a specifically acting chromatin-remodelling protein complex. How PBAF

activates transcription remains unclear, but the *in vitro* assay used by Tjian and colleagues has provided a valuable insight into the functional specificity that can exist among the highly related complexes involved in gene regulation.

Rachel Smallridge

References and links

ORIGINAL RESEARCH PAPER Lemon, B. *et al.* Selectivity of chromatin-remodelling cofactors for ligand-activated transcription. *Nature* **414**, 924–928 (2001)

FURTHER READING Flaus, A. & Owen-Hughes, T. Mechanisms for ATP-dependent chromatin remodelling. *Curr. Opin. Genet. Dev.* **11**, 148–154 (2001) | Havas, K. *et al.* ATP-dependent chromatin remodeling activities. *Cell Mol. Life Sci.* **58**, 673–682 (2001)

WEB SITES

Encyclopedia of Life Sciences:

<http://www.els.net/>

Chromatin remodelling and histone modification in transcription regulation

Robert Tjian's laboratory:

<http://mcb.berkeley.edu/faculty/BMB/tjianr.html> |

<http://www.hhmi.org/research/investigators/tjian.html>