WNV entry to the brain. This was confirmed by comparing the permeability of the BBB in wild-type and *Tlr3*^{-/-} mice: following WNV infection or stimulation with the viral mimic poly (I:C), the permeability was increased in wild-type mice but not in *Tlr3*^{-/-} mice. A further insight into the pathogenesis of severe disease was provided by results suggesting that TNF- α receptor 1 signalling downstream of Tlr3 promotes WNV entry into the brain.

Sporadic outbreaks of WNV infection of humans have become increasingly common over the past 5 years, particularly in North America and Europe. This new work identifying Tlr3 as the receptor allowing WNV to enter the brain can hopefully be exploited for the development of new therapeutics.

Sheilagh Molloy

References and links

ORIGINAL RESEARCH PAPER Wang, T. *et al.* Toll-like receptor 3 mediates West Nile virus entry into the brain causing lethal encephalitis. *Nature Med.* 21 Nov 2004 (doi:10.1038/nm1140)

A female sandfly (*Phlebotomus* species), which is the vector of *Leishmania major*. The females are blood suckers and transmit parasites to humans. Image courtesy of the WHO © (1975).

PARASITOLOGY

Stuck in the gut

Cutaneous leishmaniasis, caused by infection with *Leishmania major*, is the most common Old World leishmanial disease and is transmitted by the sandfly. To devise new methods of parasite control it is imperative to understand the biology of infection in both the host and the vector. In a recent report in *Cell*, Kamhawi *et al.* have identified the receptor that binds to and retains *L. major* in the insect midgut, so that it can complete its replication cycle and produce a transmissible infectious form, a crucial step in the life cycle.

L. major replicates as intracellular amastigotes in macrophages, but replication in the sandfly occurs extracellularly in the digestive tract. Amastigotes transform into procyclic promastigotes, which are not infectious but become attached to the sandfly midgut and differentiate to produce infectious swimming metacyclic promastigotes, which are transmitted to humans through insect bites. Procyclic promastigotes must be retained in the gut to complete the replication cycle and produce infectious metacyclic promastigotes. Kamhawi *et al.* show that a tandem-repeat galectin, which is expressed on the surface of sandfly midgut epithelial cells, functions to specifically bind the main surface receptor of *L. major* procyclic promastigotes and attach them to the insect midgut.

Galectins are a β -galactoside-binding family of lectins conserved from fungi to man that contain one or more carbohydrate-recognition domains (CRD) and function in immunity, homeostasis and embryogenesis. The main glycoconjugate on the surface of *L. major* promastigotes is lipophosphoglycan (LPG), which has sugar side chains including a poly- β -galactosyl epitope (Gal β 1-3 $_n$) and can bind to galectin-3 and galectin-9. Galectin-9 is thought to function as a surrogate macrophage receptor in humans.

Kamhawi *et al.* sequenced a midgut cDNA library from the *L. major* vector *Phlebotomus*

papatasi and identified a galactose-binding protein named PpGalec. PpGalec is a tandem-repeat galectin with two CRDs separated by a linker and was specifically expressed in the midgut and upregulated in adult females. Moreover, PpGalec was only found in *P. papatasi* and its close relative *Phlebotomus dubosqi* (also an *L. major* vector) and was absent from other *Phlebotomus* species. Finally, only the procyclic (not the metacyclic) promastigotes bound to PpGalec.

Using confocal microscopy of insect midgut sections stained with PpGalec-specific antisera, expression of the galectin was localized to the luminal midgut cells. Importantly, pre-treatment of dissected midguts with PpGalec-specific antisera reduced the number of parasites bound by 72%. Plus, by adding PpGalec-specific antisera to a bloodmeal containing *L. major* amastigotes, parasite retention in the sandfly was markedly reduced, confirming that PpGalec has an integral role in parasite survival in the vector.

During differentiation from procyclic to metacyclic promastigote forms, downregulation of the poly-galactosyl epitopes on LPG disrupts interactions between LPG and PpGalec and promotes release of the infectious parasites. Interactions with galectins could account for vector specificity and parasite survival within the vector.

Identifying the insect receptor for *L. major* LPG is a huge advance that might facilitate development of new control methods, such as transmission blocking vaccines, for *L. major*-sandfly and other *Leishmania* species-sand fly interactions.

Susan Jones

References and links

ORIGINAL RESEARCH PAPER Kamhawi, S. *et al.* A role for insect galectins in parasite survival. *Cell* 119, 329–341 (2004)

WEB SITES

Jesus Valenzuela's laboratory:

<http://www.niaid.nih.gov/dir/labs/LMVR/valenzuela.htm>

David Sack's laboratory:

<http://www.niaid.nih.gov/dir/labs/lpd/sacks.htm>