Despite their different time courses, AAD and Wallerian degeneration might share similar molecular mechanisms. The authors found that in 'Wallerian-degeneration slow' mice, which show delayed Wallerian degeneration, AAD was largely absent. Moreover, calpain, which is known to participate in Wallerian degeneration, also seems to be a mediator of AAD, as is shown by the ability of calpain inhibitors to prevent it *in vivo*. Additional similarities and differences between AAD and Wallerian degeneration might now be established using this technique.

Beyond the description of the degeneration process, this method can also be used to monitor the efficacy of interventions aimed at preventing degeneration or promoting regeneration, as exemplified by the use of calpain inhibitors in this report. Moreover, as the response of axons to damage is relevant not only to spinal cord injury but also to conditions such as multiple sclerosis and amyotrophic lateral sclerosis, this technique might be used to provide new insights into these diseases.

Juan Carlos López, *Chief Editor*,
Nature Medicine

References and links

ORIGINAL RESEARCH PAPER Kerschensteiner, M. *et al.* *In vivo* imaging of axonal degeneration and regeneration in the injured spinal cord. *Nature Med.* 10 April 2005 (10.1038/nm1229)

TRPC channels. The authors propose a model whereby activation of TrkB by BDNF triggers, through PLC γ , the release of calcium from intracellular stores, which provides a sufficient increase in calcium concentrations to activate TRPC channels, allowing an influx of extracellular calcium, depolarization and turning.

TRPCs are part of a large family of TRP channels. TRPs are widely expressed, and have been implicated in many, especially sensory, functions. These two *Nature* reports indicate that TRPCs have an essential role in axon pathfinding. This exciting idea now awaits confirmation *in vivo*.

Annette Markus, *Associate Editor*,
Nature Neuroscience

References and links

ORIGINAL RESEARCH PAPERS

Li, Y. *et al.* Essential role of TRPC channels in the guidance of nerve growth cones by brain-derived neurotrophic factor. *Nature* 9 March 2005 (10.1038/nature03477) | Wang, G. X. & Poo, M.-m. Requirement of TRPC channels in netrin-1-induced chemotropic turning of nerve growth cones. *Nature* 9 March 2005 (10.1038/nature03478)

FURTHER READING Henley, J. & Poo, M.-m. Guiding neuronal growth cones using Ca²⁺ signals. *Trends Cell Biol.* 14, 320–330 (2004)

WEB SITES

Poo's laboratory: <http://mcb.berkeley.edu/labs/poo/>
Institute of Neuroscience, Shanghai: <http://www.ion.ac.cn/>

Image courtesy of Lawrence Keogh

DEVELOPMENT

NF- κ B branches out

The widely expressed transcription factor nuclear factor- κ B (NF- κ B) has a well-characterized role in many cellular processes, including inflammation, immune- and stress-related responses, and the regulation of cell survival and proliferation, in all tissues. Recent studies have also highlighted several neural-specific functions for NF- κ B, including an involvement in learning and memory, although the mechanisms that underlie these processes are not well understood. Now, however, Gutierrez and colleagues, reporting in *Development*, have uncovered a new and unexpected role for NF- κ B in promoting the growth and branching of neural processes in the nervous system during development.

To investigate the function of NF- κ B in the nervous system, Gutierrez and colleagues studied the activation of NF- κ B in two well-characterized populations of developing PNS and CNS neurons: the sensory neurons of the embryonic and neonatal nodose ganglion, and layer 2 pyramidal neurons in the postnatal somatosensory cortex of mice. In both populations, preventing NF- κ B activation or its transcriptional activity substantially reduced the overall span, total neurite length and total branch number of the neurite arbours. However, there was no effect on neuronal survival, which indicates that these changes were not the consequence of a general detrimental effect on neuronal viability, nor could they

be explained by a global reduction in cellular metabolism, as there was no effect on soma size.

Importantly, this effect of NF- κ B on the promotion of neurite growth was specific to a critical phase of neuronal development. In the nodose ganglion cells, it took place following the phase of naturally occurring neuronal death, when the axons of the remaining neurons elaborate and modify their terminal arborizations; and in the neurons of the somatosensory cortex it occurred when the dendritic arbours grow rapidly and establish functional connections.

It remains to be determined whether NF- κ B exerts its effects on neurite growth directly or through its action on the expression of other genes that are involved in the growth of neural processes. Nevertheless, this new role for NF- κ B sheds light on the molecular mechanisms that underlie the growth and branching of axonal and dendritic processes, which are crucial for functional development and plasticity in the nervous system. Moreover, this finding might explain the increasing evidence that NF- κ B is linked to learning and memory, as establishing and refining neuronal processes is thought to be important for these functions.

Alison Rowan

References and links

ORIGINAL RESEARCH PAPER Gutierrez, H., Hale, V. A., Dolcet, X. & Davies, A. NF- κ B signalling regulates the growth of neural processes in the developing PNS and CNS. *Development* 132, 1713–1726 (2005)