In the news

SILENCING SCRAPIE

Finding a method to target the abnormal proteins that cause prion diseases, such as scrapie, bovine spongiform encephalopathy (BSE) and Creutzfeldt–Jakob disease (CJD), could lead to new treatments or prevention strategies for these fatal diseases. A recent study suggests that harnessing RNA interference (RNAi) technology could be a promising approach.

Previous work has shown that genetic deletion of the normal form of PrP^C, the protein that becomes malformed to cause scrapie, can protect against disease development. Now, Alexander Pfeifer of the University of Bonn, Germany, and colleagues have shown that knocking out the protein in only a proportion of brain cells is effective, leading Pfeifer to suggest that this might be “a promising therapeutic option” (BBC News, 4 December 2006).

Despite these encouraging findings, the researchers urged caution. Pfeifer says: “It will take years before the method can be used on human beings” (University of Bonn, 4 December 2006). Approximately two-thirds of brain cells need to carry the transgene responsible for knocking out PrP^C to prevent disease progression. Targeting this number of brain cells in adult humans could be difficult, and improving current methods of gene delivery will be necessary. As Adriano Aguzzi from the University Hospital of Zurich, Switzerland, puts it, the researchers face “an engineering problem” (News @ nature.com, 1 December 2006).

Other researchers have highlighted the potential adverse effects of RNAi, including off-target effects and unknown consequences of deleting normal PrP^C. Qingzhong Kong from the Case Western Reserve University, USA, says that: “Much more research is needed” (BBC News, 4 December 2006) and argues that other therapeutic interventions, such as a vaccine against prion proteins, would be easier to administer.

Katherine Whalley

DEVELOPMENT

Dendrites hit the spot

During development of the nervous system, countless connections must be formed between neurons. How axons and dendrites come together to make the functional circuits of the brain is a subject of endless fascination. A new insight comes from Mumm *et al.*, who have found evidence that dendrites actively seek out specific targets in the developing brain.

Developing axons use various strategies to ensure that they reach the correct target, but dendrites have generally been thought to be more

“...retinal ganglion cells formed dendritic arborizations in a sequential pattern, targeting specific laminae...”

promiscuous. Instead of forming selective synapses, they are believed to generate widespread arborizations and connections that are then refined through removal, or ‘pruning’, of the inappropriate synapses and branches.

Mumm *et al.* used *in vivo* time-lapse imaging to investigate how the dendrites of retinal ganglion cells in the zebrafish form lamina-specific dendritic arborizations in the inner plexiform layer of the retina. Zebrafish embryos develop from an egg to a larva with a functioning visual system in just 5 days, and because they are transparent they are perfect for studying this kind of problem. By using transgenic zebrafish that expressed fluorescent dyes in specific neurons, the authors could watch as the dendritic arbor developed.

Remarkably, they found that many of the retinal ganglion cells formed dendritic arborizations in a

REPAIR

Rescue of vision

Degeneration of photoreceptors in the retina will cause irrevocable blindness and so far no treatment is available to rescue the sight of patients with this condition. Among the different treatment strategies, transplantation of retinal cells is closely pursued and investigated. In a recent study by MacLaren, Pearson *et al.*, the transplantation of photoreceptor precursor cells resulted in their functional integration into host retinas.

Previously, transplantation of either neural-derived stem cells or immature retinal cells into an adult retina has not been successful. It was believed that the mature environment prevents immature cells making appropriate connections and therefore functionally wiring up with the brain. By assessing the survival, differentiation and functional integration of photoreceptor precursor cells at different stages of differentiation, the authors have identified a time window in which these cells can be successfully transplanted and restore at least partial sight in mouse models of retinal degeneration.

Initially, the authors isolated immature mouse retinal cells on postnatal day 1 (P1), using green fluorescent protein (GFP)-expressing mice in order to discriminate between donor and host cells. Isolated P1 cells were transplanted into GFP-negative wild-type P1 littermates. The authors

sequential pattern, targeting specific laminae and working from the inner to the outer retina. This indicates that, rather than relying solely on dendritic pruning to establish accurate connections, some neurons can project dendrites towards specific targets.

were able to show that three weeks after transplantation P1 cells not only survived, but had also integrated into the recipient retina and had the morphology of mature photoreceptors. Strikingly, the same results were obtained when P1 cells were transplanted into adult retinas, indicating that the mature retina did not pose an obstacle to successful transplantation.

As P1 retinas are composed of a mixture of cells at different stages of differentiation, MacLaren, Pearson *et al.* set out to determine which cells of the isolated mixture become photoreceptors and at which stage of differentiation cells have the best potential to integrate. With this aim, retinal cells were isolated from animals at different developmental stages, namely from different embryonic stages through to adult, and transplanted into wild-type adult hosts. Only cells isolated between P1 and P7 showed robust integration. Specifically, the study showed that only postmitotic cells and not proliferating progenitor cells were able to integrate successfully into recipient retinas. When donor retinal cells were isolated from a transgenic mouse in which the expression of GFP was driven by the transcription factor *Nrl*, which is specifically expressed in postmitotic rod precursors, transplantation of only the *Nrl-Gfp*-positive cells yielded the

The authors also found that the retinal ganglion cells formed stratified arborizations in laminae in which amacrine cell neurites had already formed laminated plexuses. The neurites of amacrine cells might therefore 'prepattern' the inner plexiform layer and provide cues that guide the dendrites of ganglion cells to the correct laminae.

These findings add to the view of dendrites as complex and sophisticated structures that are gradually emerging from the shadow of the axon. Further work will lead to a better understanding of how dendrites can seek out appropriate targets during development.

Rachel Jones

ORIGINAL RESEARCH PAPER Mumm, J. S. *et al.*
In vivo imaging reveals dendritic targeting of laminated afferents by zebrafish retinal ganglion cells. *Neuron* **52**, 609–621 (2006)

same number of integrated cells as in experiments in which all isolated cells were transplanted. This provided the primary evidence that rod precursors differentiate into mature rod photoreceptors and functionally integrate into the recipient retina.

To assess this strategy as a viable therapeutic tool, the authors transplanted GFP-positive cells from *Nrl-Gfp* mice (rod precursors) into recipient retinas from three different mouse models of inherited retinal degeneration. Not only did the donor cells integrate and differentiate into rod photoreceptors, they were also able to restore light sensitivity in the recipient retinas, showing that they formed functional connections to the brain.

This study clearly shows that there is a very specific developmental time window during which photoreceptor precursor cells are able to integrate into the host retina and make functional connections to the brain. As success does not rely on the age of the recipient, this study is the first step towards a potential therapy for patients who lose their sight due to disease of the retina.

Claudia Wiedemann

ORIGINAL RESEARCH PAPER MacLaren, R. E., Pearson, R. A. *et al.* Retinal repair by transplantation of photoreceptor precursors. *Nature* **444**, 203–207 (2006)

IN BRIEF

➤ CALCIUM SIGNALLING

Regional interaction of endoplasmic reticulum Ca^{2+} signals between soma and dendrites through rapid luminal Ca^{2+} diffusion.

Choi, Y. M. *et al.* *J. Neurosci.* **26**, 12127–12136 (2006)

Although the role of the endoplasmic reticulum (ER) in Ca^{2+} signalling events is well established, the structural and functional organization of the neuronal ER is poorly understood. Choi *et al.* provide strong evidence that the ER in dendrites and the soma is interconnected. Using fluorescent Ca^{2+} markers, they measured changes in Ca^{2+} concentration in distinct regions of isolated dopamine neurons after local or whole cell stimulation. The results point towards the ER as a continuous, signal-integrating organelle in which Ca^{2+} can move rapidly between distinct parts of the neuron.

➤ REPAIR

The Nogo–Nogo receptor pathway limits a spectrum of adult CNS axonal growth.

Cafferty, W. J. & Strittmatter, S. M. *J. Neurosci.* **26**, 12142–12150 (2006)

Several factors expressed in the CNS inhibit axonal outgrowth; these include Nogo-A, which is mainly expressed by glial cells, and the axonal Nogo receptor NgR1. The authors assessed the regenerative capability of two strains of transgenic mice that failed to show an enhanced regenerative phenotype after spinal hemisection in previous studies: *nogo-a^{atg/atg}* and *ngr1^{-/-}*. This study used another lesion paradigm, unilateral pyramidotomy, and showed regeneration in both strains verified by pronounced collateral sprouting of intact corticospinal tract fibres across the midline into the denervated spinal cord. Identification of the molecular basis for this difference might reveal clues to efficient therapies after spinal cord injury.

➤ BEHAVIOUR

Male chimpanzees prefer mating with older females.

Muller, M. N. *et al.* *Curr. Biol.* **16**, 2234–2238 (2006)

Men's preference for younger women has been explained by the assumption that it makes more sense to form a long-term mating bond with a female who will be fertile for a long time to come. If this is true, male animals that do not form long-lasting partnerships, as is the case for chimpanzees, should not prefer younger over older females. The authors show that male chimpanzees in fact prefer older females over younger ones, although the reason for this particular preference remains elusive.

➤ CELLULAR NEUROSCIENCE

Distinct patterns of striatal medium spiny neuron activity during the natural sleep–wake cycle.

Mahon, S. *et al.* *J. Neurosci.* **26**, 12587–12595 (2006)

Medium spiny neurons (MSNs), the output cells of the striatum, have been characterized by their rhythmic transitions from hyperpolarized to depolarized activity states. Most studies revealing such MSN activity patterns have been performed in anaesthetized animals. Mahon *et al.* show that, in awake animals, MSNs display non-rhythmic membrane potential fluctuations, which result in irregular firing patterns. By contrast, during slow-wave sleep, their intracellular activity resembles that seen in anaesthetized animals.