DIABETIC NEPHROPATHY

CHOLESTEROL AND PODOCYTE INJURY

Researchers report that accumulation of cholesterol in podocytes has a central role in podocyte injury and might contribute to the development and/or progression of diabetic kidney disease. Sandra Merscher-Gomez and colleagues also found that induction of cholesterol efflux using cyclodextrin preserved human podocyte function *in vitro* and reduced albuminuria in a mouse model of progressive diabetic kidney disease.

To identify key pathways that are activated in podocytes in diabetic kidney disease, the researchers cultured normal human podocytes in the presence of sera from patients with type 1 diabetes mellitus with and without diabetic kidney disease and from healthy controls. They found that cell blebbing and intracellular cholesterol levels were increased and expression of the ATP-binding cassette transporter 1 (*ABCA1*)—a mediator of cholesterol efflux—was decreased in podocytes exposed to sera from patients with diabetic kidney disease compared with podocytes exposed to sera from diabetic patients with normoalbuminuria or from controls.

In further experiments, Merscher-Gomez *et al.* showed that cholesterol efflux was impaired in podocytes exposed to sera from patients with diabetic kidney disease and that *ABCA1* was downregulated in kidney biopsy samples from patients with diabetic kidney disease ($n=70$) compared with levels in kidney biopsy samples from healthy living kidney donors ($n=32$). Cyclodextrin treatment prevented cholesterol accumulation, cytoskeleton remodelling, cell blebbing and apoptosis in podocytes treated with sera from patients with diabetic kidney disease. In mice with diabetic kidney disease, cyclodextrin treatment also had beneficial effects, including a reduction in albuminuria and renal cortical cholesterol content and improvement in fasting insulin levels.

“Our findings indicate that cyclodextrin can protect podocytes in diabetic kidney disease,” concludes Merscher-Gomez.

“Future research will focus on moving the use of cyclodextrin for the treatment of diabetic kidney disease into clinical practice, and on the identification of new compounds that target reverse cholesterol transport for the treatment of diabetic kidney disease.”

Ellen F. Carney

Original article Merscher-Gomez, S. *et al.* Cyclodextrin protects podocytes in diabetic kidney disease. *Diabetes* doi:10.2337/db13-0399