Researchers have identified a number of single nucleotide polymorphisms (SNPs) that could be used to predict risks of recurrence and survival for patients with bladder cancer. “Frequent, painful screenings are needed to identify recurrences, so the ability to identify patients at high risk could help to improve quality of life,” explains Angeline Andrew of Norris Cotton Cancer Center, NH, USA, lead author of the study.

Andrew and colleagues hypothesized that bladder cancer prognosis might be affected by inherited variations in genes that are involved in carcinogenic processes, such as apoptosis, proliferation and DNA repair. To test their theory, they genotyped DNA samples from 563 patients diagnosed with bladder cancer between 1994 and 2001, identified in the New Hampshire State Cancer Registry. They reviewed medical records to obtain detailed clinical information, including recurrences, progression, tumour size, multiplicity and treatment.

Half of the patients in this cohort experienced recurrence at least once during the study period, and 40% died. Two SNPs were associated with a shorter time to first bladder cancer recurrence: rs2238151 in the aldehyde dehydrogenase 2 (*ALDH2*) gene (hazard ratio [HR] 2.03) and rs5742714 in the insulin-like growth factor 1 gene (HR 1.61). *ALDH2* is an enzyme involved in alcohol metabolism and, interestingly, patients with the variant SNP who drank alcohol were less likely to experience recurrence than those who never drank (HR 0.11).

Other SNPs were found to be predictive of survival. Patients with non-muscle-invasive bladder cancer who were heterozygous for a SNP in the gene for DNA repair protein *XRCC4* experienced longer survival (HR 0.52) than those with the wild type allele, whereas a polymorphism in tumour suppressor *RB1CC1* was associated with shorter survival than wild type for patients with non-muscle-invasive or muscle-invasive disease (HR 2.29 overall).

SNP–SNP and SNP–treatment interactions were also detected. Patients with polymorphisms in two immune-response-related genes (*IL15RA* and *MBL2*) were approximately four times more likely to experience recurrence than patients with neither. Andrew and her team hypothesized that patients with variant alleles of genes related to the immune response would exhibit a differential response to immunotherapy treatment. Indeed, time to recurrence was shorter for patients treated with immunotherapy who harboured SNP rs3176879 in the gene for vascular cellular adhesion molecule 1.

This population-based study has generated a large amount of data on the associations between SNPs and bladder cancer prognosis, which the authors hope will be replicated in other studies and investigated further. “Future work will include identifying factors that are modifiable after diagnosis, which could prevent recurrences in those patients at higher genetic risk,” concludes Andrew.

Sarah Payton

Original article Andrew, A. S. et al. Genetic polymorphisms modify bladder cancer recurrence and survival in a USA population-based prognostic study. *BJU Int.* doi:10.1111/bju.12641