BLADDER CANCER PATIENT-DERIVED MURINE XENOGRAPHS

Patient-derived xenograft (PDX) platforms could help to develop and personalize bladder cancer treatments, according to a recent article published in *PLoS ONE*.

Pan *et al.* collected tumour specimens from patients with bladder cancer, which were then implanted into immunodeficient NOD Scid gamma (NSG) mice, establishing an orthotopic PDX model by subsequent injection of a single-cell suspension of implanted PDXs into the mouse bladder. Overall, 22 PDXs were generated, which demonstrated morphology identical to the source tumour and 92–97% homology in terms of genetic aberrations on whole-exome sequencing. Mutation analysis was carried out to identify potential druggable targets.

Mice bearing PDXs were treated using molecularly targeted therapies (MTT) matched to selected genetic aberrations identified: *FGFR3*, *EphB4*, *Src*, *HER2/3* and *PIK3CA*. Next, the team identified drugs that could be repurposed to target the products of the specific genes, for example, the dual EGFR/HER2 inhibitor lapatinib, which is approved for breast cancer therapy. They noted discrepancies in response in groups of NSG mice bearing different PDXs that both expressed HER2, and further use of the PDX platform to screen for effective MTTs matching other cancer-specific genetic aberrations showed that different PDX responses were also observed with drugs targeting other mutations.

The PDX platform was also used to screen for effectiveness of the commonly used first-line chemotherapy drugs cisplatin, gemcitabine and both drugs in combination. Although some of the PDXs screened were resistant to cisplatin and some to gemcitabine, PDXs were more sensitive to the combination treatment and, in many cases, resistance to one drug could be overcome by the other.

The team are now testing this method in patients. “A patient has been treated with three lines of therapy ... with treatment decisions based on the efficacy studies in patient-derived xenografts,” explains lead author Chong-xian Pan. “The cancer response to the three lines of therapy between the patient’s cancer and her derived xenografts were exactly the same.” The team are now contacting pharmaceutical companies to discuss potential clinical trials based on these xenografts.

Annette Fenner

Original article Pan, C.-x. *et al.* Development and characterization of bladder cancer patient-derived xenografts for molecularly guided targeted therapy. *PLoS ONE* 10, e0134346 (2015)