Phosphoglycerate dehydrogenase (PHGDH) is a key regulator of the hypoxia-inducible factor (HIF)2 α pathway in renal tumours and might provide a treatment target for patients resistant to HIF2 α antagonists, according to a paper published in *Cancer Research*.

HIF affects metabolism in renal cell carcinoma (RCC), accelerating glycolysis and, therefore, disease progression, as RCC is dependent on aerobic glycolysis for ATP production. HIF2 α antagonists have been shown to increase progression-free survival (PFS) in patients previously treated with tyrosine kinase inhibitors. However, patients can develop resistance to these agents and an alternative approach to targeting this pathway could, therefore, be clinically useful.

Yoshino and colleagues established sunitinib-resistant clear cell (cc)RCC cells by injecting 786-o cells into mice and treating them with sunitinib until resistance was acquired. The cells were then harvested and HIF2 expression confirmed in those which would be used in further experiments (termed SU-R-786-o). They then used CRISPR–Cas9 to knock out HIF2 α in SU-R-786-o cells. Known HIF target genes were downregulated in the knockout cells and, furthermore, most genes previously shown to be upregulated in wild-type 786-o cells after HIF2 α antagonist treatment were also upregulated in the knockout cells without treatment, suggesting that the knockout cells mimicked treatment with the antagonist. Knockout cells were also morphologically different from their wild-type counterparts, with a rounder shape than the normal spindle shape of the cell line.

Combined RNA sequencing and proteomics studies showed that PHGDH was upregulated in knockout cells. Pathway analysis identified the serine biosynthesis pathways as being enriched. When PHGDH expression was suppressed in the HIF2 α -knockout SU-R-786-o cells using transfection with si-PHGDH, cell proliferation was inhibited. Two novel PHGDH inhibitors — CBR-5884 and NCT-503 — were used in cell proliferation assays and showed potent inhibitory effects on the knockout cells. Furthermore, CBR-5884 also induced apoptosis in the knockout cells and in a number of other ccRCC cell lines. NCT-503 treatment *in vivo* significantly suppressed tumour growth compared with vehicle treatment, which was likely to be due to reduced cell proliferation, as shown using Ki67 immunohistochemistry.

Overexpression of PHGDH in wild-type 786-o cells also affected cell morphology in a dose-dependent manner; those cells which demonstrated only weak overexpression did not change shape, whereas strong overexpression led to a round cell shape.

Analysis of The Cancer Genome Atlas data showed that patients with PHGDH amplification had worse overall and disease-free survival than those without ($n = 500$; $P < 0.0003$ and $P < 0.0329$, respectively.) In addition, PHGDH expression was a predictor of overall survival.

Normal serine biosynthesis was altered in SU-R-786-o and HIF2 α -knockout 786-o cells, with accumulation of 3-phosphoserine, serine, and glycine observed. The tricarboxylic acid cycle was also activated in these cells as well as the sunitinib-resistant cells compared with wild-type cells.

The data from these studies suggest that resistance to previously developed HIF2 α inhibitors could be overcome by targeting at a different point in the pathway. PHGDH could also be a relevant treatment target in patients with PHGDH gene amplification.

Annette Fenner

ORIGINAL ARTICLE Yoshino, H. et al. 1 PHGDH as a key enzyme for serine biosynthesis in HIF2 α -targeting therapy for renal cell carcinoma. *Cancer Res.* <http://dx.doi.org/10.1158/0008-5472.CAN-17-1589> (2017)