CORRIGENDUM

ATM-mediated KDM2A phosphorylation is required for the DNA damage repair

L-L Cao, F Wei, Y Du, B Song, D Wang, C Shen, X Lu, Z Cao, Q Yang, Y Gao, L Wang, Y Zhao, H Wang, Y Yang and W-G Zhu

Oncogene (2016) 35, 402; doi:10.1038/onc.2015.311**Correction to:** *Oncogene* (2016) 35, 301–313; doi: 10.1038/onc.2015.81; published online 30 March 2015

Since the publication of the above article the authors have requested a change to the author affiliations as per below:

L-L Cao^{1,4}, F Wei^{1,4}, Y Du¹, B Song¹, D Wang¹, C Shen¹, X Lu¹, Z Cao¹, Q Yang¹, Y Gao², L Wang¹, Y Zhao¹, H Wang¹, Y Yang¹ and W-G Zhu^{1,3}¹Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education), State Key Laboratory of Natural and Biomimetic Drugs, Beijing Key Laboratory of Protein Posttranslational Modifications and Cell Function, Department of

Biochemistry and Molecular Biology, School of Basic Medical Sciences, Peking University Health Science Center, Beijing, China;

²Protein Chemistry Facility, School of Biological Sciences, Tsinghua University, Beijing, China and³Peking University-Tsinghua University Joint Center for Life Sciences, Beijing, China. Correspondence: Professor W-G Zhu, Department of Biochemistry and Molecular Biology, Peking University Health Science Center, Xueyuan Road 38#, Beijing 100191, China. E-mail: zhuweiguo@bjmu.edu.cn⁴These authors contributed equally to this work.

The authors wish to apologise for any inconvenience caused.