

ERRATUM

A new role of SNAI2 in postlactational involution of the mammary gland links it to luminal breast cancer development

S Castillo-Lluva, L Hontecillas-Prieto, A Blanco-Gómez, M del Mar Sáez-Freire, B García-Cenador, J García-Criado, M Pérez-Andrés, A Orfao, M Cañamero, JH Mao, T Gridley, A Castellanos-Martín and J Pérez-Losada

Oncogene (2015) **34**, 4797–4798; doi:10.1038/onc.2015.322

Correction to: *Oncogene* (2015) **34**, 4777–4790; doi:10.1038/onc.2015.224; published online 22 June 2015

Following the online publication of this article we were notified that the headings Snai2 WT^{ERBB2+} and Snai2 KO^{ERBB2+} in Table 1 had been inserted in the incorrect columns. The corrected table appears overleaf.

We apologise for any inconvenience caused by this error.

This error has now been rectified, and the corrected article appears in this issue. The html and online pdf versions have also been rectified, and now carry the corrected paper.

Table 1. Pathophenotypes studied in nulliparous and parous *Snai2* WT^{ERBB2+} and *Snai2* KO^{ERBB2+} female mice

		<i>Snai2</i> WT ^{ERBB2+}	<i>Snai2</i> KO ^{ERBB2+}	P-value
A. Temporal stages				
<i>Tumor latency (weeks)</i>				
Latency interval (n)	Nulliparous Parous	28.8–90.9 (n = 41) 21.6–61 (n = 24)	29.7–114 (n = 33) 13.1–80.8 (n = 45)	
Median (IR)	Nulliparous Parous	47.7 (13.8) 40 (14.82)	54.1 (21.04) 40.4 (13.21)	0.0127 NS
	P-value	0.0038	0.0004	
<i>Duration of disease (weeks)</i>				
Disease interval (n)	Nulliparous Parous	4.28–23.86 (n = 36) 6.7–20 (n = 24)	4.57–21.14 (n = 22) 5.14–22.29 (n = 45)	
Median (IR)	Nulliparous Parous	12.9 (12.75) 13.3 (7.46)	10.36 (10.39) 11.6 (5.14)	NS NS
	P-value	NS	NS	
<i>Lifespan (weeks)</i>				
Lifespan interval (n)	Nulliparous Parous	39.4–90.9 (n = 41) 20–79.1 (n = 24)	38.3–114 (n = 32) 32.1–93 (n = 45)	
Median (IR)	Nulliparous Parous	60.42 (14.28) 53.38 (17.07)	66.14 (32.93) 50 (16.14)	0.0145 NS
	P-value	0.0089	0.0009	
B. Tumor traits				
<i>Tumor number</i>				
Incidence percentage (rate)	Nulliparous Parous	87.8% (36/41) 87.5% (21/24)	66.67% (22/33) 76.09% (35/46)	0.0453 NS
	P-value	NS	NS	
Multiplicity (2 or more)	Nulliparous Parous	88.89% (32/36) 90.48% (19/21)	59.09% (13/22) 94.29% (33/35)	0.0204 NS
	P-value	NS	0.0017	
Median (IR)	Nulliparous Parous	3 (2) 3 (4)	1 (3) 3 (3.25)	0.0012 NS
	P-value	NS	0.0076	
<i>Local growth rate</i>				
Median (IR; n)	Nulliparous Parous	0.32 (0.17; n = 59) 0.24 (0.21; n = 46)	0.33 (0.136; n = 26) 0.32 (0.27; n = 72)	NS NS
	P-value	NS	NS	
<i>Metastasis</i>				
Incidence % (rate)	Nulliparous Parous	63.64% (21/33) 90.47% (19/21)	50% (9/18) 61% (20/33)	NS 0.0276
	P-value	0.0535	NS	
Multiplicity (2 or more)	Nulliparous Parous	100% (21/21) 78.9% (15/19)	100% (9/9) 90% (18/20)	NS NS
	P-value	0.0424	NS	
Median (IR)	Nulliparous Parous	5 (8.5) 8 (18)	1 (5.25) 3 (10)	NS 0.0349
	P-value	NS	NS	

Abbreviations: IR, interquartile range; n, number of mice; NS, non-significant *P* values. Tumor traits were compared with the Mann–Whitney *U*-test, and the temporal stages of the disease were evaluated using the Kaplan–Meier estimator and the Log-Rank test. The incidence and multiplicity of tumors and metastases were compared with Fisher's exact test. The numbers of mice used in the experiments are shown in the respective figures along the manuscript. The *n* values shown in the local growth rate represent all tumors analyzed.