Dear Editor

Pragmatic randomised controlled trial

I welcome the article by Dr Kernick and colleagues in the journal (*Prim Care Respir* 2002 **1** (1) 6-8) which is an admirable attempt to evaluate the as yet unproven worth of asthma clinics in primary care. The design being pragmatic is appropriate but a number of points need to be made in my view

Firstly, just because a study is "real-life" or pragmatic there is no excuse for reduced research standards. Whilst blinding of patients to interventions may not be feasible wherever possible there should be as much blinding and rigor as possible. For instance, data collection should be undertaken by someone blinded to treatment allocation or questionnaires should be completed away from the study setting. Furthermore, extending the length of these studies to 6 months or even a year would permit the "halo" effects of being an intervention group to wear off.

Secondly, whilst admiring the value of looking at the proportion of responders, (a factor often missed in clinical trials where mean data only is reported), it is important to remember random effects. Guyatt¹ has suggested a simple methodology for taking this into account looking at deteriorators and improvers. In an intervention without bias and no clinical effect these numbers should be equal whereas an effective intervention should both increase improvers and decrease deteriorators. The author should therefore report the proportion deteriorating by clinically significant amounts as well. It is likely that the trend found towards benefit in the intervention group may have been statistically significant if evaluated in this way.

David Price, GPIAG Chair in Primary Care Respiratory Medicine

1. Guyatt GH, Juniper EF, Walter SD, *et al* Interpreting treatment effects in randomised trials. *BMJ* 1998 Feb **3** 3316 (7132): 690-9

Sir

Revision of BTS guidelines for the treatment of Asthma

I am writing in response to your letter in the December issue of the GPIAG journal inviting comments on the BTS Guidelines. I also noted the disappointing results of asthma management in the UK mentioned by Mike Thomas in the March issue of the journal. There are, of course, numerous audits and surveys which support this

At present the BTS guidelines recommend the use of short acting β_2 agonists as step 1 treatment and by implication the first drug to use in newly diagnosed asthmatics. These bronchodilator drugs give temporary relief but the symptoms soon return and may even worsen, requiring even more β_2 agonist inhalation but never really achieving proper control

This clearly is not the intention of the guidelines which state that the drug combination or "step" should be appropriate to the severity of the disease. Unfortunately, doctors do not always assess severity accurately and are often reluctant to prescribe inhaled steroids. Patients are also reluctant to take them as they do not give immediate relief. Thus many patients struggle to control their asthma without anti-inflammatory treatment, never reaching their optimum lung function, continuing with chronic disability and risking long-term lung damage leading to chronic obstructive disease

Author's Reply

Dear Editor

The question from Professor Price is an important one - "just because a study is real life or pragmatic, is it an excuse to reduce research standards?"

This comment opens up a much broader agenda and reflects a paradigm that assumes there is an absolute truth "out there" that can be obtained by the application of rigorous scientific endeavour. I would argue that solutions are always satisfactory, never optimal and that we live in an environment of "bounded rationality" where we have to do the best we can against a background of limited resources and conflicting objectives. It seems that the burgeoning academic literature has little relevance or impact on decision-makers as they muddle through in an environment of conflicting objectives, ambiguity and limited room for manoeuvre. We need more "quick and dirty" research that at least helps to make sense of the world rather than rigorous studies that are invariably out of date and bear little relationship to the contingencies of the real world

David Kernick

Another problem resulting from the first use of β_2 agonists is that the patient thinks that this is the proper treatment for asthma and that inhaled steroids are an optional extra. If steroids were given to newly diagnosed asthmatics they may have to wait a few days to obtain symptomatic benefit but they would soon regard these drugs as the proper treatment for asthma. Furthermore, they would understand the importance of taking emergency oral steroids early in an attack.

The BTS classification of severity of asthma in "steps" by the amount of treatment prescribed is unhelpful. It would be better to adopt the American system

Mild intermittent asthma
Mild persistent asthma
Moderate persistent asthma
Severe asthma

A completely fresh approach to the treatment of asthma is therefore required with steroids being used as soon as the diagnosis has been confirmed. A large dose should be used initially and then gradually reduced to a level where the minimum dose is used to control symptoms. β_2 agonists are used as needed for breakthrough wheezing. Exacerbations are treated by increasing the dose of steroids and adding bronchodilators if necessary.

Those subject to attacks should carry their own supply of oral steroids and should not hesitate to take them in time to stop an attack

This approach has been used successfully in Finland where reduction of 74% of hospital admissions has been achieved (see Thorax 2001; 56:806-814).

Some GPs (and nurses) in the UK follow this method but they are swimming against the tide of popular opinion generated by the BTS guidelines. Perhaps they should receive some encouragement from the GPIAG

Dr George Strub