References

- [1] Asthma – the Australian Government perspective. Accessed on 15 January 2006 at www.health.gov.au/internet/wcms/publishing.nsf/Content/health-pq-asthma-index.htm.
- [2] National Asthma Action Plan 1999–2002 Australian Government 2001. Accessed on 15 January 2006 at [www.health.gov.au/internet/wcms/Publishing.nsf/Content/health-pq-asthma-pubs-naap.htm/\\$FILE/naap.pdf](http://www.health.gov.au/internet/wcms/Publishing.nsf/Content/health-pq-asthma-pubs-naap.htm/$FILE/naap.pdf).
- [3] The National Asthma Council Australia. See strategic documents and other resources at www.nationalasthma.org.au/html/strategy/home/index.asp accessed on 15 January 2006.
- [4] The Australian Asthma Monitoring Centre. See the latest report Asthma in Australia 2005 accessed on 15 January 2006 at www.asthamonitoring.org/.

doi:10.1016/j.pcrj.2006.04.118

ABS17: Linking evidence to policy – lessons from the national evaluation of the Asthma 3+ Visit Plan in Australia

Justin Beilby^a, Nicholas Zwar^b, Teresa Burgess^a, Iqbal Hasan^b, Gawaine Powell Davies^b, Sophie Couzos^c, Scott Davis^c

^a University of Adelaide, North Terrace, Adelaide, 5005 Australia ^b University of NSW, Australia ^c NACCHO, Australia

Introduction: The Asthma 3+ Visit Plan (rewarding systematic asthma care in primary care) is unique to Australia. Based on the Six Step Asthma Plan, it comprises three GP visits over four months for patients with moderate to severe asthma. GPs receive an incentive payment on completion of the three visits. **Objectives:** The Aim was to establish the degree of uptake of the Plan. **Objectives** included identifying barriers and enhancers to uptake for consumers and GPs. **Subjects:** GPs, GP organisations and consumers (including non English speaking and Aboriginal and Torres Strait Islander peoples).

Methods:

- Statistical analysis of Medicare GP Payment data
- A semi structured phone interview of all 120 National GP Organisations
- A GP Survey (of a stratified sample of GPs)
- Consumer interviews
- Focus groups with non English speaking Australians
- Aboriginal and Torres Strait Islanders uptake analysis

Data from all these elements were analysed and triangulated and this paper reports on this extensive analysis. **Results:** Uptake remains low and has not increased notably since a peak six months after introduction in 2001. Key barriers identified were:

- Structure of the incentive
- Lack of consumer education
- General Practice systems

The current Plan is not appropriate for Aboriginal and Torres Strait Islander or CALD populations. **Conclusion:** A 'one size fits all' policy does not meet the range of needs across the Australian community. The value of rewarding systematic asthma care in primary care was however demonstrated. Flexibility, multi-disciplinary participation and a recognition of differing practice circumstances is required and efficient general practice systems are demonstrated as the key to systematic chronic disease interventions. Consumer education is needed around preventive health care.

Conflict of interest and funding

Funding: Australian Government Department of Health and Ageing. No conflict of interest declared.

doi:10.1016/j.pcrj.2006.04.119

ABS18: Role of hypertonic saline in bronchoprovocation for the diagnosis of bronchial asthma

S.M. Abdullah, Al Manun, Mahiuddin Ahmed, Mustafizur Rahman
NIDCH, Mokhali, Dhaka, 1216, Bangladesh

Introduction: Bronchoprovocation is performed to evaluate bronchial hyper-responsiveness. Almost all asthma patients demonstrate bronchial hyper-responsiveness. **Materials and method:** 50 non-smoking adults with complaints of breathlessness, cough, chest tightness & positive family history of bronchial asthma, baseline FEV₁ > 70% of predicted (mean FEV₁ = 84, median value = 81.7) were included in this study. Patients underwent bronchoprovocation using methacholine followed by hypertonic saline, 7 days apart or vice versa. **Results:** Of 50 patients, 34 were male, 16 were female with a mean age of 26.44 years. With 1 mg/ml of methacholine fall of FEV₁ > 20%, 10–20%, <10% was observed in 36 (72%), 8 (16%) & 6 (12%) patients respectively. With 14% hypertonic saline >20% fall was observed in 32 (64%), >10–20% in 4 (8%), 5–10% in 2 (4%) & <5% in 2 (4%) patients. No response was observed in 10 patients. **Conclusion:** This study shows that hypertonic saline is an equally effective but more specific bronchoprovocative agent than methacholine in bronchial asthma. Since hypertonic saline is very cheap and easily available, we propose that it should be considered as a reliable bronchoprovocative agent for the diagnosis of bronchial asthma in poor countries.

Conflict of interest and funding

None declared.

doi:10.1016/j.pcrj.2006.04.120

ABS19: Parents' perceptions of asthma in children aged 2–5 years

Margaret Cuffwright

Salford Primary Care Trust, Sandringham House-Salford Primary Care Trust, Windsor Street, Salford, Manchester, M5 4DG, United Kingdom

Introduction: There is reluctance on the part of clinicians to diagnosis asthma in young children consequently parents of children with asthma express anxieties about how to manage their child's symptoms. **Objective:** To explore the significance of a diagnosis on parents. **Methods:** The purposive sample consisted of parents of 28 children with newly diagnosed asthma aged 2–5 years. Data were analysed using the principles of grounded theory. **Results:** During the pre-diagnosis phase parents learned that asthma is variable. Conflicting advice from professionals influencing their view about preventative treatment. Parents had few concerns about steroids although this view changed by the 2nd interview. **Conclusion:** Parents views of asthma were shaped by the events leading up to the diagnosis and these influenced how they managed the condition now and in the future.

Conflict of interest and funding

Funded by North West Office of the Department of Health.

doi:10.1016/j.pcrj.2006.04.121

ABS20: Developing a community based spirometry service for east London

Jill Goddard

Tower Hamlets PCT, City University, Bancroft Road, London, E1 4DG, United Kingdom

Introduction: Spirometry is the gold standard for the diagnosis and classification of severity in chronic obstructive pulmonary disease (COPD). It also provides a valuable tool