

and scattering at about  $3,350 \text{ cm}^{-1}$ . The coupling constants range from the order of magnitude of solid *n*-decanol for methanol and ethanol to about 50% larger for the higher members of the series.

These observations provide a direct spectroscopic argument for the "polymeric" structure of liquid alcohols and suggest grounds for reconsidering earlier studies which failed to account for the possibility of multiple -OH stretching modes. The unexplained secondary splittings and the effect of increasing molecular weight on the spectra suggest areas of further investigation and raise the possibility that Raman spectroscopy may soon become a more powerful and widely applied tool for the study of hydrogen-bonded liquids than infrared spectroscopy was two decades ago.

H. A. WELLS\*

Air Force Materials Laboratory,  
Wright-Patterson Air Force Base,  
Ohio 45433

Received August 17, 1972.

\* Present address: Institute of Paper Chemistry, Appleton, Wisconsin, USA.

- <sup>1</sup> Pimentel, G. C., and McClellan, A. L., *The Hydrogen Bond*, 75 (Freeman, San Francisco and London, 1960).
- <sup>2</sup> Jakobsen, R. J., Mikawa, Y., and Brash, J. W., *Nature*, **215**, 1071 (1967).
- <sup>3</sup> Sheppard, N., in *Hydrogen Bonding* (edit. by Hadzi, D.), 85 (Pergamon, London, 1959).
- <sup>4</sup> Sandorfy, C., *Can. J. Spectrosc.*, **17**, 1 (1972).
- <sup>5</sup> Luck, W. A. P., and Ditter, W., *Ber. Bunsenges. phys. Chem.*, **72**, 365 (1968).
- <sup>6</sup> Venkateswarlu, K., and Mariam, S., *Acta phys. austriaca*, **15**, 367 (1962).
- <sup>7</sup> Wagner, J., *Z. phys. Chem.*, **45**, 69 (1940).

## Can Spin Avert Singularities?

TRAUTMAN has suggested<sup>1</sup> that the introduction of intrinsic spin effects into general relativity through the Einstein-Cartan torsion theory may avert a gravitational singularity. In this note we suggest that the singularity is avoided only because of the high symmetry of the model used.

We shall not describe the Einstein-Cartan theory here (see ref. 1 for a bibliography), except to point out that for a classical description of spin, with spin density tensor

$$s^a{}_{bc} = \eta_{bcde} u^d u^e s^a \quad (1)$$

the usual relativistic description of the fluid flow<sup>2</sup> is valid. (In equation 1,  $u^a$  is the 4-velocity of the spinning matter, and  $s^a$  is the spacelike spin vector of the fluid satisfying  $s^a u_a = 0$ .) But one modification has to be made. The usual vorticity vector  $\omega^a = \frac{1}{2} \eta^{abcd} u_c u_d u_b$  which gives the angular velocity of the fluid with respect to a Fermi-propagated frame becomes  $\omega^a + s^a$ . Hence the Raychaudhuri equation<sup>2</sup> takes the form

$$\theta + \frac{1}{3} \theta^2 + 2[\sigma^2 - (\omega_a + s_a)(\omega^a + s^a)] + \rho = 0 \quad (2)$$

for dust of density  $\rho$ , where  $\theta = u^a{}_{;a}$  is the expansion of the fluid, and  $\sigma$  is the shear scalar. In Trautman's quasi-Friedmann model,  $\sigma = \omega^a = 0$ , and conservation of matter and spin implies  $\rho = \rho_0 R^{-3}$ ,  $s = s_0 R^{-3}$ , where  $R(t)$  is the usual Robertson-Walker scale length, and  $s = (s^a s_a)^{1/2}$ . It follows that for small  $R$ , the repulsive spin term  $-2s^2 \sim R^{-6}$  in equation 2 dominates the attractive  $\rho \sim R^{-3}$  term, and the singularity is avoided. If a perfect radiation fluid is considered,  $\rho \sim R^{-4}$ , which does not alter the conclusion.

But Trautman<sup>1</sup> invokes a primordial magnetic field to produce global spin alignment, and so his universe must be anisotropic. We have solved the Einstein-Cartan field equa-

tions for an axisymmetric homogeneous Bianchi I model with metric

$$ds^2 = -dt^2 + X^2(t)dx^2 + Y^2(t)(dy^2 + dz^2) \quad (3)$$

We find  $\sigma = \sigma_0 R^{-3}$  (just as in the usual relativistic models),  $\rho \sim R^{-3}$ ,  $s \sim R^{-3}$ ,  $\omega^a = 0$ , where for this model  $R = (XY^2)^{1/3}$ . It can now be seen from equation 2 that the attractive shear term  $2\sigma^2$  is of the same order of magnitude as the repulsive spin term, and indeed if  $\sigma_0 > s_0$  a singularity cannot be prevented.

In these simple models  $\sigma^2 = \lambda s^2$ , where  $\lambda$  is a constant. In Trautman's model  $H^2 \sim \rho \sim R^{-3}$  except near  $t=0$ , where  $H = \frac{1}{3}\theta$  is the usual Hubble constant. Thus  $(\sigma^2/H^2) \approx \lambda s^2/\rho = \lambda(s^2/\rho)_{t=0}(\rho/\rho_{t=0})$ . At  $t=0$   $s^2 = \rho/2$ , in Trautman's model, and  $\rho_{\text{now}}/\rho_{t=0} = 10^{-84}$  if  $\rho_{\text{now}} = 10^{-29} \text{ g cm}^{-3}$ . Thus Trautman's suggestion for avoiding the singularity is only applicable if  $(\sigma^2/H^2)_{\text{now}} \lesssim 10^{-84}$ . But the best observational evidence is  $(\sigma^2/H^2)_{\text{now}} \lesssim 10^{-6}$  and so, unless the Universe is very isotropic indeed, a singularity cannot be avoided in this way.

We thank Bernd Schmidt and Martin Walker for a helpful discussion. Since this communication was submitted Copzynski<sup>3</sup> has independently constructed the same Bianchi I model.

JOHN STEWART  
PETR HÁJÍČEK

Max-Planck-Institut für  
Physik und Astrophysik,  
8000 München 40,  
Föhringer Ring 6,  
Germany

Received April 16, 1973.

- <sup>1</sup> Trautman, A., *Nature phys. Sci.*, **242**, 7 (1973).
- <sup>2</sup> Ellis, G. F. R., in *General Relativity and Cosmology*, XLVII Corso (Academic Press, 1971).
- <sup>3</sup> Copzynski, W., *Phys. Lett.*, **A**, **43**, 63 (1973).

## Reference Abbreviations

FROM September 1 it is *Nature's* intention that all abbreviations of references should conform to the style of the *World List of Scientific Periodicals*, fourth ed. (Butterworth, 1963-65). The changeover will be gradual and authors submitting manuscripts from now on are asked to ensure that the references are written appropriately.

Editorial, Advertising and Publishing Offices of NATURE

MACMILLAN JOURNALS LIMITED  
4 LITTLE ESSEX STREET, LONDON WC2R 3LF  
Telephone Number: 01-836 6633. Telegrams: Phusis London WC2R 3LF  
Telex 262024

MACMILLAN JOURNALS LIMITED  
711 NATIONAL PRESS BUILDING  
WASHINGTON DC 20004  
Telephone Number: 202-737 2355. Telex 64280

Advertisement Department  
MACMILLAN JOURNALS LIMITED  
4 LITTLE ESSEX STREET, LONDON WC2R 3LF  
Telephone Numbers: UK 01-836 6633. USA 202-737 2355

Subscription Department  
MACMILLAN JOURNALS LIMITED  
BRUNEL ROAD, BASINGSTOKE, HANTS RG21 2XS  
Telephone Number: Basingstoke 29242

Classified advertisements  
T. G. SCOTT & SON, LIMITED  
1 CLEMENT'S INN, LONDON WC2A 2ED  
Telephone Number: 01-242 6264/01-405 4743  
Telegrams: Textualist London WC2A 2ED  
Registered as a newspaper at the Post Office

Copyright © Macmillan Journals Limited, August, 6 1973