20. Meyer, K., and Rapport, M. M. In: F. F. Nord: *Advances in Enzymology, Hyaluronidases*, Vol. 13, pp. 199-236 (Interscience Publishers, Inc., New York, 1952).
21. Morse, B. S., and Nussbaum, M.: The detection of hyaluronic acid in the serum and urine of a patient with nephroblastoma. *Amer. J. Med.*, **42**: 996 (1967).
22. Österlin, S. W., and Jacobson, B.: The synthesis of hyaluronic acid in vitreous. *Exp. Eye Res.*, **7**: 497 (1968).
23. Pessac, B., and Defendi, V.: Cell aggregation: Role of acid mucopolysaccharides. *Science*, **175**: 898 (1972).
24. Powars, D. R., Allerton, S. E., Beierle, J., and Butler, B. B.: Wilms' tumor clinical correlation with circulating mucin in three cases. *Cancer*, **29**: 1597 (1972).
25. Robinson, H. C., and Hopwood, J. J.: The alkali cleavage and borohydride reduction of cartilage proteoglycans. *Biochem. J.*, **133**: 457 (1973).
26. Saito, H., Yamagata, T., and Suzuki, S.: Enzymatic methods for the determination of small quantities of isometric chondroitin sulfates. *J. Biol. Chem.*, **243**: 1536 (1968).
27. Schiller, S., Slover, G. A., and Dorfman, A.: The synthesis of hyaluronic acid by cell-free extracts of embryonic skin. *Biochem. Biophys. Res. Commun.*, **5**: 344 (1961).
28. Silbert, J. E.: Incorporation of ^{14}C and ^3H from labeled nucleotide sugars into polysaccharide in the presence of a cell-free preparation from cartilage. *J. Biol. Chem.*, **239**: 1310 (1964).
29. Tomasi, T. B., Jr., Robertson, W. V. B., Naeye, R., and Reichlin, M.: Serum hyperviscosity and metabolic acidosis due to circulating hyaluronic acid. *J. Clin. Invest.*, **45**: 1080 (1966).
30. Tomida, M., Koyama, H., and Ono, T.: Hyaluronic acid synthetase in cultured mammalian cells producing hyaluronic acid. *Biochim. Biophys. Acta*, **338**: 352 (1974).
31. Toole, B. P., Jackson, G. and Gross, J.: Hyaluronate in morphogenesis: Inhibition of chondrogenesis *in vitro*. *Proc. Natl. Acad. Sci. U. S. A.*, **69**: 1384 (1972).
32. Van Praag, D., Stone, A. L., Richter, A. J. and Farber, S. J.: Composition of glycosaminoglycans (mucopolysaccharides) in rabbit kidney. II. Renal cortex. *Biochim. Biophys. Acta*, **273**: 149 (1972).
33. Varadi, D. P., Cifonelli, J. A. and Dorfman, A.: Acid mucopolysaccharides in normal urine. *Biochim. Biophys. Acta*, **141**: 103 (1967).
34. Wasteson, Å., Westermark, B., Lindahl, U., and Ponten, J.: Aggregation of feline lymphoma cells by hyaluronic acid. *Int. J. Cancer*, **12**: 169 (1973).
35. The abbreviations used are: $\Delta\text{di Ch-S}$, 3-O- β - Δ -4,5-glucuronosyl-*N*-acetyl-D-galactosamine-6-O-SO₄; $\Delta\text{di Ch-4S}$, 3-O- β - Δ -4,5-glucuronosyl-*N*-acetyl-D-galactosamine-4-O-SO₄; $\Delta\text{di Ch}$, 3-O- β - Δ -4,5-glucuronosyl-*N*-acetyl-D-galactosamine; $\Delta\text{di HA}$, 3-O- β - Δ -4,5-glucuronosyl-*N*-acetyl-D-glucosamine.
36. The authors are grateful to Drs. Emily Pang and John Moehr for providing us with the Wilms' tumor, urine, and plasma supplies.
37. This investigation was supported by NIH Grants AM-05996, HD-04583 and HD-09402 and by the Home for Destitute Crippled Children.
38. Dr. John J. Hopwood is a Kennedy Fellow in the Neurosciences. Present address: Department of Chemical Pathology, Adelaide Children's Hospital, North Adelaide, S. A. 5006 (Australia).
39. Requests for reprints should be addressed to: Albert Dorfman, M.D., University of Chicago, Department of Pediatrics, 950 East 59th Street, Hospital Box 413, Chicago, IL 60637 (USA).
40. Received for publication February 15, 1977.
41. Accepted for publication April 5, 1977.

Copyright © 1978 International Pediatric Research Foundation, Inc.
0031-3998/78/0012-0001\$02.00/0

Printed in U.S.A.

Pediat. Res. **12**: 56-57 (1978)

Letter to the Editor: Metabolic Studies in Tyrosinemia

NEIL R. M. BUIST,⁽¹⁾ JACK H. FELLMAN, AND NANCY KENNAWAY

University of Oregon, Health Sciences Center, Portland, Oregon

The article by K. F. Faull and his colleagues entitled "Metabolic Studies on Two Patients with Non-hepatic Tyrosinemia Using Deuterated Tyrosine Loads" (3) raises several points which appear to us to require comment. The term "non-hepatic tyrosinemia" is inappropriate for a disorder in which hepatic enzymes are alleged to be deficient and only serves to complicate the present nomenclature. The tyrosinemia which is associated with liver damage and a Fanconi syndrome is clinically quite distinct from the tyrosinemia associated with eye and skin lesions and mental retardation without liver damage. In order to distinguish these two clinical entities we have called them type 1 and type 2, respectively (2).

The enzyme assays in Dr. Faull's study are critically inadequate. Since liver contains a nonspecific mitochondrial transaminase (aspartate aminotransferase) capable of accepting tyrosine as substrate (11) any meaningful investigation of tyrosine aminotransferase must be carried out on supernatant fractions of liver homogenates (7, 8). Granner's method requires the use of diethyldithiocarbamate as an inhibitor of hydroxylase activity in fresh liver samples and this was not employed. Moreover, saturating concentrations of substrate were not present in the assay for *p*-hydroxyphenylpyruvate hydroxylase activity even if all the [^{14}C]tyrosine was converted to ketoacid. Finally, the choice of a paper chromatographic procedure is baffling when

more simple, sensitive, and accurate methods already exist (5, 10, 12).

The [^3H]tyrosine experiments are predicated on the retention of the ^3H on the phenolic ring during its metabolic passage. This must be constant for any meaningful interpretation of these experiments. Under physiologic conditions, ^3H exchange at the *ortho* position of tyrosine could be considerably greater than under the strong acid conditions employed by the authors to establish stability of the labeling. We thus have no means of interpreting the reported urinary metabolic results which might vary with storage and even with the pH of the urine. Indeed, the ^3H study may be more misleading than instructive. The [^3H]tyrosine load (120 mg/kg) could not dissolve in 100 ml as described by the authors; moreover, at the pH of ileal fluid, the solubility is only about 750 mg/liter. The stool content of ^3H compounds is not mentioned and failure of absorption could account for a substantial proportion of the load. Some evidence for this is apparent from the 165-fold increase in urinary *p*-hydroxyphenylacetate after ^3H -tyrosine load in their normal subject. It has been established that the bulk, if not all, of the urinary *p*-hydroxyphenylacetate originates from gut bacterial action on tyrosine (1, 6). Furthermore, the blood levels of [^3H]tyrosine in patients continued to increase for 8 hrs, which also suggests delayed absorption of the amino acid from the gut.

The statement that the "pattern of urinary metabolites indicated a persistent defect in *p*-hydroxyphenylpyruvic acid oxidase" is simplistic. It certainly appears that both Faull's patients handled the [²H]tyrosine load very differently but to use this data to postulate substrate inhibition of *p*-hydroxyphenylpyruvate hydroxylase *in vivo* is fallacious. The so-called substrate inhibition is a contrivance of the enzyme assay in which oxygen inactivation of the enzyme occurs in the presence of the ketoacid. At physiologic pO₂ no inhibition is observed (5). The urinary metabolite patterns and the different handling of a tyrosine load in the two patients could just as well be explained by a difference in the amount of residual cytosol tyrosine aminotransferase activity. In this condition, the ketoacid accumulates as a result of mitochondrial transamination of tyrosine in tissues which lack the hydroxylase (4, 9).

In summary, it is regrettable that a decisive investigation of the enzyme defects was not achieved. It is clear that these patients did not have tyrosinemia type 1 but it remains uncertain as to whether they both had tyrosinemia type 2 or whether one had a primary defect in *p*-hydroxyphenylpyruvate hydroxylase.

REFERENCES AND NOTES

1. Borud, O., Midtredt, T., and Gjessing, L. R.: Urinary phenolic compounds in gnotobiotic and conventional rats on a free diet, and before and after L-DOPA loading on a milk diet. *Acta Pharmacol. Toxicol.*, **30**: 185 (1971).
2. Buist, N. R. M., Kennaway, N. G., and Fellman, J. H.: Disorders of

- tyrosine metabolism. In: W. H. Nyhan: *Heritable Disorders of Amino Acid Metabolism*, p. 160 (John Wiley, New York, 1974).
3. Faull, K. F., Gan, I., Halpern, B., Hammond, J., Im, S., Cotton, R. G. H., Danks, D. M., and Freeman, R.: Metabolic studies on two patients with non-hepatic tyrosinemia using deuterated tyrosine loads. *Pediat. Res.*, **11**: 631 (1977).
4. Fellman, J. H., Buist, N. R. M., Kennaway, N. G., and Swanson, R. E.: The source of aromatic ketoacids in tyrosinaemia and phenylketonuria. *Clin. Chim. Acta*, **39**: 243 (1972).
5. Fellman, J. H., Fujita, T. S., and Roth, E. S.: Assay, properties and tissue distribution of *p*-hydroxyphenylpyruvate hydroxylase. *Biochim. Biophys. Acta*, **284**: 90 (1972).
6. Fellman, J. H., Roth, E. S., and Fujita, T. S.: Decarboxylation to tyramine is not a major route of tyrosine metabolism in mammals. *Arch. Biochem. Biophys.*, **174**: 562 (1976).
7. Fellman, J. H., Vanbellinghen, P. J., Jones, R. T., and Koler, R. D.: Soluble and mitochondrial forms of tyrosine aminotransferase: Relationship to human tyrosinemia. *Biochemistry*, **8**: 615 (1969).
8. Kennaway, N. G., and Buist, N. R. M.: Metabolic studies in a patient with hepatic cytosol tyrosine aminotransferase deficiency. *Pediat. Res.*, **5**: 287 (1971).
9. Kennaway, N. G., Buist, N. R. M., and Fellman, J. H.: The origin of urinary *p*-hydroxyphenylpyruvate in a patient with hepatic cytosol tyrosine amino transferase deficiency. *Clin. Chim. Acta*, **41**: 157 (1972).
10. Lindblad, B.: Radiochemical assays for *p*-hydroxyphenylpyruvate hydroxylase activity in human liver. *Clin. Chim. Acta*, **34**: 113 (1971).
11. Miller, J. E., and Litwack, G.: Purification, properties, and identity of liver mitochondrial tyrosine aminotransferase. *J. Biol. Chem.*, **246**: 3234 (1971).
12. Raheja, M. C., Dylewski, I., and Crawhill, J. C.: Estimation of *p*-hydroxyphenylpyruvate hydroxylase activity based on ¹⁴CO₂ evolution from the ¹⁴C-labeled substrate. *Can. J. Biochem.*, **51**: 172 (1973).
13. Correspondence should be addressed to: Neil R. M. Buist, M.D., Professor of Pediatrics and Medical Genetics, University of Oregon, Health Sciences Center, Portland, OR 97201 (USA).

Copyright © 1978 International Pediatric Research Foundation, Inc.
0031-3998/78/0012-0001\$02.00/0

Printed in U.S.A.

Pediat. Res. **12**: 57-58 (1978)

Letter to the Editor: Reply to Dr. Buist

D. M. DANKS

Genetics Research Unit, Royal Children's Hospital Research Foundation, Parkville, Victoria, Australia

Thank you for the opportunity of replying to the criticisms of Dr. Buist and colleagues. The terminology of tyrosinemia is indeed confusing. Perhaps we should have used the term "tyrosinemia without liver damage."

We recognize deficiencies in the enzyme assays and therefore laid rather little weight upon these results in our argument; however, there were clear differences between the two patients. We can firmly dismiss the suggestion that deuterium atoms have been lost from the phenolic ring by chemical exchange during storage because such a process would alter the ratio of the D₂ to the D₁ peaks on mass spectrometry, producing an increase in the D₁ peak and a decrease in the D₂ peak. The ratio of these peaks was monitored in all samples and did not change significantly.

Absorption of the tyrosine loads used was clearly slow and probably incomplete. Parenteral administration of a smaller load would undoubtedly be preferable. Nonetheless, as our critics themselves point out, the studies of the control subject give us some idea of the amount of phenolic acids contributed

by absorption of the products of bacterial metabolism and this is clearly far too small to have any influence on the overall results obtained.

We are left with the principal observation of our study basically unaltered by the criticisms raised. One patient excreted approximately the same very large proportion of the administered label in the urine whether he was tested with high serum tyrosine or with serum tyrosine reduced by dietary treatment. In the other patient the dietary control resulted in a 100-fold reduction of the proportion of the label excreted in the urine as tyrosine and tyrosyl derivatives.

In considering these results it is important to note that the serum tyrosine level was still 2-3 times the upper limit of normal in both patients while on dietary treatment and so the rate-limiting step of tyrosine catabolism was already overloaded before the labeled tyrosine load was administered. Consequently, one should expect the handling of the load to be comparable in the two patients under these circumstances if indeed the two patients had defects of the same enzyme as