Response

JAMES E. BOWES

MRS. BUMPERS RELINQUISHED HER TIME TO DR. BOWES, WHO MADE THE FOLLOWING GUEST PRESENTATION

End Measles, Inc., Indianapolis, Indiana, USA

I have been in the business of bringing the product to the consumer for 22 years; *i.e.*, vaccinations to the consumer. In 1960, I helped break the polio epidemic in Rhode Island by organizing a mass immunization campaign. There were over 103 reported cases of polio. That was the second to the last major epidemic in this nation.

Rhode Island was also the site for the first statewide oral polio Sunday. This was accomplished on three different Sundays, 2 months apart. Eighty percent of the total population in the state turned out in one day; 680 thousand people on each of those separate Sundays with no federal, state, or local funds.

I helped conduct the first statewide Single Day Measles Immunization Campaign in the country in Rhode Island in January of 1966. Despite the six inches of snow on the ground at 6:00 a.m., we immunized 33 thousand children on that one day. We were advised by the manufacturer not to conduct a mass immunization program because of the unknown risks involved. We had only 2% adverse reactions.

I also served as a consultant to the Detroit program where 174 thousand children turned out on one Sunday in October of 1966. We had volunteers assigned to run the clinics; a volunteer doctor in every one of them.

At the annual meeting of the American Medical Association in 1967, we conducted a scientific exhibit where we were able to press a button which indicated the visitor's town code and consequently there would appear on the screen the number of measles cases which could be expected in that particular town during the next year, how many patients could be hospitalized, child school-days absent, cost factors, etc. At the same time, in 20 sec, the typewriter made a printout which the questioner could take with him. These cost/risk results were published by the Center for Disease Control.

For 12 years, we have kept a register of the number of children who are susceptible to measles in approximately 3 thousand countries around the world.

Dr. Sabin has already spoken about the various obstacles to immunizations. In 1970, in one of our published newsletters entitled "Measles Memo", which went to 2400 people, including all members of Congress, governors, health officers, and members of the media, it spoke of the "hard to reach" children (Fig. 1). It expressed the pleas of a child, saying "I would be easy to reach if my parents could afford it, if there was a school program, if the local health officers were concerned", and finally, "if I were big enough to speak up for my right to health".

Six years ago, nine young epidemiologists in the nation met for a 2-day work conference in Atlanta to discuss what our national stance should be on immunizations. Out of this meeting there was drafted a resolution for a national federal commitment on behalf of childhood immunizations. News stories were released and published in various medical journals. However, there was not enough funding to support a full ongoing program.

Four years ago, I founded a not-for-profit organization called End Measles, Inc. Of the 400 foundations contacted, I heard from four, and received a total of \$6400. Recently End Measles, Inc., with the help of computer analysts, studied the measles cases in 18 states and in the last 4 years, along with the rates of immunizations, and we have come up with a method of predicting epidemics 3 years in advance in 356 communities in this country.

Whether they will occur or not depends on adequate immunization programs. The risk of measles still exists. Thirteen states have had more measles this year than last.

In this study we also developed a concept which the health officer can use to indicate when he is going to have a measles epidemic. We call it a Measles Epidemic Threshold (MET) (Fig. 2). This establishes the number of measles cases in a given population which would indicate the beginning of an epidemic. For instance, in my community of 700 thousand people, the MET would be 70 cases. One starts counting the cases when children get back into school in the fall. If the number should reach 50 or 60 cases, then there could be an epidemic. For a small community of 100,000 people, the MET would be 25 cases. If we hope to have any effect in eliminating measles from this country, we can never allow the number of measles cases to rise above 26/100,000 people.

I have seen a lot of epidemics and have predicted some to people in my own area. The year after I came to the Indianapolis Health Department, there were 1200 measles cases in an area where I was not allowed to enact our prevention program during the previous year. With the concept of MET points, it might be easier to convince the medical community.

I agree with Dr. Mortimer and Dr. Sabin about most of the obstacles to effective immunization programs. We are all living with such obstacles. In a private survey of 600 public health nurses 6 years ago, I asked all kinds of questions about measles immunization programs. Twenty percent answered that the obstacle was within the medical community: "the medical society won't let us do it." In most cases, the obstacles are within our own family; our own medical family. The medical family is broad, and extends to the superintendent of schools. When we searched the school health records of children in our 140 schools more than a year ago, we found that only 47-50% of the children were completely immunized.

We prevailed upon a school to allow us to initiate a program of exclusion in the beginning of our program. We provided the principal with a list of "bad guys"; students who had one or more immunization deficiencies. We then sent a letter home telling the parents to take the child to the doctor. Failing that, the child would be recommended for exclusion, effective in 2 weeks. In a school of 700, usually we would have 150 children on the bad guy list. The exclusion letter was then sent out. By noon on Monday there were only 25 left and the next day only four. Mothers would come to school Monday morning with notes from their doctors saying their children had the necessary shots. The biggest obstacle was to convince the principal that he was not going to get a lot of phone calls. After all, the schools have a service that parents want, and they will do what is necessary to keep their children there.

In March of 1970, I put out a Measles Memo entitled "What Happened?" We have had two previous opportunities to do so as shown on the Memo's bar graph (Fig. 3). Measles was low in 1968, but we did not respond fully. Rubella was the concern the next year. Measles went down again in 1974. Now we finally have the opportunity to eradicate measles, as Dr. Hinman says, if we take action.

The federal and state governments cannot do it alone. Private physicians cannot do it alone. They need a catalyst to help them get the product to the customer.

MEASLES MEMO

& ERADICATION PROGRESS REPORT

VOLUME 4, NO. 10

NOVEMBER 23, 1970

HARD TO REACH Who, Me?

Well, I'd Be Easy To Reach

If my parents could afford to take me to a private physician.

If my parents knew that preventive immunizations would save me from a lot of sick days and not make them lose time from work.

If I lived in a state that has a school entrance immunization law.

If I go to a school with an active school health program.

If I didn't live in a community that lets epidemics happen.

If I lived in a community where the health officer had a deeper concern for us "hard-to-reach."

If I lived in an area that had more active well-baby clinics.

If I didn't live in the "inner city."

If I didn't live so far away from the only immunization center.

If I were BIG enough to speak out for my "RIGHT-TO-HEALTH."

And That's Why There Are 10 Million
Hard-To-Reach Kids Like Me

Measles epidemic threshold
(M E T)

The following table gives the number of measles cases during a single measles season occurring in a given city or county population that indicates the imminent possibility of a measles epidemic.

<u>Population</u>		<u>MET</u>		
10,000 -	15,000	6	measles cases	
15,001 -	20,000	7	"	"
20,001 -	25,000	8	"	"
25,001 -	30,000	9	"	"
30,001 -	35,000	10	"	"
35,001 -	40,000	11	"	"
40,001 -	45,000	12	"	"
45,001 -	50,000	13	"	"
50,001 -	55,000	14	"	"
55,001 -	60,000	15	"	"
60,001 -	65,000	16	"	"
65,001 -	70,000	17	"	"
70,001 -	75,000	18	"	"
75,001 -	80,000	19	"	"
80,001 -	85,000	20	"	"
85,001 -	90,000	21	"	"
90,001 -	95,000	22	"	"
95,001 -	100,000	23	"	"
100,001 -	125,000	25	"	"
125,001 -	150,000	30	"	"
150,001 -	200,000	35	"	"
200,001 -	250,000	40	"	"
250,001 -	300,000	45	"	"
300,001 -	400,000	50	"	"
400,001 -	500,000	55	"	"
500,001 -	750,000	60	"	"
750,001 -	1,000,000	70	"	"
1,000,001 -	2,000,000	100	"	"
2,000,001 -	4,000,000	190	"	"
4,000,001 -	6,000,000	300	"	"
6,000,001 -	8,000,000	430	"	"

When the above cases have occurred in a city or county, it is time to:

M E C - - Measles Epidemic Control
with the intensive measles
immunization campaign.

End Measles, Inc.
August 9, 1978

MEASLES MEMO

& ERADICATION PROGRESS REPORT

VOLUME 4, NO. 2

MARCH 2, 1970

fewer than 1,000 by 1970

MEDICAL NEWS

JAMA, Jan 29, 1968 • Vol 203, No 5

Measles Expected To Decline Again

Measles Resurgence
Sparks New Campaign
To Immunize Children

Health Officials Had Thought
Disease Was Licked, but It
Hits in Poor Neighborhoods

By JONATHAN SPIVAK
Staff Reporter of THE WALL STREET JOURNAL
WASHINGTON—McBride
Past. Hein

Plan to Eradicate Measles in 1967 Detailed by PHS

PUBLIC HEALTH ASSOCIATION
Measles Eradication Campaign.
Etiology of Hip Fractures.
Effects of Prenatal Fluoridation.

Medical Tribune—World Wide Report
SAN FRANCISCO—Details of the measles
eradication campaign launched by the
United States Public Health Service
with the goal of eliminating the disease
from this country by the end of 1967
were disclosed here at the annual meet-
ing of the American Public Health As-
sociation.

THE WALL STREET JOURNAL
Friday, February 20, 1970

Figure 1
REPORTED CASES OF MEASLES BY 4-WEEK PERIOD, USA
EPIDEMIOL. YEAR 1969-70 COMPARED WITH 1966-67 AND 1968-69

"With a concerted effort, it is possible to erad-
icate measles in 1967."

William H. Stewart
SURGEON GENERAL,
U.S. PUBLIC HEALTH SERVICE

'66 Measles Cases At 21-Year Low

Medical News—World Wide Report
ATLANTA, GA.—The number of measles
cases reported in the United States in
1966 was the lowest in 21 years, ac-
cording to the National Communicable
Disease Center here of the U.S. Public
Health Service.

Mass Immunization Can End Measles by 1969

ATLANTIC CITY, N.J. Measles can be
eradicated by 1969 if mass-immuniza-
tion programs are used. If not, it will
take until 1974, according to
AMERICAN JOURNAL OF HYGIENE
1620 AUGUST 1967

WHAT HAPPENED?