ANNUAL MEETING

European Society for Paediatric Research

Leuven, Belgium, September 3-6, 1979*

HORMONAL INTERACTION AND GLUCOSE HOMEOSTASIS

M. CORNBLATH

National Institutes of Health, Bethesda, MD., USA

Throughout development, the endocrine factors involved in glucose homeostasis vary in relation to age and maturation. The endocrine-metabolic milieu in utero changes dramatically at birth and then gradually throughout infancy and childhood. The available data demonstrate a unique response of the developing infant in regulating glucose homeostasis in both health and disease. Hormone values alone, do not adequately explain the vulnerability of the neonate or infant to hypoglycemia. Additional studies involving recent advances in our understanding of hormone action through receptor sites, feed-back mechanisms and enzyme regulation will be necessary to extend our understanding of the basic mechanisms involved.

In the fetus, carbohydrate balance depends upon the maternal supply of glucose, via the placenta and, as far as is known, on fetal production of insulin and cortisol. Little data are available concerning the interactions or effects of hypothalamic factors or other hormones on fetal carbohydrate homeostasis. However, almost immediately after birth, specific, yet unique responses of multiple endocrine secretions occur. The blood glucose values tend to be lower in the neonatal period than later in the presence of normal values for cortisol, low, for insulin, and high, for glucagon, hGH, T4 and reverse T3.

The variety of hypoglycemic syndromes in the neonatal and early infancy periods elucidate some unique aspects of hormonal secretion and actions. Hyperinsulinism, in utero whether transient as in the infant of the diabetic mother, or persistent due to malformation of islet cell formation and function, can produce a number of effects before and after birth. In the primate model insulin alone has been shown to induce macrosomia, extramedullary hematopoiesis, lipogenesis and glycogen storage. There are all present in the infant of the diabetic mother, but rapidly disappear over a short period of time. In addition, increased insulin receptor sites have been demonstrated on the monocytes of cord blood of infants of diabetic mothers. However, in the infant born with an abnormality of islet cell function or formation, e.g. nesidioblastosis, the profound hypoglycemia can be life threatening even in the presence of pituitary and adrenal hormonal responses. Some data would suggest that the malformation involves not only the insulin producing beta cell, but the alpha, delta and pancreatic polypeptide cells as well and thus a total paracrine dysfunction occurs which interferes with glucose homeostasis. Preliminary data on infants with persistent hyperinsulinemia from birth and subsequently treated either operatively or with diazoxide suggest that increased insulin receptor sites may persist. Thus, the abnormality noted in hyperinsulinism may have far reaching effects beyond those usually attributed to insulin alone.

The syndrome of congenital hypopituitarism is another example of unique endocrine-glucose interrelationships in the neonatal period. This syndrome clearly indicates that the absence of growth hormone and cortisol may result in profound, even fatal, hypoglycemia in the first hours after birth. Yet, other infants with the same condition may maintain normoglycemia and later develop, ketotic hypoglycemia between 1 and 5 years of age. Hypothalamic factors have been found deficient in some of these patients leading to an inadequate feedback mechanism of carbohydrate homeostasis.

Ketotic hypoglycemia is the most prevalent form of low blood glucose between 1 and 5 years of age. Some of these were infants who had been hypoglycemic in the neonatal period as SGA babies, or because of panhypopituitarism or isolated growth hormone deficiency, while others had no previous evidence of carbohydrate intolerance. Fasting ketosis followed by hypoglycemia may reflect inadequate normal feedback mechanisms which regulate hormones responsible for glycogenolysis and gluconeogenesis. These abnormal responses disappear beyond age 7 or 8. Thereafter the child begins to show normal hormonal-glucose relationships.

Glucose homeostasis is the result of complex interactions that change profoundly throughout development. Their complex nature require the application of recently developed concepts to understand the underlying mechanisms.

THE CONTROL OF CARBOHYDRATE METABOLISM IN THE LIVER

H.G. HERS

Laboratoire de Chimie Physiologique, Université de Louvain, and International Institute of Cellular and Molecular Pathology, Brussels, Belgium

Glucose occupies a unique position in intermediary metabolism for two reasons:

1. it is the substrate of glycolysis which is the only pathway to produce ATP in anaerobic life
2. it is the major and usually the unique substrate for brain metabolism.

The second reason is by far the most important in the human body and one major role of the liver is to maintain a normal level of glycaemia. The control of the level of glycaemia by the liver involves the storage of glucose as glycogen and the formation of glucose from endogenous source by gluconeogenesis.

The control of glycogen metabolism in the liver

As explained in detail in a recent review (Hers, Ann. Rev. Biochem., 45, 167-189, 1976) the rate limiting steps of glycogen synthesis and breakdown in the liver are catalyzed by glycogen synthase and glycogen phosphorylase. Each of these enzymes exist in two forms: one active (a) and the other one inactive (b). The a and b forms are interconvertible through phosphorylation by kinases and dephosphorylation by phosphatases. Glucose is readily diffusible inside of the liver cell. It binds to phosphorylase a which can be considered as the glucose receptor of the liver. The glucose bound phosphorylase a is rapidly converted to phosphorylase b by phosphorylase phosphatase. A first effect of glucose is to decrease and eventually to arrest glycogen degradation in the liver. Furthermore, phosphorylase a is a strong inhibitor of synthase phosphatase, the enzyme that activates glycogen synthase. Therefore, the disappearance of phosphorylase a as a result of glucose binding allows the activation of glycogen synthase and the initiation of glycogen synthesis. Various ionic and hormonal effects including those of insulin appear to be explained by a primary action on the inactivation of phosphorylase. Glucagon causes glycogenolysis by the intermediary of cyclic AMP, activation of protein kinase, phosphorylase kinase and phosphorylase. Each step of this glycogenolytic pathway has been observed in hepatocytes incubated in the presence of the hormones. In contrast, catecholamines and other agonists including vasopressin and angiotensin stimulate glycogenolysis by another mechanism, possibly involving a change in calcium concentration. Insulin counteracts the effect of these glycogenolytic agents by acting at various steps of the glycogenolytic cascade.

The control of gluconeogenesis and the futile cycles

Futile cycle is a metabolic interconversion of which the net balance is the consumption of ATP. Three such cycles exist on the pathway of glycolysis and gluconeogenesis (1) at the level of glucose/glucose 6-phosphate interconversion (2) at the level of fructose 6-phosphate/fructose diphosphate interconversion (3) at the level of pyruvate/phosphoenolpyruvate interconversion. The best known mechanism by which glucagon stimulates gluconeogenesis is the inactivation, by phosphorylation, of pyruvate kinase, preventing futile recycling between phosphoenolpyruvate and pyruvate. The control of the recycling between fructose 6-phosphate and fructose diphosphate is still badly understood but appears to be controlled by glucagon also. Recycling between glucose and glucose 6-phosphate occurs continuously and is controlled only by the concentration of substrate: it plays a major role in the control of glucose uptake and glucose output by the liver.

THE TRANSPORT OF VITAMIN D

R. BOUILLON and H. VAN BAELEN

Rega Institute, University of Leuven, Leuven, Belgium

Vitamin D and its hydroxylated metabolites are transported in the blood, bound to a transport protein. For this purpose, cartilaginous fish and amphibia use lipoproteins but bony fish, reptiles, birds and mammals have a specific vitamin D-binding protein, called DBP. We have isolated DBP from the serum of man, rat and chick and compared their characteristics.

GENERAL CHARACTERISTICS OF DBP

	Man	Rat	Chick
Molecular weight	56000	52000	60000
Sedimentation coefficient	4.1 S	4.1 S	4.1 S
Amino acid analysis	known	known	known
Electrophoretic mobility	alpha	alpha	beta
Isoelectric point	± 5	± 5	± 6
Thermostability increased by excess ligand	+	+	+
Binding characteristics			
K_a for 25OHD ₃ (4C,pH 7.4)	$10^8 M^{-1}$	$10^8 M^{-1}$	$10^8 M^{-1}$
n binding sites	1	1	1
Tendency to polymerisation	-	+	-
Interaction with a cytosolic protein	+	+	+

The concentration of DBP is high in the three species (6 to 13 μM) so that less than 3% of all binding sites are normally occupied. The regulation of the serum concentration is quite dissimilar in the three species. There is little age-dependent variation in man and chick but an important increase in serum DBP in the growing rat. Estrogens increase the DBP level in man (and probably also the chick) whereas in the rat androgens increase the DBP concentration. The vitamin D status does not influence the DBP level in any of the species. Human and rat DBP are genetically heterogenous, coded by a pair of codominant autosomes. Human DBP was therefore previously known as "group-specific component or 6c".

The significance of the serum transport protein for vitamin D can probably be searched in its high affinity and high capacity for 25-hydroxyvitamin D. DBP therefore functions as a storage protein accumulating 25-hydroxyvitamin D during periods of access to vitamin D and releases it, thereafter, very slowly for biological activation or inactivation in the kidney. DBP is also very important in the placental transfer of 25-hydroxy-vitamin D: since the human neonatal DBP level is only half that of the mother, the neonatus also has only half the circulating reserve of 25-hydroxyvitamin D. The DBP level remains constant in most diseases, except in cirrhosis of the liver and nephrotic syndrome. Hypovitaminosis D occurs in the latter disease due to the continuous loss of 25-hydroxyvitamin D together with the loss of DBP in the urine.

Serum DBP also interacts with an ubiquitous intracellular protein to form the 6S intracellular binding protein but the significance of this complexation remains to be explored. Finally, DBP is also used in the competitive protein binding assay to measure 25-hydroxyvitamin D in biological fluids. This assay is especially useful to detect (latent) hypovitaminosis D and vitamin D intoxication.

VIRUS REPLICATION - AN INTRODUCTION

A. BILLIAU

Rega Institute, University of Leuven,
Leuven, Belgium

Viruses can only multiply in an internal cellular environment. They lack subcellular organelles such as nuclei, mitochondria, ribosomes as well as cytoplasmic components that are necessary for the synthesis of their own structural components: nucleic acids (RNA or DNA), proteins, carbohydrates and lipids. Thus, viruses cannot multiply but must be replicated by the cells that they invade. The essential steps of virus replication are similar for all virus classes: first, a virus particle (virion) must attach to the outer surface of the host cell. Next, it penetrates the cell membrane and enters the cytoplasmic environment. Some or all of the outer surface layers of the virus (envelope and capsid) are removed, so that the viral genome (DNA or RNA) becomes accessible to the cellular organelles and enzymes which will initiate the replication process. Some of the genetic information is used to synthesize virus-specific enzymes (polymerases) which are necessary for the replication of the viral nucleic acid. Other parts of the viral genome code for the structural proteins. The newly synthesized viral nucleic acid and proteins are then assembled (maturation) into new viral particles which may leave the cells either by simple cell lysis or by a budding process at the membranes of the outer cell surface or in the endoplasmic reticulum.

This basic strategy of virus infections leaves ample space for variations depending on whether the viral genome is composed of DNA or RNA, on whether it is double- or single-stranded, linear or circular, monolithic or fragmented. The parental RNA may either be directly used by ribosomes (+ strand viruses), or it may first need transcription into a complementary strand (- strand viruses). Enveloped viruses probably require the synthesis of specific carbohydrates and lipids.

Depending on the genetic make-up of the host cell and on the conditions of infection, the virus cycle may be productive (i.e. many new infectious particles are released by each infected cell) or it may be restrictive, or even abortive. Some viruses may remain latently present in cells over a long time and over a large number of generations. Several mechanisms for such chronicity of virus infection have been discovered. Certain taxa of viruses have the unique capability to integrate their genome, or a fragment thereof, into that of the cell. Chronic infections may also be maintained by the generation of defective-interfering (DI) particles, which contain only part of the virus genome, but which are replicated when they are present in cells, along

with complete virus particles. In general the small size of the DI-genome allows it to be replicated more rapidly than the intact genome and hence competes strongly for the polymerases. In certain conditions, DI-particles can alter the lytic pathway of infection and rapidly establish persistent infections in which the cells survive but carry in them considerable amounts of virus antigens. This is an important area in virology to-day as it may be related to the development of long-term degenerative diseases which may have a viral etiology.

SLOW VIRUS DISEASES OF THE CENTRAL NERVOUS SYSTEM

V. TER MEULEN

Institute für Virologie und Immunbiologie, University of Würzburg
Würzburg, F.R. Germany

During the last 3 decades a new concept of an infectious process has been derived by studying certain Central Nervous System (CNS) diseases in animals. These infections are characterized by an incubation period lasting for many months to years and by a predictable, protracted clinical course usually leading to death. It has been found that these disorders, which are referred to as slow virus diseases, are related to unconventional and conventional agents, both of which have been associated with naturally occurring diseases in animals and man. The unconventional agents, which have not been visualized, isolated or characterized reveal unusual biological and physico-chemical properties, unlike any known virus group. These agents are associated with scrapie in sheep, encephalopathy in mink and Kuru or Creutzfeldt-Jakob disease in man. The conventional viruses isolated from slow virus diseased brain material, such as Visna, Subacute Sclerosing Panencephalitis (SSPE). Progressive Rubella Panencephalitis (PRP) or Progressive Multifocal Leukoencephalopathy resemble classical viruses with typical structural, physico-chemical and biological characteristics. For this group of viruses, progress has been made in the understanding of the infectious process, since the isolates provide an experimental basis for virological and immunological studies in these diseases.

From a pediatric point of view, only SSPE and PRP are of interest, since the other slow virus diseases are usually not seen in children. SSPE is caused by a measles virus infection. Measles virus (referred to SSPE virus) has been isolated from CNS and lymphoid tissue. Moreover the patients reveal a pronounced antimeasles hyperimmunresponse which is pathognomonic for this disorder.

Virological studies indicate that SSPE viruses are related but not identical with typical virus isolated from acute measles. At the present state of investigation, SSPE is considered a late complication of acute measles infection. PRP is associated with a persistent rubella virus infection of the CNS. Virological and immunological data reveal great similarities to SSPE. However, information on the pathogenetic mechanism of this disease process are not available.

Previous investigations on slow virus diseases have demonstrated the different ways a virus infection can result in a chronic CNS disorder. Moreover, they have shown that virus-host interactions exist which lead to diseases lacking the common characteristics of an infectious process. Obviously, many virological and immunological problems have to be solved before the complexity of these disorders are understood, but it already can be expected, that other human diseases will be found which are caused by a slow virus infection.

VIRAL INFECTIONS OF THE CENTRAL NERVOUS SYSTEM AND LOCAL SYNTHESIS OF OLIGOCLONAL ANTIBODIES

B. VANDVIK

Department of Neurology, Rikshospitalet, Oslo, Norway

Oligoclonal IgG, synthesized locally in the central nervous system, is commonly demonstrable in the cerebro-spinal fluid in subacute sclerosing panencephalitis and some other viral central nervous system infections and in multiple sclerosis.

Extensive studies of cerebro-spinal fluid and brain tissue in subacute sclerosing panencephalitis showed that the bulk of the oligoclonal IgG is measles virus-specific antibody. A principally similar association of oligoclonal cerebrospinal fluid-IgG with virus-specific antibody activities was observed in progressive rubella virus panencephalitis, herpes simplex encephalitis and mumps meningitis. These findings indicate that the occurrence of oligoclonal IgG in the cerebro-spinal fluid in viral central nervous system infections reflects a specific antibody response to virus antigens in the central nervous system.

In some cases of subacute sclerosing panencephalitis and other central nervous system infections, a local synthesis of small amounts of oligoclonal antibodies to viruses other than that causing the disease was observed. This suggests that the immunizing drive caused by a virus infecting the central nervous system sometimes may lead to an activation of cell clones producing antibodies to unrelated antigens. Such mechanisms may be relevant in explaining the local synthesis in the central nervous system of oligoclonal antibodies to various viruses in multiple sclerosis.