- injury and partial hepatectomy. *Int. J. Cancer*, 22: 181 (1978).
12. Nayak, N., Das, P., Bhuyan, U., and Mittal, A.: Localization of α -fetoprotein in human and rat fetal livers: An immunohistochemical method using horseradish peroxidase. *J. Histochem. Cytochem.*, 22: 4 (1974).
 13. Nayak, N. and Mittal, I.: The dynamics of α -fetoprotein and albumin synthesis in human and rat liver during normal ontogeny. *Am. J. Pathol.*, 86: 359 (1977).
 14. Oledzka-Slotwinska, H. and Desmet, V.: Morphologic and cytochemical study on neonatal liver "giant" cell transformation. *Exp. Mol. Pathol.*, 10: 162 (1969).
 15. Postlethwaite, A. E., Jackson, B. K., Beachey, E. H., and Kang, A. H.: Formation of multinucleated giant cells from human monocyte precursors: Mediation by a soluble protein from antigen and mitogen-stimulated lymphocytes. *J. Exp. Med.*, 155: 168 (1982).
 16. Purtilo, D. T. and Yunis, E. J.: Alpha-fetoprotein: Its immunofluorescent localization in human fetal liver and hepatoma. *Lab. Invest.*, 25: 291 (1971).
 17. Ruebner, B. and Miyai, K.: The pathology of neonatal hepatitis and biliary atresia with particular reference to hemopoiesis and hemosiderin deposition. *Ann. N.Y. Acad. Sci.*, 3: 375 (1963).
 18. Sell, S.: Distribution of α -fetoprotein and albumin containing cells in the livers of Fischer rats fed four cycles of N-2-fluorenyl-acetamide. *Cancer Res.*, 38: 3107 (1978).
 19. Sternberger, L., Hardy, P., Cuculis, J., and Meyer, H.: The unlabeled antibody enzyme method of immunohistochemistry: Preparation and properties of soluble antigen-antibody complex (horseradish peroxidase-antihorseradish peroxidase) and its use in identification of spirochetes. *J. Histochem. Cytochem.*, 18: 315 (1970).
 20. Sutton, J. and Weiss, L.: Transformation of monocytes in tissue culture into macrophages, epithelioid cells, and multinucleated giant cells. *J. Cell. Biol.*, 28: 303 (1966).
 21. Williams, C. A. and Chase, M. W.: Production of antiserum. In: *Methods in Immunology and Immunochemistry*, Vol. 1, Preparation of Antigens and Antibodies. pp. 197-306 (Academic Press, New York, 1967).
 22. Requests for reprints should be addressed to: Dr. J. M. Andres, Dept. of Pediatrics (Gastroenterology), Box J-296, J. Hillis Miller Health Center, University of Florida College of Medicine, Gainesville, FL 32610.
 23. Supported in part by a University of Florida Sponsored Research Award.
 24. Received for publication October 4, 1982.
 25. Accepted for publication May 12, 1983.

Letter to the Editor

H. M. ROBINSON

Tropical Metabolism Research Unit, University of the West Indies, Kingston, Jamaica

The excellent review by Brasel (1), "Endocrine Adaptation to Malnutrition" is a thorough attempt to look for generalities in a confusing field. It is possible that one of the points of confusion Brasel leaves unresolved contains within it information that may help to reduce the confusion, and suggests a need for greater liaison between investigators.

Brasel points out that studies on growth hormone in malnutrition generally report high levels of this hormone, with exceptions, in cases where growth hormone levels were originally normal or diminished. The studies of Monckeberg *et al.* (2) and Robinson *et al.* (3) were cited. Both of these studies do more than this. Robinson *et al.* finds abnormally high growth hormone levels in malnourished children showing zero or negative weight gain. These levels are 2-5 times higher than during dietary treatment. Dietary treatment initiates the recovery or catch-up growth phase. Both Monckeberg *et al.* and Robinson *et al.* also report on growth hormone levels in children being treated for existing malnutrition.

Similar reports have been given by Beas *et al.* (4) and Godard (5). They point out that during treatment and/or recovery from protein-energy malnutrition, lower growth hormone values were found in children in whom growth rate was very slow [Robinson *et al.* (3) and Godard (5)] or in those in whom recovery was not yet initiated [Monckeberg *et al.* (2) Beas *et al.* (4)] and higher growth hormone levels were found in those showing satisfactory growth rates [Robinson *et al.* (3) and Godard (5)]. The administration of growth hormone to the children who were not growing significantly increased growth rates [Monckeberg *et al.* (2)]. At no time during the catch-up growth phase [Robinson *et al.* (3)] did the growth hormone levels approach the abnormally high values shown in the malnourished group.

Malnutrition is a state defined anthropometrically. As such it is a measure of an anthropometric state achieved and not the rate of change of nutritional status or growth. The results of Monckeberg *et al.* (2), Robinson *et al.* (3), Beas *et al.* (4) and Godard (5) indicate that this rate of change, shown during recovery, is significantly related to growth hormone levels.

It would help in the interpretation of all results on malnourished

children if clinical status could be specified more precisely than is possible by currently accepted anthropometric classification schemes, or by traditional clinical divisions into named disease states. It would be illogical to ask that only growth rates should be given, although they are clearly important. It is also reasonable to assume that food intakes would be important determinants of hormonal profile, and also that patients can be not only recovering, but if studied early enough after the initiation of malnutrition, their hormonal profile may also reflect their rate of deterioration.

It is difficult to see how coherence can be obtained. Is it unreasonable to suggest a discussion among principal investigators is needed to agree on an ideal list of clinical data for any malnourished child studied, and to propose the requirement for a single archive where copies of such new data could be lodged by all investigators, and made accessible to all workers in the field, permitting more easily the detection of trends of the type reported by Monckeberg *et al.* (2), Beas *et al.* (4), Godard (5), and Robinson *et al.* (3)?

REFERENCES AND NOTES

1. Brasel, J.: *Pediatr. Res.*, 14: 1299 (1980).
2. Monckeberg, F., Donoso, G., Oxman, S., Pak, N., and Meneghello, J.: *Pediatrics*, 31: 58 (1963).
3. Robinson, H., Cocks, T., Kerr, D., and Picou, D. In: 'Endocrine Aspects of Malnutrition'. Ed. L. I. Gardner and P. Amacher, 1973, 45-72.
4. Beas, F., Contreras, I., Maccioni, A., and Arenas, S.: *Br. J. Nutr.*, 26: 169 (1971).
5. Godard, C.: In: 'Endocrine Aspects of Malnutrition'. Ed. L. I. Gardner and P. Amacher, 1973, 19-30.

Response

JO ANNE BRASEL

Harbor-UCLA Medical Center, UCLA School of Medicine, Torrance, California

I completely agree with Dr. Robinson that much of the difficulty in evaluating the data on growth hormone levels in malnutrition is related to the fact that the subjects are often in differing states of malnutrition and/or recovery. It would have been very helpful to me in my review if the types of additional information on the patients and their condition, which is mentioned in the letter, had been available. I can only support the proposal that investigators in the field develop a format for recording such data for the use of others with great enthusiasm. Good luck!