

- head of bacteriophage T4. *Nature*, 227: 680 (1970).
- Burnette, W. N.: Western blotting: electrophoretic transfer of proteins from sodiumdodecylsulfate-polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein A. *Analyt. Biochem.*, 112: 195 (1981).
 - Pinto, M., Robine-Leon, S., Appay, M. D., Keding, M., Triadou, N., Dus-saulx, E., Lacroix, B., Simon-Assmann, P., Haffen, K., Fogh, J., and Zwi-baum, A.: Enterocyte-like differentiation of a human colon malignant epi-thelial cell line (Caco-2) grown in standard culture conditions. *Bio Cell*, 47: 323 (1983).

- Auricchio, S., Caporale, C., and Santamaria, F.: Human fetal intestinal brush border sucrase, oligoaminopeptidase and dipeptidylaminopeptidase. *Pediatr. Res.*, 15: 1193 (1981).
- Stamatoyannopoulos, G. and Nienhuis, A. W.: Cellular and molecular regu-lation of hemoglobin switching. (Grune and Stratton, New York, 1979).
- Whalen, R. G., Butler-Browne, G. S., Sell, S., and Gros, F.: Transitions in contractile protein isozymes during muscle cell differentiation. *Biochimie*, 61: 625 (1979).
- Triadou, N. and Audran, E.: Interaction of the brush-border hydrolases of the human small intestine with lectins. *Digestion*, 27: 1 (1983).

Letter to the Editor

EDWARD F. BELL, EKHARD E. ZIEGLER, AND GILBERT B. FORBES

Department of Pediatrics, University of Iowa, Iowa City, Iowa, USA,
and Department of Pediatrics, University of Rochester, Rochester,
New York, USA

The volume of distribution of bromide is often used to estimate the volume of extracellular water in the body. The "corrected" bromide space (CBS) is assumed to equal the volume of extra-cellular water. To calculate the CBS, the dose of bromide (Br dose) is divided by the equilibrium concentration of bromide in extracellular fluid ($[Br]_e$), with a correction for the amount that is assumed to be distributed in non-extracellular sites (principally erythrocytes):

$$CBS = \frac{Br \text{ dose} \times 0.90}{[Br]_e}$$

Because $[Br]_e$ cannot be measured directly, it is calculated from the concentration of bromide in serum ($[Br]_s$) by applying two additional correction factors: the Donnan equilibrium factor 0.95 (8, 11, 12), and the concentration of water in serum (7, 8, 12), which is approximately 94% (12). Thus,

$$[Br]_e = \frac{[Br]_s}{0.95 \times 0.94}$$

and, therefore

$$CBS = \frac{Br \text{ dose}}{[Br]_s} \times 0.90 \times 0.95 \times 0.94$$

These corrections are not needed when one uses bromide to estimate total body chloride (10) because the Donnan factors for these two elements are the same, and both are measured in the same serum sample.

Several authors (1, 4, 7, 9, 13, 14) have applied one or two of these three factors incorrectly, multiplying rather than dividing $[Br]_s$ by the Donnan factor and the serum water correction. The result in most cases has been a large, systematic overestimation of extracellular water. When both the Donnan factor and the serum water correction are used incorrectly (1, 4, 9, 13, 14) the resulting errors are additive; extracellular water is then system-atically overestimated by 25–31%, depending on the factor used for the serum water correction (0.92–0.94). It should be noted, parenthetically, that even with correct application of these factors there is a slight overestimation of the bromide space if no correction is made for bromide lost in urine before the serum sample is obtained.

Table 1 shows the corrected bromide space results for infants as reported (1, 4, 9, 13, 14) and the values that would have been obtained with correct application of the Donnan and serum water factors. The mean extracellular water volume calculated by Brans *et al.* (1) was 335 ml/kg, higher than the "expected" volume of 286 ml/kg (ref. 1, Table 3). They concluded that "infants with congenital heart disease appear to have . . . slightly

Table 1. Corrected bromide space (CBS) of infants

Authors	Subjects	Mean CBS as reported (ml/kg)	Correctly calculated* mean CBS (ml/kg)
Cassady (4)	TermAGA†	376	296
	Premature AGA†	424	334
Brans <i>et al.</i> (1)	Infants with CHD‡	335	264
	Thornton <i>et al.</i> (13)	Term	504
Thornton <i>et al.</i> (14)	Premature	640	504
	Normocythemic	510	402
	Polycythemic	520	409
Cheek <i>et al.</i> (9)	TermV§	358	273
	TermC	354	270

* Using same Donnan factor and serum water correction used by original authors: Donnan factor 0.95 (1, 4, 9, 13, 14); serum water factor 0.934 (1, 4, 13, 14) or 0.92 (9).

† AGA, appropriate birth weight for gestational age, less than 24-h-old.

‡ CHD, congenital heart disease.

§ V, vaginally delivered, 24-h-old.

|| C, delivered by cesarean section, 24-h-old.

expanded volumes of extracellular water" (1). This conclusion seems to have been drawn erroneously as a result of the afore-mentioned errors in correcting the bromide space (unless the "expected" volume was also calculated incorrectly).

The extent of these errors is not known because many authors report only that corrections were made for the Donnan equilib-rium and serum water, without giving details of the calculations. It must be presumed, however, that other papers (2, 3, 5, 6) citing the method of Cassady (4) or Thornton *et al.* (14) also contain bromide space results that were incorrectly calculated. In some cases extracellular water was incorrectly estimated from bromide space and then subtracted from total body water to yield intracellular water (1, 2, 5, 6, 9, 14). This, of course, led to underestimation of intracellular water by the same amount. For example, the mean intracellular water of 328 ml/kg reported by Brans *et al.* (1) should have been 399 ml/kg.

As far as we know, the most recent experimental paper to correctly use and describe the three corrections for bromide space was published in 1954 (8). Most of our information regarding the extracellular water of infants is based on bromide space measurements. Many of these data are demonstrably wrong (1, 7, 9, 13, 14); others cannot be judged because the details of bromide space correction were not given. We conclude that infants contain less extracellular water than was previously thought.

REFERENCES AND NOTES

- Brans, Y. W., Dweck, H. S., Harris, H. B., Parr, G. V. S., Bailey, P. E., Kirklín, J. W., and Cassady, G.: Effect of open-heart surgery on the body composition of infants and young children. *Pediatr. Res.*, 15: 1024 (1981).

2. Brans, Y. W., Shannon, D. L., and Hunter, M. A.: Maternal diabetes and neonatal macrosomia. III. Neonatal body water estimates. *Early Hum. Dev.*, 8: 307 (1983).
3. Brans, Y. W., Sumners, J. E., Dweck, H. S., Bailey, P. E., and Cassady, G.: Feeding the low-birthweight infant: orally or parenterally? II. Corrected bromide space in parenterally supplemented infants. *Pediatrics*, 58: 809 (1976).
4. Cassady, G.: Bromide space studies in infants of low birth weight. *Pediatr. Res.*, 4: 14 (1970).
5. Cassady, G.: Effect of cesarean section on neonatal body water spaces. *N. Engl. J. Med.*, 285: 887 (1971).
6. Cassady, G. and Milstead, R. R.: Antipyrine space studies and cell water estimates in infants of low birth weight. *Pediatr. Res.*, 5: 673 (1971).
7. Cheek, D. B.: Estimation of the bromide space with a modification of Conway's method. *J. Appl. Physiol.*, 5: 639 (1953).
8. Cheek, D. B.: Observations on total body chloride in children. *Pediatrics*, 14: 5 (1954).
9. Cheek, D. B., Wishart, J., MacLennan, A. H., Haslam, R., and Fitzgerald, A.: Hydration in the first 24 h of postnatal life in normal infants born vaginally or by caesarean section. *Early Hum. Dev.*, 7: 323 (1982).
10. Forbes, G. B., Reid, A. F., Bondurant, J., and Etheridge, J.: Changes in total body chloride during growth. *Pediatrics*, 17: 334 (1956).
11. Gamble, J. L., Jr. and Robertson, J. S.: Volume of distribution of radioactive chloride in dogs; comparisons with sodium, bromide and inulin spaces. *Am. J. Physiol.*, 171: 659 (1952).
12. O'Brien, D., Ibbott, F. A., and Rodgerson, D.: *Laboratory Manual of Pediatric Micro-Biochemical Techniques*. 4th ed. pp. 78-80 (Harper and Row, New York, 1968).
13. Thornton, C. J., Shannon, D. L., Hunter, M. A., and Brans, Y. W.: Dynamic skinfold thickness measurements: a noninvasive estimate of neonatal extracellular water content. *Pediatr. Res.*, 16: 989 (1982).
14. Thornton, C. J., Shannon, D. L., Hunter, M. A., Ramamurthy, R. S., and Brans, Y. W.: Body water estimates in neonatal polycythemia. *J. Pediatr.*, 102: 113 (1983).

Response

YVES W. BRANS, GEORGE CASSADY, AND DONALD B. CHEEK

Perinatal Research Laboratory, Departments of Pediatrics and of Obstetrics and Gynecology, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA; Perinatal Research Laboratory and Division of Perinatal Medicine, Department of Pediatrics, University of Alabama in Birmingham, Birmingham, Alabama, USA; and Department of Obstetrics and Gynecology, The University of Adelaide, Adelaide, South Australia

Doctors Bell, Ziegler, and Forbes must be congratulated for realizing that over a period of 30 yr the various factors used to correct the bromide space have been at times *apparently* misapplied (3). The correct equation should be written as follows:

Corrected bromide space

$$= \frac{\text{Amount of bromide administered}}{\text{Bromide concentration in plasma}} \times 0.90 \times 0.95 \times 0.94$$

where 0.90 corrects for the amount of intracellular bromide; 0.95 corrects for the Donnan equilibrium; and 0.94 corrects for the proportion of water in plasma.

Their letter triggered an extensive review of data in our three laboratories. Dr. Cassady recalculated the data from all studies performed in his laboratory (1, 2, 4-6) and found that the correct formula was indeed used. Dr. Cheek also used that formula in all his work (7-9). Unfortunately, typographic errors in the "Materials and Methods" sections of several articles suggested that some of the correction factors were misapplied (1, 4, 6, 9). Dr. Bell and his colleagues are therefore incorrect in concluding that "infants contain less extracellular water than was previously thought" and the "corrected" values given in his table are in fact erroneous. In Dr. Brans's laboratory, however, the incorrect formula reported by Cassady (4) and Cheek (9) was used:

Corrected bromide space

$$= \frac{\text{Amount of bromide administered}}{\text{Bromide concentration in plasma}} \times \frac{0.90}{0.95 \times 0.94}$$

An overestimation of both the corrected bromide space (CBS) and the interstitial water (IW) resulted (10). Because the same error was applied to the two study groups being compared (preterm *versus* term neonates), the overestimation did not affect the comparisons. Likewise, in the regression analysis, the errors

affected the slope and y-intercept of the regression lines, but not the correlation coefficient nor the $S_{y \cdot x}$. For the record, however, corrected values are as follows:

CBS (ml/kg): 400 ± 67.0 in term *versus* 501 ± 61.0 in preterm babies

IW (ml/kg): 352 ± 63.0 in term *versus* 449 ± 59.4 in preterm babies

$\% \Delta \text{MT} = 0.064 \text{ CBS} - 1.3$

$\% \Delta \text{SS} = 0.081 \text{ CBS} - 7.2$

$\Delta \text{AMT} \times 10^4 = 0.087 \text{ CBS} - 5.8$

$\Delta \text{ASS} \times 10^4 = 0.077 \text{ CBS} + 0.2$

$\% \Delta \text{MT} = 0.066 \text{ IW} + 1.0$

$\% \Delta \text{SS} = 0.083 \text{ IW} - 4.0$

$\Delta \text{AMT} \times 10^4 = 0.090 \text{ IW} - 2.7$

$\Delta \text{ASS} \times 10^4 = 0.080 \text{ IW} + 3.0$

These corrections do not affect the validity of the conclusions of the study.

We cannot agree that significant amounts of bromide are lost in the urine during the 3 h of the study. In our experience urine losses amount to less than 1% of the administered dose of bromide. Finally, Bell *et al.*'s conclusion that "infants contain less extracellular water than was previously thought" is not supported by the facts.

REFERENCES AND NOTES

1. Brans, Y. W., Dweck, H. S., Harris, H. B., Parr, G. V. S., Bailey, P. E., Kirklín, J. W., and Cassady, G.: Effect of open-heart surgery on the body composition of infants and young children. *Pediatr. Res.*, 15: 1024 (1981).
2. Brans, Y. W., Sumners, J. E., Dweck, H. S., Bailey, P. E., and Cassady, G.: Feeding the low-birthweight infant: orally or parenterally? II. Corrected bromide space in parenterally-supplemented infants. *Pediatrics*, 58: 809 (1976).
3. Bell, E. F., Ziegler, E. F., and Forbes, G. B.: Errors in bromide space correction (letter to the editor). *Pediatr. Res.*, (in press).
4. Cassady, G.: Bromide space studies in infants of low birth weight. *Pediatr. Res.*, 4: 14 (1970).
5. Cassady, G.: Effect of cesarean section on neonatal body water spaces. *N. Engl. J. Med.*, 285: 887 (1971).
6. Cassady, G. and Milstead, R. R.: Antipyrine space studies and cell water estimates in infants of low birth weight. *Pediatr. Res.*, 5: 673 (1971).
7. Cheek, D. B.: Estimation of the bromide space with a modification of Conway's method. *J. Appl. Physiol.*, 5: 639 (1953).
8. Cheek, D. B.: Observations of total body chloride in children. *Pediatrics*, 14: 5 (1954).
9. Cheek, D. B., Wishart, J., MacLennan, A. H., Haslam, R., and Fitzgerald, A.: Hydration in the first 24 h of postnatal life in normal infants born vaginally or by caesarean section. *Early Hum. Dev.*, 7: 323 (1982).
10. Thornton, C. J., Shannon, D. L., Hunter, M. A., and Brans, Y. W.: Dynamic skinfold thickness measurements: a non-invasive estimate of neonatal extracellular water content. *Pediatr. Res.*, 16: 989 (1982).