R Fein, W THAL, H-J Otto, M Gunkel

Dept. Paed. Med. Akad. Magdeburg, GDR

The combined ventilation and perfusion scanning with ^{133}Xe -gas and ^{99m}Tc Human albumin microspheres was carried out in 133 children at the age from 1 month to 16 years. The findings were compared with clinical x-ray and bronchologic ones as well as with parameters for the function of the lung FVC and FEV₁.
In children with bronchiectasia ($n = 24$) there exist distinct perfusion and exhale disturbances in form of trapped gas, reduced exhalation and half value time of the ^{133}Xe -clearance in the bronchiectasial areas, proved bronchographically.
In children with deforming bronchitis ($n = 18$) we only observed slight perfusion disturbances and twice discreet exhale troubles without trapped gas.
In children with cystic fibrosis ($n = 26$) regional obstructions could prematurely be proved, they were not yet caught spirometrically. Typical findings were diffuse perfusion- and ventilation disturbances (trapped gas) mainly in the apex areas.
With agenesis of A. pulmonalis we observed no perfusion and reduced FRC of the side affected with no abnormality in exhaling, as well as in case with agenesis of the lung an absent perfusion and ventilation.

W Dorsch and M Ettl

Kinderpoliklinik of the University, Munich

Late bronchial reactions (LBR) of neonatal immunized rabbits upon allergen inhalation challenges represent an important animal model of human LBR (Am.Rev.Resp.Dis.126:493,1982). Previously we described "compressed air" (CA) as sensitive parameter of bronchial obstruction in guinea pigs (Pflügers Arch.391:236,1981). Now we present a method for measuring CA and airway resistance (Raw) in rabbits.

Both compartments of a body plethysmograph (head chamber = HC and body chamber = BC) are separated by a water filled rubber cuff around the head of the animal. In both chambers there are identical gas conditions with respect to volume, humidity (100%) and temperature (37°C). Volume changes in HC and BC are measured by pressure transducers and plotted on an XY recorder. From this graph CA can be read off as described previously. Raw is determined as follows: Respiratory flow is calculated by differentiation of HC volume changes with time (dV/dt) and compared with pressure changes in both chambers ($P_{\text{tot}} = P_{\text{HC}} + P_{\text{BC}}$). The resulting graph is identical to the well known α -curve of one chamber plethysmography. The ratio between P_{tot} and alveolar pressure (" β curve") is determined by a special occlusion technique.

Both parameters, CA and Raw, are certainly useful for studying LBR in an animal model. They can be obtained by noninvasive methods.

J.C. de Jongste, E.J. Duiverman, H.J. Neijens, K.F.

Kerrebijn. Erasmus University Rotterdam/Sophia Children's Hospital - Dept. of Pediatric Respiratory Disease.

A 10-year old girl presented with deep cyanosis, right cardiac failure and generalised seizures. The medical history was unremarkable except for neonatal meconium aspiration and a generalised seizure at the age of 4. After treatment of the acute phase, the diagnosis of primary alveolar hypoventilation was established by the following findings: 1. Persistent hypoxia (SaO_2 40-100%) and hypercapnia (pCO_2 60-100 mm Hg), especially during quiet sleep, with prolonged episodes of apnea; 2. impaired ventilatory CO_2 -response; 3. normal pulmonary function without airway obstruction; 4. no signs of myopathy with normal EMG and diaphragmatic EMG; 5. improvement of ventilation during infusion of almitrine (a chemoreceptor agonist). There were no abnormalities on 24-hours EEG registration. Additional findings were right ventricular hypertrophy with pulmonary hypertension and mental retardation with cerebral atrophy on CT scan. Both were probably the consequence of prolonged hypoxia. Maintenance therapy included doxapram, almitrine and caffeine together with oxygen during sleep. Because this medication was ineffective in preventing nocturnal apnea, mechanical ventilatory assistance was instituted during sleep by means of an external cuirass ventilator. At present the patient has been treated for 18 months with caffeine and cuirass ventilation at night. No complications were seen; the therapy proved effective in preventing nocturnal hypoxia and resulted in a marked improvement in mental status, a normal growth and pubertal development and normalisation of cardiac dimensions.

F.Midulla, F.Macri, G.Ciofetta, V.Colloridi, A.Ceccamea, R.Ronchetti. V Clin Ped, Rome University "La Sapienza".

Up to now 400 cases of PAVF have been described. The PAVF are most often found isolated, mainly in the left inferior lobe. In only the 27% of the cases the cyanosis starts in the neonatal period. We describe a 4 month-old child with diffuse, bilateral PAVF, who manifested cyanosis in the first month of life. At birth there were no signs of respiratory distress. Since the first month of age the mother noticed peripheral cyanosis. On admission she was in good general conditions with an evident peripheral cyanosis. Routine examinations showed only a mild absolute polycythemia. At rest PaO_2 was 41 mmHg, PaCO_2 34 mmHg and pH 7.4. With the hyperoxo test was calculated a right to left shunt of 60%. The cardiac catheterization showed that the PaO_2 in the pulmonary veins was around 40 mmHg and that there were no cardiac malformations; the injection of Tc99 labelled albumin macroaggregates demonstrated the passage of a considerable amount of particles through the pulmonary circulation and the pulmonary angiography showed the injection of multiple small vessels placed in the peripheral coat of lung parenchyma. After death lung pathology confirmed the presence of multiple pulm vascular lacunae.

T. LOÓS

Dept. of Pediatric Pneumology of County Hospital Nyíregyháza 4412 (Hungary)

A 2-year-old boy was sent to our hospital because of repeated cough and chronic pneumonia in the left lung. On his chest X-ray picture there is a homogeneous shadow left behind the heart. At the bronchoscopic investigation, signs of chronic bronchitis were found. On the left bronchography there are two different abnormalities. On the one hand the alteration behind the heart (which was considered a chronic pneumonia) crushes the bronchi of the left lower lobe and on the other hand there is a congenital malformation too, namely the lingula inferior bronchus arises separately between the upper and the lower lobe bronchi. In addition we found other congenital disorders too in the same child. In newborn age he underwent a surgical reconstruction on a left hernia diaphragmatica. The shadow behind the heart was caused not by a chronic pneumonia, but by a left thoracic kidney. His stomach was also irregularly located, not on the left side, but to the right of the middle line. Finally we found klinodaktyly on both thumbs.

EPRS—Abstract Read by Title

J.C. de Jongste, E.J. Duiverman, H.J. Neijens, K.F.

Kerrebijn. Erasmus University Rotterdam/Sophia Children's Hospital - Dept. of Pediatric Respiratory Disease.

A 10-year old girl presented with deep cyanosis, right cardiac failure and generalised seizures. The medical history was unremarkable except for neonatal meconium aspiration and a generalised seizure at the age of 4. After treatment of the acute phase, the diagnosis of primary alveolar hypoventilation was established by the following findings: 1. Persistent hypoxia (SaO_2 40-100%) and hypercapnia (pCO_2 60-100 mm Hg), especially during quiet sleep, with prolonged episodes of apnea; 2. impaired ventilatory CO_2 -response; 3. normal pulmonary function without airway obstruction; 4. no signs of myopathy with normal EMG and diaphragmatic EMG; 5. improvement of ventilation during infusion of almitrine (a chemoreceptor agonist). There were no abnormalities on 24-hours EEG registration. Additional findings were right ventricular hypertrophy with pulmonary hypertension and mental retardation with cerebral atrophy on CT scan. Both were probably the consequence of prolonged hypoxia. Maintenance therapy included doxapram, almitrine and caffeine together with oxygen during sleep. Because this medication was ineffective in preventing nocturnal apnea, mechanical ventilatory assistance was instituted during sleep by means of an external cuirass ventilator. At present the patient has been treated for 18 months with caffeine and cuirass ventilation at night. No complications were seen; the therapy proved effective in preventing nocturnal hypoxia and resulted in a marked improvement in mental status, a normal growth and pubertal development and normalisation of cardiac dimensions.

J Storr, N A P Evans, W Lenney

Royal Alexandra Hospital for Sick Children, Brighton, England

There is little evidence that any medication is helpful in acute viral croup but few studies have attempted objective measurement of the stridor before and after treatment. We have devised a small microphone which can be easily attached to the child's neck by double sided adhesive tape. The microphone predominantly responds to vibration and is unresponsive to external sounds. Tape recordings were made before and 15 minutes after nebulised therapy. The response to nebulised adrenaline, ipratropium bromide, xylometazoline, clemastine and placebo were observed in a double blind manner. Inspiratory and expiratory components were separated by listening to the tape whilst concurrently viewing the output on a BBC micro-computer video display unit (VDU). The measurement was expressed by integrating the sound amplitude over the period of inspiratory noise. There was little response to any medication in children under 2 years of age but the response increased with age. Improvement in the microphone trace correlated well with clinical improvement. Conclusion - We present evidence to show that medications (especially adrenaline) can be helpful in acute viral croup in children over 2 years of age.