

# Postheparin Plasma Lipoprotein and Hepatic Lipase Activities in Hyperinsulinemic Infants of Diabetic Mothers and in Large-for-Date Infants at Birth

LIISA M. ROVAMO, MARJA-RIITTA TASKINEN, TIMO KUUSI, AND KARI O. RAIVIO

*Children's Hospital, Second and Third Departments of Medicine, and Department of Obstetrics and Gynecology, University of Helsinki, Helsinki, Finland*

**ABSTRACT.** To study postheparin plasma lipase activities in nonfed newborn infants immediately after birth and to investigate the possible influence of fetal hyperinsulinemia on lipoprotein lipase activity, we measured lipoprotein and hepatic lipase activities in 55 macrosomic newborn infants: group I consisted of 21 infants born to mothers with insulin-dependent diabetes. The infants were hyperinsulinemic at birth and had hypoglycemia and poor lipolysis at the age of 2 h. Group II consisted of 18 infants born to mothers with gestational diabetes. Group III consisted of 16 large-for-date infants born to nondiabetic mothers. The mean postheparin plasma lipoprotein lipase activities at 2 h of age were similar (mean 36  $\mu\text{mol}$  free fatty acids/ml/h; SEM 15) in groups I-III. Lipoprotein lipase activity correlated negatively with cord-serum triglycerides (range 0.13-1.2 mmol/liter) but did not correlate with serum insulin (range 5.4-524  $\mu\text{U/ml}$ ) or C-peptide (range 0.6-21.0  $\mu\text{g/liter}$ ). Hepatic lipase activity was somewhat higher in group I (mean 68  $\mu\text{mol}$  free fatty acids/ml/h; SEM 23) than in groups II and III (mean 55  $\mu\text{mol}$  free fatty acids/ml/h; SEM 14). Hemoglobin Alc was the only important factor explaining the difference in hepatic lipase activities between groups. Lipoproteins and apolipoproteins A-I, A-II, and B were similar in all three groups. We conclude that in large-for-date infants lipoprotein lipase is active at birth without exogenous fat induction, and that these infants are capable of hydrolyzing fat, their main source of energy, immediately after birth. In addition, we conclude that postheparin plasma lipoprotein lipase activity is not affected by fetal hyperinsulinemia. (*Pediatr Res* 20: 527-531, 1986)

## Abbreviations

FFA, free fatty acids  
Hb, hemoglobin

Soon after birth fat becomes the major metabolic fuel (1): breast-milk fat is readily absorbed through the bowel wall as chylomicrons and liver produces endogenous lipoproteins into

Received July 17, 1985; accepted January 30, 1986.

Reprint requests Liisa Rovamo, M.D., Children's Hospital, University of Helsinki, Stenbäckinkatu 11, SF-00290 Helsinki, Finland.

Supported by The Foundation for Pediatric Research, Heikki and Hilma Honkanen's Foundation, Foundation of Nutrition Research, Finnish Cultural Foundation, Sigrid Juselius Foundation, The Academy of Finland, and Huhtamäki Inc., Leiras Pharmaceuticals.

the circulation (2). Triglycerides in chylomicrons and very low-density lipoproteins of plasma are hydrolyzed by lipoprotein lipase (3, 4). The lipase is located on the vascular endothelial surface of adipose and muscle tissue. It is released into the circulation by heparin. Heparin also releases hepatic lipase from the vascular endothelial surface of the liver. In contrast to lipoprotein lipase, the function of hepatic lipase in lipoprotein metabolism has not been fully established (4, 5).

On the basis of animal experiments, lipoprotein lipase activity of newborn infants has been suggested to be low at birth and to increase postnatally (1, 6, 7). If so, infants at birth might be incapable of using fat effectively for energy. Previously, we have shown that at the age of 1 to 4 days postheparin plasma lipoprotein lipase is already at the adult level in breast-milk-fed term and preterm infants; their hepatic lipase activity is about three times the adult value (8, 9). On the other hand, we found that lipoprotein lipase activity increases during the first days of life whereas hepatic lipase activity remains unchanged in infants receiving fat-containing parenteral nutrition (10).

Fetal hyperinsulinemia is associated with macrosomia, hypoglycemia, and inhibition of lipolysis in infants born to diabetic mothers (11). In adults, insulin is known to induce lipoprotein lipase (3, 12), and lipoprotein lipase activity of adipose tissue has been reported to be high in obese adults whose basal serum insulin level is elevated (13, 14). This suggests that fetal hyperinsulinemia associated with maternal diabetes mellitus may induce lipoprotein lipase activity perinatally.

To investigate the levels of postheparin plasma lipases immediately after birth in large-for-date infants and to study the possible inducing effect of intrauterine insulin on lipoprotein lipase, we measured lipoprotein and hepatic lipase activities in hyperinsulinemic infants born to diabetic mothers, in infants born to gestationally diabetic mothers, and in large-for-date infants born to nondiabetic mothers.

## MATERIALS AND METHODS

**Subjects.** A total of 55 infants were studied. They belonged to one of the following three groups:

Group I consisted of 21 infants born to insulin-dependent diabetic mothers (three of White's class A/B (15), eight of class B, six of class C, and four of class D). The antenatal and obstetric care of the mothers was as previously described (16-18). During the last trimester of pregnancy the mean Hb Alc was 6.7% of total hemoglobin [range 5.6-8.9% Hb; mean + SD for normal pregnant women in the 3rd trimester is in our laboratory 5.0 + 0.5% Hb (19)]. During the week preceding delivery the mean Hb Alc was 6.4% Hb (range 5.0-8.2% Hb).

After ascertainment of lung maturity by amniocentesis (20),

labor was induced in eight mothers whereas 13 mothers underwent elective cesarean section. On the morning of intended delivery, the mothers received 12 U of short-acting insulin subcutaneously, followed by a glucose infusion at a rate of 4 mg/kg/min; subsequent doses of insulin were adjusted to maintain normoglycemia.

Gestational ages of 14 boys and 7 girls ranged from 34 to 38 wk (mean 37.6), birth weights from 2660 to 4780 g (mean 3891 g), and relative birth weights from -0.5 to 4.5 SD units (mean 2.2). Relative birth weight refers to the deviation of the individual birth weight from the mean birth weight for gestational age (21) divided by the corresponding SD (21). One infant had an Apgar score of 3 at 1 min but 7 at 5 min of age; all others scored from 7 to 10 at 1 min of age.

Group II consisted of 18 infants born to mothers with gestational diabetes: blood glucose concentration exceeded 140 mg/dl at 2 h in an oral glucose (50 g) tolerance test. The mothers were treated by diet alone. At the time of delivery the mean Hb Alc was 5.4% of total Hb (range 4.7-6.0% Hb). During delivery the mothers received glucose infusion at a rate of 4 mg/kg/min but no insulin. Sixteen infants were born through the vaginal route and two by cesarean section. Gestational age of 12 boys and six girls ranged from 37 to 42 wk (mean 39.4), birth weights from 3140 to 4670 g (mean 3820 g), and relative birth weights from -0.3 to 2.7 SD units (mean 1.1). Apgar scores ranged from 7 to 10 at 1 min of age.

Group III consisted of 16 large-for-date infants born to mothers whose oral glucose tolerance tests were normal. At the time of

delivery the mean Hb Alc was 5.7% of total Hb (range 4.7-6.5% Hb). Twelve of the mothers received glucose infusion at a rate of 0.2-2.5 mg/kg/min during delivery. Thirteen infants were born through the vaginal route and three by cesarean section. Gestational ages of nine boys and seven girls ranged from 39 to 42 wk (mean 41.2), birth weights from 4500 to 5970 g (mean 4811 g), and relative birth weights from 2.0 to 5.8 SD units (mean 2.7). Apgar scores ranged from 7 to 10 at 1 min of age.

The purpose of the investigation was explained to both parents and the investigation was carried out with their consent. The experimental protocol was approved of by the Ethical Committee, Department of Obstetrics and Gynecology, University of Helsinki. Ethical considerations precluded the study of infants with normal birth weights born to nondiabetic mothers.

*Experimental protocol.* Immediately after birth a sample of mixed cord blood was deproteinized for the measurement of glucose and  $\beta$ -hydroxybutyrate concentrations (groups I and II). Another sample of cord blood was centrifuged at +4° C. Part of the serum was stored at -20° C for the assays of insulin and C-peptide, FFA, and apolipoproteins. The rest of the serum was ultracentrifuged for the separation of the lipoprotein classes.

During the first hours of life the infants were under careful observation: blood pressure, respiratory rate, heart rate, and heel and rectal temperatures were recorded every 10 min; all infants remained in good condition.

At the age of 2 h, a blood sample was taken through an indwelling peripheral vein catheter for the measurement of blood glucose and  $\beta$ -hydroxybutyrate, serum insulin or free insulin,

Table 1. Hormonal and metabolic parameters at birth and 2 h of age: mean (range)

	n	Cord blood	n	2 h	p value*
Serum C-peptide ( $\mu$ g/liter)					
Group I	20	5.9 (1.1-21)			
Group II	18	2.0 (0.7-4.7)			
Group III	16	1.8 (0.6-3.7)			
p value†		$p < 0.001$			
Serum insulin ( $\mu$ U/liter)					
Group I (free insulin)	21	120 (7.2-520)	19	26 (0.7-77)	$p < 0.001$
Group II	18	30 (5.4-100)	15	13 (2.9-30)	$p < 0.05$
Group III	16	27 (7.5-100)	16	29 (4.5-130)	NS
p value†		$p < 0.001$		$p < 0.05$	
Blood glucose (mmol/liter)					
Group I	18	5.9 (2.2-12)	20	1.2 (0.5-2.5)	$p < 0.0001$
Group II	14	5.9 (1.8-9.5)	18	2.1 (1.1-3.0)	$p < 0.0001$
Group III			16	2.1 (1.3-3.0)	
p value†		NS		$p < 0.0001$	
Serum triglycerides (mmol/liter)					
Group I	21	0.41 (0.13-1.2)	19	0.59 (0.36-0.88)	$p < 0.05$
Group II	18	0.41 (0.22-0.65)	18	0.63 (0.14-0.97)	$p < 0.05$
Group III	15	0.39 (0.21-0.58)	16	0.60 (0.18-0.84)	$p < 0.05$
p value†		NS		NS	
Serum FFA (mmol/liter)					
Group I	21	0.20 (0.06-0.58)	20	0.27 (0.05-0.82)	NS
Group II	18	0.18 (0.07-0.28)	18	0.70 (0.13-1.7)	$p < 0.001$
Group III	16	0.22 (0.10-0.41)	16	0.77 (0.20-1.7)	$p < 0.001$
p value†		NS		$p < 0.0001$	
Blood $\beta$ -hydroxybutyrate (mmol/liter)					
Group I	16	0.80 (0.13-3.2)	20	0.21 (0.41-0.39)	$p < 0.05$
Group II	14	0.63 (0.21-1.5)	17	0.21 (0.14-0.27)	$p < 0.05$
Group III			16	0.24 (0.18-0.34)	
p value†		NS		NS	

\* Comparing cord-blood and 2-h values.

† Comparing groups.

FFA, and triglycerides, and plasma basal activities of lipoprotein and hepatic lipases. Thereafter, a heparin (Medica, Helsinki, Finland) bolus of 100 IU/kg was injected through the catheter. Fifteen minutes later a blood sample was taken for the measurement of lipase activities. Then the infants received their first feeding of breast milk. As shown previously (8–10), the infants showed no signs of bleeding tendency or other side-effects resulting from heparin administration. Five infants of group I and one of group III received intravenous glucose because of hypoglycemia during the first days of life.

**Analytical methods.** Serum C-peptide, serum insulin, and free insulin were measured with radioimmunoassay methods (22, 23). Serum FFA concentration was measured with an enzymatic method (Nefa C-Test, Wako, Wako Pure Chemical Industries LTD, Japan). Blood  $\beta$ -hydroxybutyric acid concentration was measured with a fluorometric method (24).

Separation of serum lipoproteins was done using a Ti-50.3 rotor in a Beckman L7-70 ultracentrifuge (Beckman Instruments, Inc., Palo Alto, CA). Lipoproteins (very low-density lipoproteins, low-density lipoproteins, high-density lipoproteins 2, and high-density lipoproteins 3) were separated by sequential spinning at densities 1.006, 1.063, 1.125, and 1.21 g/ml for 18, 24, 65, and 65 h, respectively. Triglyceride and cholesterol concentrations in serum and in lipoprotein fractions were measured with enzymatic methods (kit nos. 297771 and 187313, Boehringer Diagnostica GmbH, Mannheim, West Germany). Phospholipid concentrations in serum and lipoprotein fractions were measured as inorganic phosphate (25) and protein concentrations with the Lowry method (26). Serum apolipoprotein A-I, A-II and B concentrations were measured with a radial immunodiffusion method (27, 28).

The lipoprotein and hepatic lipase activities of postheparin plasma were measured with the immunochemical method of Huttunen *et al.* (29); lipoprotein lipase was measured after inactivating hepatic lipase with a specific antiserum and hepatic lipase was measured at 1 M NaCl concentration which inactivates lipoprotein lipase, no serum was added.

**Statistical methods.** Statistical analyses were performed by means of BMDP statistical software. We used one-way analysis of variance and unpaired *t* test for comparing groups I–III and the paired *t* test for comparing cord-blood and 2-h values. In addition, we computed correlation coefficients between lipases and the other parameters measured and performed covariance analyses when the other parameters were found to correlate mutually.

## RESULTS

**Hormonal and metabolic parameters.** The parameters were measured to evaluate hyperinsulinemia at birth and to assess the extent of hypoglycemia and lipolysis at the age of 2 h in our nonfed infants. Table 1 compares the values measured in cord blood and at the age of 2 h.

The results indicate that the infants born to insulin-dependent diabetic mothers were hyperinsulinemic at birth and that they became hypoglycemic and had low FFA levels suggesting poor lipolysis at the age of 2 h.

**Cord serum lipoproteins and apoproteins.** The mean serum concentrations of triglycerides, cholesterol, phospholipids, and proteins in lipoprotein classes of cord serum (Table 2) were similar in groups I–III. Also, the concentrations of apolipoproteins A-I, A-II, and B (Table 2) were similar in all groups.

**Postheparin plasma lipoprotein lipase at the age of 2 h.** There was no measurable lipase activity in the blood samples taken before the heparin administration. Fifteen minutes after the heparin bolus of 100 IU/kg, the mean activity of lipoprotein lipase, expressed in  $\mu\text{mol FFA/ml/h}$ , was 38.9 in group I, 35.7 in group II, and 33.0 in group III (Fig. 1A); the values were not statistically different ( $p > 0.5$ ). Taking all three groups together, postheparin-plasma lipoprotein lipase activities at the age of 2 h

Table 2. Lipoprotein-lipids and apoproteins A-I, A-II, and B in cord serum (mean  $\pm$  SEM)\*

	Group		
	I	II	III
Triglycerides (mg/dl)	35.8 $\pm$ 5.0	35.9 $\pm$ 2.9	34.1 $\pm$ 2.7
VLDL†	8.7 $\pm$ 2.2	8.2 $\pm$ 1.4	7.4 $\pm$ 1.3
LDL	14.5 $\pm$ 1.6	12.2 $\pm$ 1.2	12.9 $\pm$ 1.2
HDL	13.8 $\pm$ 2.4	14.5 $\pm$ 1.5	13.9 $\pm$ 1.2
HDL2	10.3 $\pm$ 2.9	6.6 $\pm$ 0.9	8.2 $\pm$ 0.9
HDL3	6.3 $\pm$ 1.2	7.9 $\pm$ 1.2	5.7 $\pm$ 0.7
Cholesterol (mg/dl)	69.1 $\pm$ 3.8	69.2 $\pm$ 3.8	66.7 $\pm$ 3.6
VLDL	2.5 $\pm$ 0.5	2.5 $\pm$ 0.5	2.3 $\pm$ 0.6
LDL	31.5 $\pm$ 1.8	29.3 $\pm$ 2.2	30.3 $\pm$ 2.1
HDL	38.3 $\pm$ 2.5	36.0 $\pm$ 2.6	35.1 $\pm$ 2.1
HDL2	22.3 $\pm$ 1.8	19.3 $\pm$ 2.1	20.8 $\pm$ 1.6
HDL3	15.9 $\pm$ 1.0	16.8 $\pm$ 1.6	16.4 $\pm$ 1.1
Phospholipids (mg/dl)	91.7 $\pm$ 5.6	84.0 $\pm$ 3.1	80.2 $\pm$ 4.4
VLDL	2.5 $\pm$ 0.6	2.5 $\pm$ 0.4	2.5 $\pm$ 0.4
LDL	23.5 $\pm$ 1.7	20.1 $\pm$ 0.9	19.4 $\pm$ 1.1
HDL	65.8 $\pm$ 4.3	61.4 $\pm$ 2.6	58.3 $\pm$ 3.6
HDL2	32.0 $\pm$ 2.8	26.4 $\pm$ 2.5	25.2 $\pm$ 1.7
HDL3	23.0 $\pm$ 1.4	24.3 $\pm$ 1.7	21.3 $\pm$ 1.5
Proteins (mg/dl)			
VLDL	6.8 $\pm$ 1.5	12.3 $\pm$ 3.2	11.7 $\pm$ 2.1
LDL	38.7 $\pm$ 5.9	32.9 $\pm$ 3.3	38.4 $\pm$ 5.4
HDL	103 $\pm$ 8.8	119 $\pm$ 11	120 $\pm$ 11
HDL2	44.8 $\pm$ 4.8	42.8 $\pm$ 4.5	53.8 $\pm$ 9.4
HDL3	58.7 $\pm$ 5.1	76.5 $\pm$ 11	71.9 $\pm$ 7.4
Apo A-I (mg/dl)	96.4 $\pm$ 4.7	103 $\pm$ 3.5	97.2 $\pm$ 2.3
Apo A-II (mg/dl)	28.0 $\pm$ 1.6	30.9 $\pm$ 1.1	27.8 $\pm$ 1.0
Apo B (mg/dl)	29.2 $\pm$ 1.3	27.7 $\pm$ 1.2	27.4 $\pm$ 0.9

\* The number of infants was 19 in group I, 17 in group II, and 15 in group III.

† VLDL, very low-density lipoproteins; LDL, low-density lipoproteins; HDL, high-density lipoprotein.

in these nonfed infants ranged from 12.7 to 81.9  $\mu\text{mol FFA/ml/h}$  (Fig. 1A). In a group consisting of 34 healthy medical students the mean postheparin-plasma lipoprotein lipase activity was 17.4  $\mu\text{mol FFA/ml/h}$  (range 6.6–35.2) (8). The lipoprotein lipase activities of our infants correlated negatively with total triglycerides ( $r = -0.38$ ,  $p < 0.01$ ), very low-density lipoprotein-triglycerides ( $r = -0.33$ ,  $p < 0.05$ ), and low-density lipoprotein-triglycerides ( $r = -0.45$ ,  $p < 0.01$ ) of cord blood.

The infants of group I were hyperinsulinemic. Nevertheless, their lipoprotein lipase activities were comparable to values observed in infants in groups II and III. Taking all groups together, lipoprotein lipase activity was independent of cord-serum C-peptide and insulin. In addition, lipoprotein lipase activity was independent of birth weight and gestational age.

**Postheparin plasma hepatic lipase at the age of 2 h.** There was no measurable lipase activity in the blood samples taken before the heparin administration. Fifteen minutes after the heparin bolus, the mean hepatic lipase activity, expressed in  $\mu\text{mol FFA/ml/h}$ , was 68.3 in group I, 58.7 in group II, and 52.3 in group III (Fig. 1B). Hepatic lipase activity was in group I somewhat higher ( $p < 0.05$ ) than in groups II and III.

Taking all three groups together, hepatic lipase activity correlated negatively ( $r = -0.43$ ,  $p < 0.01$ ) with gestational age, but positively with cord serum C-peptide ( $r = 0.33$ ,  $p < 0.05$ ) and




Fig. 1. Postheparin plasma lipoprotein (A) and hepatic lipase (B) activities of newborn infants. In group I the infants were born to insulin-dependent diabetic mothers and in group II to gestationally diabetic mothers. Group III consisted of large-for-date reference infants. Horizontal lines indicate the mean lipase activities. In  $B$   $p < 0.05$  when comparing groups I–III.

the Hb A1c of mother ( $r = 0.54$ ,  $p < 0.01$ ). However, gestational age, C-peptide, and Hb A1c correlated mutually. Covariance analysis showed that Hb A1c was the only important ( $p < 0.05$ ) factor explaining the variation of hepatic lipase.

#### DISCUSSION

In animals, lipoprotein lipase activity has been found to increase postnatally (1, 6). We found, however, that at the age of 2 h the mean lipoprotein lipase activity in our infants was higher than in term and preterm infants at the age of 1 to 4 days (mean + SD,  $19.7 \pm 10.5$   $\mu\text{mol FFA/ml/h}$ ) or in adults (mean + SD,  $17.4 \pm 6.3$   $\mu\text{mol FFA/ml/h}$ ) (8, 9). Most of our infants were, however, large-for-date (relative birth weight  $> 2.0$  SD units) and thus had more fat tissue, which might account for the higher lipoprotein lipase activity. Lipoprotein lipase activity was, however, independent of birth weight.

As in animals (6), exogenous fat has been suggested to induce lipoprotein lipase activity in man (10, 30). However, in our infants lipoprotein lipase was high although they had not been fed.

Lipoprotein lipase activity at the age of 2 h correlated negatively with cord-serum triglyceride concentration. This implies that lipoprotein lipase can hydrolyze fat at birth. Thus, active lipoprotein lipase may help to prevent hypoglycemia by producing glycerol for gluconeogenesis and FFA for oxidation in tissues. In agreement, administration of fat has been shown to increase blood glucose levels in hypoglycemic small-for-date newborn infants (31) and in starved newborn rats (32). Furthermore, lipoprotein lipase may be significant even for fetal fat storage by shuttling triglycerides for peripheral uptake, as has been proposed to occur in rats with active lipoprotein lipase in adipose tissue at birth (1, 33).

In the infants born to insulin-dependent diabetic mothers (group I) the mean cord-serum insulin level was five times and C-peptide level three times the levels found in the other two infant groups studied. At the age of 2 h, when lipoprotein lipase activity was measured, there was still a profound postnatal effect of insulin in group I: glucose and FFA levels were low although the difference in insulin concentrations between groups I and III had already disappeared. There were, however, no differences in

lipoprotein lipase activities between groups I–III, although insulin is known to induce lipoprotein lipase in adults (3, 12). The infants born to insulin-dependent diabetic mothers had a lower gestational age (37.6 wk) than large-for-date infants (41.2 wk), which might mask the inducing effect of insulin. On the other hand, lipoprotein lipase activity was high in almost all of our infants whereas insulin has been shown to be capable of increasing lipoprotein lipase activity in adult adipose tissue only if the basal lipoprotein lipase activity is low (34, 35). Therefore, the similarity of lipoprotein lipase activities in groups I–III can be explained by assuming saturation of insulin induction.

In the hyperinsulinemic infants of group I the mean hepatic lipase activity was somewhat higher than in groups II and III. The mean activity of postheparin-plasma hepatic lipase groups II and III was as in term and preterm infants (mean + SD,  $57.5 \pm 27.7$   $\mu\text{mol FFA/ml/h}$ ) but almost three times higher than in adults (mean + SD,  $23.1 \pm 10.6$   $\mu\text{mol FFA/ml/h}$ ) (8, 9). According to covariance analysis the only important factor explaining the variation of hepatic lipase in groups I–III was Hb A1c of the mothers. A high HbA1c level is an indicator of poor balance of diabetes, which often produces macrosomia of infants during the last trimester of pregnancy (11, 18). This provides an explanation for the positive correlation between hepatic lipase activity and the level of HbA1c, because hepatic lipase activity is released from the liver, which is large in macrosomic infants.

Cord-serum lipoproteins in our infants were as in normal term infants (36–40). In addition, cord-serum lipoproteins and apolipoproteins were similar in all three groups, although infants born to diabetic mothers have been reported to have higher cholesterol and lower triglyceride levels than normal term infants (41, 42). One possible explanation for this discrepancy is a difference in gestational age: most of our infants were born at term whereas in previous reports infants have been preterm (41, 42).

In conclusion, lipoprotein lipase was active in large-for-date newborn infants immediately after birth and without exogenous fat induction. High lipase activity and its negative correlation with cord-serum triglycerides implies that these newborn infants can efficiently hydrolyze fat at birth. Although insulin is known to induce lipoprotein lipase in adults, lipoprotein lipase activity was independent of serum insulin level in our hyperinsulinemic infants.

*Acknowledgments.* The authors thank Mrs. Sirkka Runeberg, Miss Paula Teräväinen, Mrs. Leena Lehtikainen, and Mrs. Hannele Hilden for their skillful technical assistance.

## REFERENCES

- Hahn P 1978 Lipids. In: Stave U (ed) *Perinatal Physiology*, 2nd ed. Medical Book Co, New York, pp 397-423
- Levene MI 1981 Tolerance of parenteral fat in the preterm neonate. In: Wosdorp, Ric, Soeters (eds) *Clinical Nutrition 81'*, Churchill Livingstone, New York, pp 97-104
- Cryer A 1981 Tissue lipoprotein lipase activity and its action in lipoprotein metabolism. *Int J Biochem* 13:525-541
- Jackson RL 1983 Lipoprotein lipase and hepatic lipase. In: Boyer PD (ed) *The Enzymes*. Academic Press, New York, pp 141-181
- Jansen H, Birkenhager JC 1981 Liver lipase-like activity in human and hamster adrenocortical tissue. *Metabolism* 30:428-430
- Das JB, Joshi IK, Philippart AI 1982 Postnatal development of lipid-clearing enzymes in the suckling animal. *J Pediatr Surg* 17:914-919
- Rosseneu M, van Biervliet JP, Bury J, Vinaimont N 1983 Isolation and characterization of lipoprotein profiles in newborns by density gradient ultracentrifugation. *Pediatr Res* 17:788-794
- Rovamo L, Taskinen MR, Kuusi T, Nikkilä EA, Ehnholm C, Raivio KO 1984 Postheparin plasma lipase activities and plasma lipoproteins in newborn infants. *Pediatr Res* 18:642-647
- Rovamo L, Nikkilä EA, Taskinen MR, Raivio KO 1984 Postheparin plasma lipoprotein and hepatic lipases in preterm neonates. *Pediatr Res* 18:1104-1107
- Rovamo L 1985 Postheparin plasma lipases and carnitine in infants during parenteral nutrition. *Pediatr Res* 19:292-297
- Cowett RM, Schwartz R 1982 The infant of the diabetic mother. *Pediatr Clin North Am* 29:1213-1231
- Sadur CN, Eckel RH 1982 Insulin stimulation of adipose tissue lipoprotein lipase. Use of the euglycemic clamp technique. *J Clin Invest* 69:1119-1125
- Nilsson-Ehle P 1981 Impaired regulation of adipose tissue lipoprotein lipase in obesity. *Int J Obesity* 5:695-699
- Pykälistö OJ, Smith PH, Brunzell JD 1975 Determinants of human adipose tissue lipoprotein lipase. Effect of diabetes and obesity on basal- and diet-induced activity. *J Clin Invest* 56:1108-1117
- White P 1965 Pregnancy and diabetes, medical aspects. *Med Clin North Am* 49:1015-1024
- Teramo K, Kuusisto AN, Raivio KO 1979 Perinatal outcome of insulin-dependent diabetic pregnancies. *Ann Clin Res* 11:146-155
- Teramo K, Ämmälä P, Ylinen K, Raivio KO 1983 Pathologic fetal heart rate associated with poor metabolic control in diabetic pregnancies. *Obstet Gynecol* 61:559-565
- Ylinen K, Raivio K, Teramo K 1981 Haemoglobin A<sub>1c</sub> predicts the perinatal outcome in insulin-dependent diabetic pregnancies. *Br J Obstet Gynaecol* 88:961-967
- Ylinen K, Hekali R, Teramo K 1981 Haemoglobin A<sub>1c</sub> during pregnancy of insulin-dependent diabetics and health controls. *J Obstet Gynaecol* 1:223-228
- Hallman M, Teramo K 1979 Amniotic fluid phospholipid profile as a predictor of fetal maturity in diabetic pregnancies. *Obstet Gynecol* 54:703-709
- Usher R, McLean F 1969 Intrauterine growth in live-born caucasian infants at sea level: standards obtained from measurements in 7 dimensions of infants born between 25 and 44 weeks of gestation. *J Pediatr* 74:901-910
- Desbuquois B, Aurbach GD 1971 Use of polyethylene glycol to separate free and antibody-bound peptide hormones in radioimmunoassays. *J Clin Endocrinol* 33:732-738
- Kuzuya H, Blix PM, Horwitz DL, Steiner DF, Rubenstein AH 1977 Determination of free and total insulin and C-peptide in insulin-treated diabetics. *Diabetes* 26:22-29
- Olsen C 1971 An enzymatic fluorimetric micromethod for the determination of acetoacetate, 3-hydroxybutyrate, pyruvate and lactate. *Clin Chem Acta* 33:293-296
- Bartlett GR 1959 Phosphorus assays in column chromatography. *J Biol Chem* 234:466-469
- Lowry OH, Rosebrough NJ, Farr AI, Randall RJ 1951 Protein measurement with the folin phenol reagent. *J Biol Chem* 193:265-275
- Cheung MC, Albers JJ 1977 The measurement of apolipoprotein A-I and A-II levels in men and women by immunoassay. *J Clin Invest* 60:43-50
- Curry MD, Gustafson A, Alaupovic P, McConathy WJ 1978 Electroimmunoassay, and radial immunodiffusion assay evaluated for quantification of human apolipoprotein B. *Clin Chem* 24:280-286
- Huttunen JK, Ehnholm C, Kinnunen PKJ, Nikkilä EA 1975 An immunochemical method for the selective measurement of two triglyceride lipases in human postheparin plasma. *Clin Chim Acta* 63:335-347
- Taskinen MR, Tulikoura I, Nikkilä EA, Ehnholm C 1981 Effect of parenteral hyperalimentation on serum lipoproteins and on lipoprotein lipase activity of adipose tissue and skeletal muscle. *Eur J Clin Invest* 11:317-323
- Mestyan J, Rubecz I, Soltesz G 1976 Changes in blood-glucose, free fatty acids and amino-acids in low-birth-weight infants receiving intravenous fat emulsion. *Biol Neonate* 30:74-79
- Ferre PR, Pegorier JP, Girard JR, Marliss EB 1977 Evidence that fatty acid oxidation stimulates gluconeogenesis in newborn rats. *Biochem Soc Trans* 5:982-990
- Cryer A 1985 Lipoprotein lipase and the uptake of lipid by adipose cells during development. *Reprod Nutr Dev* 25:255-270
- Taskinen MR, Nikkilä EA 1979 Lipoprotein lipase activity of adipose tissue and skeletal muscle in insulin-deficient human diabetes. *Diabetologia* 17:351-356
- Yki-Järvinen H, Taskinen MR, Koivisto VA, Nikkilä EA 1984 Response of adipose tissue lipoprotein lipase activity and serum lipoproteins to acute hyperinsulinaemia in man. *Diabetologia* 27:364-369
- Ginsburg BE, Zetterström R 1980 Serum cholesterol concentrations in early infancy. *Acta Paediatr Scand* 69:581-585
- Hardell LI Serum lipids and lipoproteins at birth based on a study of 2815 newborn infants. I. Concentrations and distributions of triglycerides and cholesterol. *Acta Paediatr Scand* 285:5-10
- McConathy WJ, Lane DM 1980 Studies on the apolipoproteins and lipoproteins of cord serum. *Pediatr Res* 14:757-761
- Strobl W, Widhalm K, Kostner G, Pollak A 1983 Serum apolipoproteins and lipoproteins during the first week of life. *Acta Paediatr Scand* 72:505-509
- van Biervliet JP, Vinaimont N, Caster H, Vercaemst R, Rosseneu M 1981 Plasma apoprotein and lipid patterns in newborns: influence of nutritional factors. *Acta Paediatr Scand* 70:851-856
- Andersen GE, Lous P, Friis-Hansen B 1979 Hyperlipoproteinemia in newborn infants. A study of 1025 families. *Acta Paediatr Scand* 68:683-690
- Hardell LI Serum lipids and lipoproteins at birth based on a study of 2815 newborn infants. II. Relations between materno-foetal factors and the concentrations of triglycerides and cholesterol. *Acta Paediatr Scand [Suppl]* 285:11-20