Synergistic Action of Triiodothyronine and Hydrocortisone on Epinephrine-Induced Reabsorption of Fetal Lung Liquid

P. M. BARKER, M. MARKIEWICZ, K. A. PARKER, D. V. WALTERS, AND L. B. STRANG

Department of Paediatrics, University College and Middlesex School of Medicine, Rayne Institute, London, WC1E 6JJ, United Kingdom

ABSTRACT. The influence of triiodothyronine and hydrocortisone on maturation of the response to epinephrine that leads to reabsorption of lung liquid was investigated in nine chronically catheterized fetal sheep. Experiments were performed on thyroidectomized fetal sheep at 116–120 d gestation, well before the reabsorptive response to epinephrine is normally seen. After i.v. administration of either triiodothyronine (60 µg/d) or hydrocortisone (10 mg/d) for 3 d (three fetuses in each case), all fetuses continued to secrete lung liquid during exposure to epinephrine (secretion rate = 5.9 ± 3.2 mL/h in triiodothyronine-treated and 4.4 ± 1.9 mL/h in hydrocortisone-treated fetuses). However, when the two hormones were administered together in the same doses to three fetuses, a striking reabsorptive response to epinephrine was seen (absorption rate = -12.3 ± 3.6 mL/h), similar to that observed in the mature fetus. Induction of this capacity to reabsorb lung liquid may be of importance in the management of respiratory problems of the newborn infant. (*Pediatr Res* 27: 588–591, 1990)

Abbreviations

T₃, triiodothyronine
T₄, thyroxine
TRH, thyrotropin-releasing hormone

The secretion of lung liquid by the pulmonary epithelium maintains expansion and promotes development of the fetal lung (1, 2). At birth, however, effective aeration of this organ is dependent on rapid reabsorption of the liquid from its alveolar spaces (3, 4). The switch from secretion to absorption depends on a reversal in direction of net ion transport across the pulmonary epithelium, brought about by a change from active chloride transfer into the alveolar lumen (2) to the active transport of sodium from the lumen towards the interstitium (5). This change is normally activated by a rise in fetal plasma epinephrine brought about by the stress of labor and delivery (4). In the fetal sheep (gestation = 147 d), a reabsorptive response to infusion of i.v. epinephrine in the fetus is seen from about 130 d (3, 4), and this response increases more than 20-fold in the last 2 wk of gestation. We have undertaken an investigation of hormones that might control the maturation of this reabsorptive response. We have shown previously that thyroidectomy prevents development of the reabsorptive response (6) and that this can be restored by the infusion of T₃ to the thyroidectomized fetus (7).

However, precocious exposure of the fetus to T₃ at an early gestation did not significantly advance the time of appearance of this reabsorptive capacity, from which it could be concluded that some other factor must play a role in determining its maturation during gestation. Steroid hormones, alone or in combination with thyroid hormones, are known to be important in determining the maturation of structural aspects of the lung and its surfactant system in the fetus (8, 9). Our experiments were undertaken to examine the influence of hydrocortisone alone and in combination with T₃ on the maturation of the reabsorptive response. Fetal sheep were studied at 116–120 d, which is a gestation well before that at which the reabsorptive response to epinephrine normally appears.

MATERIALS AND METHODS

Experimental preparation. The surgical preparation of chronically-catheterized fetal sheep to allow for repeated measurement of lung liquid secretion or absorption rate was identical to that described in detail by Walters and Olver (3). Nine dated, pregnant ewes were operated on at 112–115 d gestation under general anesthesia induced by thiopentone and maintained with fluothane. Fetal thyroidectomy was carried out and wide bore (inner diameter, 2.6 mm) catheters placed into each end of the incised fetal trachea, finer gauge catheters were inserted into the carotid artery and jugular vein. The catheters were brought out through the uterus and flank of the ewe and the tracheal catheters joined to form an external loop that allowed uninterrupted flow of lung liquid from the fetal lung to the larynx between experiments. After surgery, antibiotics were given for the first 48 h to the ewe (streptomycin 500 mg intramuscularly, benzyl penicillin 300 mg intramuscularly) and to the fetus (streptomycin 70 mg i.v., benzyl penicillin 50 mg i.v.). No experiments were performed during this recovery period.

Experimental procedure. The measurement of lung liquid secretion or absorption depends on the use of an impermeant tracer (¹²⁵I-albumin) added to lung liquid. Dilution of the tracer over a time period of 30 min by newly formed liquid allows accurate determination of secretion rate, whereas an increase in concentration reflects a diminution in volume, or reabsorption. Under sterile conditions, the exteriorized loop of tracheal catheter was interrupted and the distal (lung) half was connected to a glass burette into which approximately half of the lung liquid could be withdrawn by gravity. The impermeant tracer was added to this reservoir and thoroughly mixed into lung liquid by repeated raising and lowering of the burette. After half an hour of mixing, the resting secretion rate was calculated from the changing concentration of tracer in seven samples of lung liquid taken at 5-min intervals over 30 min. In the second half of the experiment, similar measurements were made during epinephrine infusion (0.5 µg/min) over a 30-min period commencing 15 min after the start of the infusion. In both control and thyroid-

Received October 30, 1989; accepted January 24, 1990.

Correspondence: Dr. D. Walters, Department of Paediatrics, University College and Middlesex Medical School, University Street, London WC1 6JJ, UK.

ectomized fetuses, infusion of epinephrine at this rate has been shown to increase resting fetal plasma epinephrine concentration from about 0.1 ng/mL to about 1.0 ng/mL (4, 6). Samples of carotid arterial blood were taken from the fetal carotid artery catheter for blood gas determinations and fetal hormone analysis before and during epinephrine infusion. Blood samples were immediately separated and the plasma stored at -40°C for later analysis. Arterial blood pressure and heart rate were measured by means of a transducer set at the estimated level of the fetal heart, which allowed for measurement of heart rate and relative changes in mean blood pressure during the experiment but precluded measurement of absolute blood pressure. At the end of the experiment, the epinephrine infusion was discontinued, sulphmethazine was mixed into the lung liquid and given i.v. (165 mg into each compartment) to reduce the risk of infection, and the tracheal loop reestablished to allow normal flow of lung liquid to the fetal larynx.

Infusion of T_3 , hydrocortisone, or T_3 and hydrocortisone. Before starting the hormone infusion, a control experiment was performed as described above to measure the secretion rates of lung liquid before and during epinephrine infusion. The infusion of T_3 alone, hydrocortisone alone, or T_3 and hydrocortisone together was then started and continued for 3 d (Grasby Syringe Driver, Grasby Medical Ltd., Watford, Herts, UK, Msl6). The daily dose of T_3 (60 $\mu\text{g}/\text{d}$, Glaxo Labs, Ltd., London, England), hydrocortisone succinate (10 mg/d, Upjohn, Crawley, England), or the two hormones in combination (same doses) were diluted in 10 mL sterile water and infused continuously over 24 h. (See Discussion for explanation of high T_3 dosage.) There were three fetuses in each treatment group. After 3 d of hormone administration, a second experiment was performed to determine any hormonal influence on resting and epinephrine-stimulated lung liquid flow.

Hormone concentrations. Fetal plasma T_3 , T_4 , and cortisol concentrations were measured using commercially available radioimmunoassay kits (Amersham International, Little Chalfont, Buckinghamshire, UK).

All values given are mean $\pm$ SD. Where paired observations were made, a paired t test is used.

RESULTS

T_3 and cortisol concentrations. At the time of the first experiment (2–4 d after surgery), fetal plasma T_4 and T_3 concentrations had fallen to very low levels (mean T_4 , 14.8 ± 5.1 ng/mL; mean T_3 , 0.12 ± 0.06 ng/mL) as a result of fetal thyroidectomy [normal values for fetuses of 120–130 d gestation: T_4 98.8 ng/mL, T_3 0.21 ng/mL (6)]. Following 3 d T_3 infusion, mean T_3 concentration rose from 0.14 ± 0.08 ng/mL to 0.96 ± 0.63 ng/mL, and, after 3 d combined T_3 and hydrocortisone infusion, from 0.06 ± 0.05 ng/mL to 0.5 ± 0.26 ng/mL. Three d after hydrocortisone infusion, plasma cortisol had risen from 1.05 ± 0.35 $\mu\text{g}/\text{dL}$ to 14.03 ± 12.82 $\mu\text{g}/\text{dL}$, and, after combined hormone infusion, from 1.03 ± 0.87 $\mu\text{g}/\text{dL}$ to 5.9 ± 1.56 $\mu\text{g}/\text{dL}$. These concentrations of both T_3 and hydrocortisone were similar to those normally seen in the fetal sheep a few days before delivery (10, 11).

Fetal blood pressure, heart rate, and arterial blood gas monitoring. All three experimental groups of fetuses showed similar rises in mean blood pressure and heart rate during epinephrine infusion both before and after exposure to hormones. Thus, the increases during epinephrine infusion before hormone administration were: blood pressure, 4.2 ± 2.9 mm Hg and heart rate, 13.3 ± 10.9 beats/min, whereas, after 3 d of hormone exposure, these increases were 3.6 ± 3.3 mm Hg and 9.5 ± 10.1 beats/min. Epinephrine infusion caused a small fall in pH (7.39 ± 0.03 to 7.37 ± 0.03 , $p < 0.05$) and a small rise in PCO_2 (45.7 ± 4.8 to 48.4 ± 5.1 , $p < 0.05$). There were no differences in these measurements between the three groups and no changes in PO_2 were detected.

The effect of T_3 , hydrocortisone, and combined T_3 and hydro-

cortisone on response of lung liquid secretion/absorption to epinephrine. In all three groups of fetuses, after 3 d of hormone exposure, there was a similar rise in resting (preepinephrine infusion) secretion rate (6.2 ± 1.4 mL/h to 9.1 ± 3.7 mL/h, $n = 9$, $p < 0.05$). In all three groups, the control epinephrine infusion before T_3 , hydrocortisone, or combined hormone exposure produced the small, expected decrease in secretion rate normally seen at this gestation (Table 1). In both the T_3 and hydrocortisone-infused groups, the secretion rates during epinephrine infusion in the 2nd experiments after 3 d hormone exposure were no different from those in the 1st prehormone experiment. However, in the three fetuses treated with a combination of T_3 and hydrocortisone, there was a most striking reabsorptive response to epinephrine at 120 d gestation, comparable to that normally seen in the euthyroid fetus at 135 to 140 d (Figure 1).

DISCUSSION

These results show that the combined action of T_3 and hydrocortisone can induce changes in the ion transport mechanism of the pulmonary epithelium in the fetal sheep, enabling it to respond to β -adrenergic stimulation at a much earlier age than is normally observed. Both these hormones are clearly necessary

Table 1. Secretion (+) or absorption (–) rates (mL/h) of fetal lung liquid before (J_{V_C}) and during (J_{V_E}) epinephrine infusion (0.5 $\mu\text{g}/\text{min}$) in nine thyroidectomized fetal sheep in single experiments performed before and 3 d after being administered T_3 , hydrocortisone (Hc), or T_3 and hydrocortisone

	Before hormone		After hormone	
	J_{V_C}	J_{V_E}	J_{V_C}	J_{V_E}
T_3 -infused group	+5.8	+2.8	+7.1	+2.4
	+8.6	+5.7	+14.4	+8.6
	+7.1	+3.5	+6.2	+6.8
Hc-infused group	+5.8	+7.4	+14.3	+6.6
	+7.2	+5.9	+9.5	+3.8
	+5.1	+1.2	+5.5	+2.9
T_3 and Hc-infused group	+7.1	+2.5	+11.1	–15.8
	+2.8	+1.8	+4.3	–12.4
	+4.0	+4.0	+4.3	–8.6

Fig. 1. Values of lung liquid secretion rate (J_v) or reabsorption rate ($-J_v$) during i.v. epinephrine infusion (0.5 $\mu\text{g}/\text{min}$) plotted against gestational age (d). The resting secretion rates before epinephrine infusion are not shown. ●, values from three thyroidectomized fetal sheep before (116 d) and 3 d after (119 d) continuous infusion of T_3 (60 $\mu\text{g}/\text{d}$) and hydrocortisone (10 mg/d). ○, mean ($\pm$ SEM) J_v for 25 nonthyroidectomized fetuses grouped in 5-d gestations from 120 to 145 d [from Brown *et al.* (4)].

for this development as neither hormone alone produced any change in the response to epinephrine.

Hormone concentrations and timing of reabsorptive response. The gestation at which the lung liquid reabsorptive response to epinephrine first appears in the fetal sheep coincides with increases in the concentrations of both thyroid and steroid hormones in fetal plasma. Maturation of the β -ring deiodination pathway required for conversion of T_4 to its active metabolite T_3 does not take place to any significant extent in the fetal sheep until after 130 d gestation (10) so that fetal plasma concentrations of T_3 remain low (0.2 ng/mL or less) in the sheep until the last 3 wk of gestation during which time a gradual rise in T_3 concentrations to about 0.5 ng/mL is seen (11). Very high doses of T_3 were needed to produce the required plasma concentrations because T_3 is cleared much more rapidly in the fetus (12) than in the postnatal animal; this is probably the result of very active conversion of T_3 to T_2 (diiodothyronine) by α -monoiodination in the placenta (13). Fetal plasma cortisol concentrations are similarly low throughout most of gestation, principally as a result of conversion by the placenta of fetal and maternal cortisol to its inactive metabolite cortisone (14). A gradual rise in fetal cortisol concentration is seen in the fetal sheep from about 130 d (15). Nathanielsz *et al.* (16) reported a later, more pronounced rise in cortisol concentrations to 10 μ g/dL only in the last few days before delivery. Thus, hormone concentrations encountered by fetuses at gestations less than 120 d in our experiments were similar to those normally seen only close to full term.

Probable site of action of T_3 and hydrocortisone. The reabsorption of lung liquid induced by epinephrine is mediated by the net flux of sodium ions across the pulmonary epithelium from the alveolar lumen (5). Since this sodium-led reabsorption can be blocked by micromolar concentrations of amiloride in lung liquid (5), it has been inferred that the "opening" or activation of sodium channels in the apical membranes of epithelial cells constitutes the crucial change underlying epinephrine-induced absorption. A similar pattern of maturation is seen when dibutyryl cAMP is used to initiate the reabsorptive response (17), and this development can be completely inhibited in the sheep by fetal thyroidectomy at 118 d (6). It follows, therefore, that the maturation of this reabsorptive response depends not on adrenoceptor development but on changes in some components beyond cAMP in the intracellular signalling system. The synergistic action of T_3 and hydrocortisone can probably be attributed to stimulation of synthesis of proteins that control, or are part of, the structure of the apical membrane sodium channels. The increase in the resting secretion rate during T_3 and hydrocortisone infusion is likely to reflect the normal rise in secretion rate seen during this part of gestation (4) rather than a T_3 /hydrocortisone-related effect, particularly as the secretion rate increased similarly in all three groups.

Clinical implications. Of all the pulmonary adaptations taking place at birth, the most obvious, the least studied, and possibly the most important is the change from the liquid-filled to the air-filled state. Similarly, it seems probable that the importance of inadequate reabsorption or continuing secretion of pulmonary liquid in preterm infants has not been adequately recognized as a cause of neonatal respiratory difficulties. It has been known for some time that T_3 and corticosteroids appear to act in concert to advance structural maturation (9) and to promote surfactant synthesis in both organ cultures of the fetal rat lung and the intact fetal rat (18). More recently, Warburton *et al.* (19) found that administration of corticosteroids and TRH together to the fetal sheep led to maturation of the surfactant system and improved lung mechanics. Both of these effects were enhanced when a β -agonist was infused in addition to the hormones. Similarly, Schellenberg *et al.* (20) have shown that a combination of T_3 , hydrocortisone, and prolactin infused into the fetal sheep increased lung distensibility and stability. Interestingly, Liggins *et al.* (21) found that although a significant increase in alveolar lavage-saturated phosphatidylcholine occurred in response to

TRH infused to sheep fetuses from 124 to 128 d gestation, improvement in lung distensibility and stability was not seen unless TRH was infused together with cortisol. In light of our findings, these results may have been partly or mainly due to the effect of these hormones on the pulmonary liquid reabsorptive system.

In the context of our experiments, very premature infants are most likely to be born at a stage when the lungs have had insufficient prenatal exposure to T_3 and corticosteroid hormones. Mean cord concentrations of both these hormones are known to be significantly lower in infants born preterm (22, 23), and it has been shown that, within the preterm group, low cord concentrations of both thyroid hormones (24) and cortisol (23) at birth are associated with a higher incidence of subsequent respiratory problems. Lucas *et al.* (25) report that 50% of T_3 concentrations in the first 36 h of life were below 0.19 ng/mL (0.3 nM), whereas the results obtained by Barker *et al.* (26) suggested that concentrations over 0.3 ng/mL may be required to induce the reabsorptive mechanism in the fetal sheep. Although it appears that many preterm infants are able to mount a substantial postnatal surge in T_3 (22), this would be too late to prevent the onset of respiratory difficulties that may be associated with poor lung liquid clearance in the first 24 h of life.

Our experiments show that the reabsorptive response to epinephrine can be induced much earlier in gestation than is normally seen, and within a short time scale (Fig. 1), by exogenous administration of T_3 and hydrocortisone. The relative independence of this aspect of lung development from any fixed developmental program and its ability to respond to hormones at a very early stage of gestation may be important factors in determining the severity of respiratory disease in those very small preterm infants who have had insufficient prenatal exposure to T_3 and hydrocortisone. Since cortisol and T_3 concentrations can be manipulated both pre- (27, 28) and postnatally, it may be possible to use these hormones to enhance the lung liquid reabsorptive capacity of the immature lung.

REFERENCES

1. Alcorn D, Adamson TM, Lambert TF, Maloney JE, Ritchie BC, Robinson PM 1977 Morphological effects of chronic tracheal ligation and drainage in the fetal lamb lung. *J Anat* 123:649-660
2. Olver RE, Strang LB 1974 Ion fluxes across the pulmonary epithelium and the secretion of lung liquid in the foetal lamb. *J Physiol* 241:327-357
3. Walters DV, Olver RE 1978 The role of catecholamines in lung liquid absorption at birth. *Pediatr Res* 12:239-242
4. Brown MJ, Olver RE, Ramsden CA, Strang LB, Walters DV 1983 Effects of epinephrine and of spontaneous labour on the secretion and absorption of lung liquid in the fetal lamb. *J Physiol* 344:137-152
5. Olver RE, Ramsden CA, Strang LB, Walters DV 1986 The role of amiloride-blockable sodium transport in epinephrine-induced lung liquid reabsorption in the fetal lamb. *J Physiol* 376:321-340
6. Barker PM, Brown MJ, Ramsden CA, Strang LB, Walters DV 1988 The effect of thyroidectomy in fetal sheep on lung liquid reabsorption induced by epinephrine or cyclic AMP. *J Physiol* 407:373-383
7. Barker PM, Strang LB, Walters DV 1989 The role of triiodothyronine (T_3) in the maturation of epinephrine-induced reabsorption of lung liquid in the fetal sheep. *J Physiol* 416:51(abstr)
8. Liggins GC 1976 Adrenocortical-related maturational events in the fetus. *Am J Obstet Gynecol* 126:931-939
9. Gross I, Dynia DW, Wilson CM, Ingleson LD, Gewolb IH, Rooney SA 1984 Glucocorticoid-thyroid hormone interactions in the fetal rat lung. *Pediatr Res* 18:191-195
10. Wu SY, Klein AH, Chopra IJ, Fisher DA 1978 Alterations in tissue thyroxine 5'-monodeiodinating activity in the perinatal period. *Endocrinology* 103:235-239
11. Fisher DA, Dussault JH, Sack J, Chopra IJ 1977 Ontogenesis of hypothalamic-pituitary thyroid function and metabolism in man, sheep and rat. *Recent Prog Horm Res* 33:59-107
12. Fraser M, Liggins GC 1988 Thyroid hormone kinetics during late pregnancy in the ovine fetus. *J Dev Physiol* 10:461-471
13. Roti E, Braverman LE, Fang S, Alex S, Emerson CH 1982 Ontogenesis of placental inner ring thyroxine deiodinase and amniotic fluid 3,3',5'-triiodothyronine concentration in the rat. *Endocrinology* 111:959-963
14. Buster JE 1983 Gestational changes in steroid hormone biosynthesis, secretion, metabolism and action. *Clin Perinatal* 10:527-552
15. Ballard PL, Kitterman JA, Bland RD, Clyman RJ, Gluckman PD, Platzker CG, Kaplan SL, Grumbach MM 1982 Ontogeny and regulation of cortico-

- steroid binding globulin capacity in the plasma of fetal and newborn lambs. *Endocrinology* 110:359-366
16. Nathanielsz PW, Comline RS, Silver M, Paisley RB 1972 Cortisol metabolism in the fetal and neonatal sheep. *J Reprod Fertil [Suppl]* 16:39-59
17. Olver RE, Ramsden CA, Walters DV 1987 The effect of dibutyryl cyclic AMP on lung liquid volume flow across the pulmonary epithelium of the fetal sheep. *J Physiol* 391:61(abstr)
18. Hitchcock KB 1979 Hormones and the lung. 1. Thyroid hormones and glucocorticoids in lung development. *Anat Rec* 194:15-40
19. Warburton D, Parton L, Buckley S, Cosico L, Enns G, Saluna T 1988 Combined effects of corticosteroid, thyroid hormones, and β -agonist on surfactant, pulmonary mechanics, and β -receptor binding in fetal lamb lung. *Pediatr Res* 24:166-170
20. Schellenberg JC, Liggins GC, Manzai M, Kitterman JA, Lee CH 1988 Synergistic hormonal effects on lung maturation in fetal sheep. *J Appl Physiol* 65:94-100
21. Liggins GC, Schellenberg JC, Manzai M, Kitterman JA, Lee CH 1988 Synergism of cortisol and thyrotropin-releasing hormone in lung maturation in fetal sheep. *J Appl Physiol* 64:1880-1884
22. Cuestas RA 1978 Thyroid function in healthy premature infants. *Pediatrics* 92:963-967
23. Sybulski S, Maughan GB 1976 Relationship between cortisol levels in umbilical cord plasma and the development of the respiratory distress syndrome in premature newborn infants. *Am J Obstet Gynecol* 125:239-243
24. Cuestas RA, Lindall A, Engel RE 1976 Low thyroid hormones and respiratory distress syndrome of the newborn. Studies on cord blood. *N Eng J Med* 295:297-302
25. Lucas A, Rennie J, Baker BA, Morley R 1988 Low plasma triiodothyronine concentrations and outcome in preterm infants. *Arch Dis Child* 63:1201-1206
26. Barker PM, Walters DV, Strang LB 1990 The role of thyroid hormones in maturation of the adrenaline-sensitive lung-liquid reabsorptive mechanism in fetal sheep. *J Physiol* 424:473-485
27. Roti E, Gnudi A, Braverman LE, Robuschi G, Emanuele R, Bandini P, Benassi L, Pagliani A, Emerson CH 1981 Human cord blood concentrations of thyrotropin, thyroglobulin, and iodothyronines after maternal administration of thyrotropin-releasing hormone. *J Clin Endocrin Metab* 53:813-819
28. Ballard PL, Liggins GC 1982 Glucocorticoid activity in cord serum: comparison of hydrocortisone and betamethasone regimes. *Pediatr Res* 101:468-471