

3 α -ANDROSTANEDIOL GLUCURONIDE AS A MARKER OF ANDROGENIC ACTIVITY IN GIRLS WITH PREMATURE PUBARCHÉ. M. Gryngarten, ME. Escobar, S. Campo, S. Ayuso, P. Bedecarrás, C. Bergadá. CEDIE. Div. de Endocrinol. Hosp. de Niños R. Gutierrez. Bs.As., Argentina.

Previous reports have postulated that 3 androstenediol glucuronide (3 α AG) constitutes a marker of the 5 reductase activity in peripheral tissues. We studied the serum levels of androgens and their metabolites in 20 girls with premature pubarche (PP), EC X: 6.6 \pm 2.3 years, and in a control group of 12 girls, 5 of them prepubertal and the other 7 pubertal. 17 hydroxiprogesterone (17OHP), Androstenedione (Δ 4A), dehydroepiandrosterone sulfate (DS), testosterone (T) and 3 AG were determined by RIA. All patients showed pubic hair Tanner grade II-III. Besides pubic hair, 13 girls had one or more associated androgenic signs like acne; or face, trunk or limbs hair. Serum T and 17OHP levels were in prepubertal range in girls with PP except in one that showed basal 17OHP of 2 ng/ml and a response to ACTH stimulation of 20, which lead to a diagnosis of non classical 21OHase deficiency.

	3 α AG	3 AG/ Δ 4A
Prepubertal C	n=5 2.08 \pm 1.0	1.47 \pm 0.79
Pubertal C	n=7 2.96 \pm 1.61	1.12 \pm 0.61
PP	n=20 6.89 \pm 4.65	2.23 \pm 1.57
PP with androgenic signs	n=13 7.40 \pm 4.70	2.36 \pm 1.88
PP without androgenic signs	n=7 5.95 \pm 4.50	1.90 \pm 0.75

The elevated 3 α AG levels found in PP agree with the pubertal levels of its precursors Δ 4A and DS. The high 3 α AG/ Δ 4A ratio found in PP associated with other androgenic signs suggests and enhanced peripheral 5 reductase activity in those patients.

SERUM LH AND FSH DURING THE FIRST TRIMESTER OF LIFE IN NORMAL (C) CHILDREN AND IN AMBIGUOUS GENITALIA. S. Chahin, E. Chaler, M. Maceiras, A. Belgorosky, MA. Rivarola. Servicio de Endocrinología. Hosp. J.P. Garrahan, Bs.As., Argentina.

Little information is available on serum LH and FSH levels during the first 3 months of life measured by a RIA which utilizes a monoclonal antibody specific for LH submit. In order to evaluate the gonadotropic profile and the maturation of the hypothalamo-pituitary-gonadal axis, serum LH and FSH were determined by IRMA in 55 control children, age 5-90 day-old, the mean chronological age (X \pm SD) was 27.9 \pm 15 for males (M) and 26.8 \pm 23.2 days for females (F). In 5 F with congenital adrenal hyperplasia (CAH), mean age 23.8 \pm 21.4 days and in 7 M babies with 46XY ambiguous genitalia (serum testosterone 4.92 \pm 2.56 nmol/L, good clinical response to androgens), mean age 15.3 \pm 10.5 days. Serum levels of LH in normal females were significantly lower (1.07 \pm 1.80 and 3.85 \pm 3.4 U/L) and those of FSH significantly higher (7.68 \pm 7.12 and 2.82 \pm 2.75 U/L) than in normal M, respectively. No correlation was found between serum LH or FSH with age (5-90 days old) in C F or M. Girls with CAH had serum FSH (0.64 \pm 1.05 U/L) significantly lower than F C (p<0.05) but serum LH (0.20 \pm 0.26 U/L) was not different. In boys with 46,XY ambiguous genitalia, serum LH (2.09 \pm 2.17) and serum FSH (2.28 \pm 2.31) were not significantly different from M C. It is concluded that, 1) serum LH and FSH do not change with age during the first trimester, 2) serum LH is higher in M and serum FSH is higher in F, and 3) inhibitory effect of adrenal androgens is present in CAH at this age.

INHIBIN GENERATION IN CRYPTORCHID (CRI) BOYS UNDER 4 YEARS OF LIFE. CA. Longui, IJP. Arnold; BB. Mendonca; W. Bloise; N. Lahlou. Unidade de Gonadas en Intersexo - FMUSP - Sao Paulo-Brasil. Fundation de Recherche en Hormologie-Fresnes- France.

Thirty cryptorchid boys with pre-scrotal testes were divided in two groups: Group I (3 bil/12 unil) received hCG (750U/m2/dose)+hMG (100U/m2/dose) 2 times weekly, during 6 weeks; Group II (5 bil/10 unil) received only hCG. Surgical correction was necessary in 28 cases, 22 of which had a testicular biopsy and subsequent histomorphological analysis. Results were obtained in basal condition and at 3rd. and 6th. weeks of treatment. Peak values are showed:

	LH	FSH	TESTO	INHIBIN	SPERM	At(u2)
UNI 39(10)	7(1.4)	370(206)	450(228)	T=67(44)	T=222(18)	
				C=11(10)	C=194(30)	
G-1						
BIL 39(16)	6(1.5)	197(122)	310(269)	24(20)	198(20)	
UNI 36(14)	4(0.6)	441(290)	257(183)	T=26(18)	T=238(20)	
				C=14(9)	C=193(23)	
G-11						
BIL 44(26)	4(0.5)	515(330)	438(199)	10(7)	207(35)	

T=topic C=cryptorch. Sperm=spermatogonia/50 tubes At=tubular area. LH(mUI/ml) FSH(mUI/ml) Testosterone (ng/dl) inhibin (UI/L). There was no advantage in hMG association. The hormonal level fell between 3rd 6th weeks (down-regulation). There was no correlation between inhibin levels and testosterone, spermatogonia or tubular area. Tubular area and number of spermatogonias were decreased on the affected side in relation to the topic testes and in bilateral cases in relation to the normal patterns.

IGF-I AND IGF-II IMMUNOREACTIVITY IN CONDITIONED MEDIA FROM PRIMARY CULTURES OF RAT SERTOLI CELLS. H. Jasper, S. Cigorraga, Endocrinology, "R.Gutierrez" Children's Hospital, Buenos Aires, Argentina.

IGF-I production by rat Sertoli cells has been reported, but evidences of IGF-II production are lacking. Trying to prove it, we measured IGF-II (as well as IGF-I) immunoreactivity in conditioned media from primary cultures of Sertoli cells from 8 day-old rats. After isolation the cells were cultured for 24 hrs in a 1:1 mixture of mediums F12: DME with 10 ug/ml transferrin, 5 ug/ml insulin, 5 ug/ml vitamin E, 4 ug/ml hydrocortisone, 100 U/ml penicillin, 2.5 ug/ml amphotericin B. The cells were then cultured for 48 hrs in medium without insulin and immediately afterwards the conditioned media and the cells were collected and analyzed separately. The RIAs employed human recombinant IGF-I and IGF-II for standard and tracer, a polyclonal anti IGF-I antibody and a monoclonal anti IGF-II antibody. To avoid interference by IGF binding proteins previous to RIAs the conditioned media were extracted with a mixture of formic acid-Tween 20-acetone. With the same purpose 8ng/tube IGF-I was added to the IGF-II RIA buffer. Recovery of added 125I IGF-II after the extraction procedure was 101.2 \pm 3.9 % (X \pm SD, n=4). Both IGFs were present in the conditioned media. Serial dilutions of the wextracted media were parallel to the standard curves. IGF-I immunoreactivity in the conditioned media was 84 \pm 20 pg/ug DNA while IGF-II immunoreactivity was 802 \pm 209 pg/ug DNA (X-SEM, n=4). Our results tend to confirm IGF-II production by rat Sertoli cells.

HORMONAL PROFILE OF CATCH UP GROWTH (CUG) IN SMALL FOR GESTATIONAL AGE (SGA) INFANTS. H. García, C. Henriquez, F. Beas, E. Fernandez, G. Ifiguez, E. Trabucco, MA. Boric, F. Barrera, R. Rubio and F. Cassorla, IDIMI. U. de Chile, Medical School and Hospital San Borja - Arriarán, Santiago, Chile.

Our purpose was to compare the endocrine profile of idiopathic SGA infants who grow rapidly during their first months of life and those who fail to do so. 36 SGA infants (17 boys and 19 girls) were followed monthly since birth, measuring the plasma levels of several hormones, at 0 and 3 months. The average birth weight was 2361 \pm 209 grs, and average length was 46 \pm 1.9 cms. Average postnatal growth was 32.6 \pm 9 grs/day; and 3.5 \pm 0.6 cm/month. According to height velocity (HV) the infants were divided into 3 groups: Group 1: (5 boys and 6 girls n:11) with a height velocity >4.1 cms/month (\pm 1 S.D.). Group 2: (11 boys, 7 girls n:18) HV between 2.9-4.1 cm/month (\pm 1 S.D.). Group 3: (6 girls, 1 boy n:7), HV below 2.9 cm/month (\pm 1 S.D.). Hormonal levels at three months of age are shown in the table.

	INSULIN	IGF-I	PRL	T3	T4	TESTOST.	E2
	UUI/ml	ng/ml	ng/ml	ng/dl	ug/dl	ng/ml	pg/ml
1.	17.8 \pm 23	92 \pm 48	19 \pm 10.0	223 \pm 25	11.6 \pm 1.5	0.9 \pm 0.8	76.4 \pm 37
2.	8.3 \pm 7.1	95 \pm 44	23 \pm 15.6	215 \pm 49	10.5 \pm 2.6	0.7 \pm 0.6	44.1 \pm 30
3.	12.0 \pm 16	57 \pm 22	17 \pm 10.6	222 \pm 30	11.5 \pm 1.1	0.2	59.3 \pm 40

CONCLUSIONS: We observed CUG in 30% of these infants during the first three months of life. We found significant differences in IGF-I levels between groups 1 and 2 vs groups 3. These results suggest that IGF-I may play a role in CUG.

JUVENILE AUTOIMMUNE THYROID DISEASE (ATD). FINE NEEDLE ASPIRATION BIOPSY (FNABT) AND IMMUNOMARCATION (PART I). J. Goldberg, V. Herzovich, J. Rossi, S. Iorcansky. Serv. Patol. y Endocrinol. Hosp. Garrahan. Bs.As. Argentina.

FNABT was done in patients with ATD (Chronic Lymphocytic Thyroiditis= CLT and Graves' Disease= GD), and processed according to the method of Hayry at al., applied to follow up transplanted organs. Cytospin samples were stained with May Grunewald-Giemsa, and HLA-DR, CD25, CD4, CD8, CD3, CD11 and surface immunoglobulines were detected by immunofluorescence. In transplanted kidneys, a corrected increment (inflammatory INDEX) (CI) 3,5 indicates immunological activation. We analyzed thyroid samples, obtaining the CI, and the phenotype of infiltrating cells. 56 FNABT were performed on 43 pts. (CLT n=23 and GD n=20), followed up from 1 y. to 4.5y. They were divided in 2 gr., according to their thyroid status. Gr. I: CLT n=29 and Gr. II: GD n=27. Results: (% \pm SD). Gr. I: CI: 6.3 \pm 0.97; B-cells: 33.9 \pm 10.5; T-cells: 61.2 \pm 11.4; CD4: 63.1 \pm 6.2; CD8: 8.34 \pm 4.6. CD4/CD8 ratio: 1.9 \pm 0.5 HLA-DR expression was found on 56.6 \pm 18.8% of the follicular cells and CD25 was expressed in 27/29 pts. Gr. II: CI: 4.2 \pm 1.3 (Gr. I vs II = p<0.001) B-cells: 58.5 \pm 16.2 (p<0.001) T-cells: 39.6 \pm 17.1 (p<0.001) CD4: 59.2 \pm 8.8 (p<0.005) CD8: 40 \pm 9.4 (p<0.02) CD4/CD8 ratio: 1.6 \pm 0.7 (p<0.02). HLA-DR was expressed on 26.6 \pm 17.3 (p<0.001) of follicular cells and CD25 was positive in 7/16 pts. All CLT showed typical cytological aspect. In GD 7/20 cytology 12/20 were normal and 1/20 adenomatous goiter (AG). The CI 5.3, the expression degree of HLA-DR on follicular cells, as well as the expression of CD 25 on lymphocytes are markers of immune activation. TLC could be considered as a form of rejection, in which cellular aggression would play an important role, being in GD mainly humoral.