

Erratum

In the article “Failure of the Loop Diuretic Torasemide to Improve Renal Function of Hypoxic Vasomotor Nephropathy in the Newborn Rabbit” by L. Dubourg *et al.* (*Pediatr Res* 47:504–508, 2000), the last two columns of Table 1 on page 506 did not appear in print and the last item in the first column should read “U_k V” rather than “UK”. The corrected table is reproduced below:

Table 1. Experimental data of the animals of groups 1–3, studied under hypoxic conditions

	Group 1 (n = 8)			Group 2 (n = 9)			Group 3 (n = 8)		
	Control (U1,U2-P3*)	Hypoxemia (U3,U4-P5)	Hypoxemia (U5,U6-P5)	Control (U1,U2-P3)	Hypoxemia (U3,U4-P5)	Hypoxemia + Torasemide (U5,U6-P5)	Control (U1,U2-P3)	Hypoxemia + Torasemide (U3,U4-P5)	Hypoxemia + Torasemide (U5,U6-P5)
pH	7.51 ± 0.02	7.48 ± 0.02	7.48 ± 0.02	7.54 ± 0.01	7.48 ± 0.01 ^b	7.49 ± 0.02 ^a	7.53 ± 0.01	7.51 ± 0.02	7.50 ± 0.02 ^a
PaO ₂ (mm Hg)	132.6 ± 5.4	43.0 ± 1.0 ^a	43.7 ± 1.0 ^a	143.6 ± 3.2	42.4 ± 0.69 ^b	43.1 ± 0.69 ^b	132.3 ± 3.3	42.2 ± 0.25 ^a	42.1 ± 0.35 ^a
PaCO ₂ (mm Hg)	35.4 ± 1.0	33.5 ± 0.6	33.1 ± 0.7 ^a	34.3 ± 0.6	33.4 ± 0.5	31.3 ± 0.3 ^b	35.6 ± 0.5	33.6 ± 0.6 ^a	34.8 ± 0.7
HCO ₃ (mmol/L)	27.7 ± 0.9	24.8 ± 0.8 ^a	24.7 ± 0.9 ^a	28.6 ± 0.6	25.0 ± 0.7 ^b	24.1 ± 0.7 ^b	29.5 ± 0.8	26.7 ± 1.1 ^a	26.9 ± 1.4 ^a
Heart rate (beats/min)	293 ± 13	297 ± 9	298 ± 8	303 ± 9	301 ± 10	320 ± 9	309 ± 6	301 ± 9	316 ± 8
MAP (mm Hg)	33.1 ± 1.8	30.0 ± 2.0 ^b	27.3 ± 1.6 ^a	42.2 ± 1.9	38.6 ± 2.2 ^{b,c}	35.1 ± 2.7 ^b	46.8 ± 1.6	40.5 ± 2.1 ^{b,d}	37.2 ± 2.2 ^{b,d}
GFR (ml/kg/min)	2.20 ± 0.22	2.00 ± 0.21	1.54 ± 0.18 ^a	2.01 ± 0.21	1.90 ± 0.21	1.39 ± 0.17 ^a	2.13 ± 0.16	1.38 ± 0.09 ^{a,c}	0.96 ± 0.12 ^{a,c}
RBF (ml/kg/min)	20.9 ± 1.30	17.6 ± 1.45	13.2 ± 1.4 ^a	16.5 ± 0.87	13.4 ± 1.0 ^{a,c}	9.0 ± 0.74 ^{b,c}	19.8 ± 1.10	12.7 ± 0.8 ^{a,c}	9.6 ± 1.0 ^{a,c}
RVR (mm Hg/ml/kg/min)	1.66 ± 0.11	1.79 ± 0.19	2.38 ± 0.43 ^a	2.59 ± 0.13	2.95 ± 0.15 ^d	4.01 ± 0.26 ^{b,d}	2.41 ± 0.13	3.25 ± 0.20 ^{a,d}	4.12 ± 0.36 ^{a,d}
UV (ml/kg/min)	0.076 ± 0.005	0.074 ± 0.005	0.057 ± 0.005 ^a	0.076 ± 0.006	0.083 ± 0.009	0.279 ± 0.046 ^{a,d}	0.076 ± 0.006	0.181 ± 0.031 ^{a,d}	0.115 ± 0.024 ^{a,c}
U _{Na} V (μmol/kg/min)	2.21 ± 0.65	1.78 ± 0.40	1.04 ± 0.22	2.00 ± 0.85	1.12 ± 0.35	28.76 ± 5.47 ^{b,e}	2.32 ± 0.99	15.89 ± 4.46 ^b	8.65 ± 3.35 ^{b,d}
U _k V (μmol/kg/min)	4.73 ± 1.21	3.66 ± 0.74	2.91 ± 0.44	7.41 ± 1.24	7.91 ± 1.42	10.32 ± 1.83 ^{b,d}	7.24 ± 0.78	8.60 ± 0.96	5.34 ± 0.77 ^{a,c}

For explanation see text (Material and Methods, Fig. 1).

The data (mean ± sem) in each group of animals are the mean of two 30-min urine collections (U1/U2, U3/U4, U5/U6) with or without the administration of torasemide (Torem[®]).

^{a,b} $p < 0.05$, $p < 0.01$, respectively, vs control same group.

^{c,d,e} $p < 0.05$, $p < 0.01$, $p < 0.001$, respectively, vs same experimental period group 1.

* P1–P5 are defined in Figure 1.