4. Likhodii SS 2001 Experiments in the rat pentylenetetrazole infusion threshold model of the ketogenic diet. *Epilepsy Res* 44:83–86
5. Winocur G, Greenwood CE 1999 The effects of high fat diets and environmental influences on cognitive performance in rats. *Behav Brain Res* 101:153–161
6. Greenwood CE, Winocur G 1996 Cognitive impairment in rats fed high-fat diets: a specific effect of saturated fatty-acid intake. *Behav Neurosci* 110:451–459
7. Lloyd HM, Rogers PJ, Hedderley DL, Walker AF 1996 Acute effects on mood and cognitive performance of breakfasts differing in fat and carbohydrate content. *Appetite* 27:151–164
8. Swoboda KJ, Specht L, Jones HR, Shapiro F, DiMauro S, Korson M 1997 Infantile phosphofructokinase deficiency with arthrogryposis: clinical benefit of a ketogenic diet. *J Pediatr* 131:932–934
9. de Vivo DC, Trifiletti RR, Jacobson RI, Ronen GM, Behmand RA, Harik SI 1991 Defective glucose transport across the blood-brain barrier as a cause of persistent hypoglycorrhachia, seizures, and developmental delay. *New Engl J Med* 325: 703–709
10. Wexler ID, Hemalatha SG, McConnell J, Buist NR, Dahl HH, Berry SA, Cederbaum SD, Patel MS, Kerr DS 1997 Outcome of pyruvate dehydrogenase deficiency treated with ketogenic diets. Studies in patients with identical mutations. *Neurology* 49:1655–1661
11. Snead OC 3rd 2004 The ketogenic diet: a cautionary note. *Pediatr Res* 55:368–369

DOI: 10.1203/01.PDR.0000142215.95720.72

Response

To the Editor: We appreciate the interest of Dr. Cunnane and Dr. Likhodii in our recent paper (1). Their letter, which raises a number of excellent points, allows us to make some additional comments about the ketogenic diet and correct one error in our paper.

The first concern by Dr. Cunnane and Dr. Likhodii is that the paper is seriously flawed because of the high fat to protein + carbohydrate ratio of 8.6:1 used in our study. While it is true that this ratio is higher than that typically used in children, we have encountered situations where over-zealous parents have approached this ratio. Moreover, our goal was not to provide an identical diet given to children but rather study the effects of ketosis on cognitive abilities in growing rats. While Dr. Cunnane and Dr. Likhodii would have preferred another diet, our use of this particular diet was not flawed. The diet has been previously used by other investigators (2–5) and, as noted in our paper, the animals tolerated the diet well and gained weight, albeit at a significantly lower rate than that seen in the control animals.

Nevertheless, as we discussed in the paper, we do share the concerns of Dr. Cunnane and Dr. Likhodii about weight gain in relationship to cognitive function. In recent work (6), we have found that caloric restriction during the 1st weeks of life leads to mild cognitive impairment. Interestingly, caloric restriction has been shown to reduce seizure susceptibility (3,5,7). We agree that future studies evaluating the ketogenic diet should use a lower ratio of fat to carbohydrate + protein and avoid the growth failure seen in our animals. The suggestions regarding future study design by Dr. Cunnane and Dr. Likhodii are excellent.

As noted by Dr. Cunnane and Dr. Likhodii, the phenomenon whereby the brain is protected during food deprivation was observed in this study. Nevertheless, brain size did differ among groups. It is not clear why they feel our comments in the text and abstract in regards to brain size are misleading.

Dr. Cunnane and Dr. Likhodii note that the cognitive impairment seen in rats and humans used lower fat concentrations than in our study. We agree and it was for this reason that we have concerns about the cognitive effects of even higher fat

concentrations. Dr. Cunnane and Dr. Likhodii will get little argument from us that in some metabolic disorders therapy with the ketogenic diet can be quite beneficial.

As we discussed in the paper, individuals with epilepsy treated successfully with the ketogenic diet with a marked reduction in seizure frequency or intensity often demonstrate dramatic improvements in cognitive function. In our study we found that significantly fewer rats on the ketogenic diet experienced spontaneous seizures than rats on the control diet (Chi-square test = 3.91, $p < 0.05$). We incorrectly stated in the paper that the results were not statistically significant and apologize for this error. The reduction in seizure susceptibility supports prior work from our laboratory that shows a dissociation between the effects of the ketogenic diet on cognition and seizure susceptibility (2).

Finally, we agree that determining the cognitive effects of the ketogenic diet requires a rigorous evidence-based approach. Regrettably, this should have been done in children years ago.

Gregory L. Holmes
Qian Zhao

Section of Neurology
Dartmouth Medical School
Neuroscience Center at Dartmouth
Lebanon, New Hampshire

Carl E. Stafstrom

Department of Neurology
University of Wisconsin School of Medicine
Madison, Wisconsin

References

1. Zhao Q, Stafstrom CE, Fu DD, Hu Y, Holmes GL 2004 Detrimental effects of the ketogenic diet on cognitive function in rats. *Pediatr Res* 55:498–506
2. Su SW, Cilio MR, Sogawa Y, Silveira DC, Holmes GL, Stafstrom CE, Silveira D 2000 Timing of ketogenic diet initiation in an experimental epilepsy model. *Brain Res Dev Brain Res* 125:131–138
3. Bough KJ, Valiyil R, Han FT, Eagles DA 1999 Seizure resistance is dependent upon age and caloric restriction in rats fed a ketogenic diet. *Epilepsy Res* 35:21–28
4. Bough KJ, Eagles DA 1999 A ketogenic diet increases the resistance to pentylenetetrazole-induced seizures in the rat. *Epilepsia* 40:138–143
5. Bough KJ, Schwartzkroin PA, Rho JM 2003 Caloric restriction and ketogenic diet diminish neuronal excitability in rat dentate gyrus in vivo. *Epilepsia* 44:752–760
6. Akman C, Zhao Q, Liu Z, Holmes GL 2004 Effect of food deprivation during early development on cognition and neurogenesis in the rat. *Epilepsy Behav* 5:446–454
7. Greene AE, Todorova MT, McGowan R, Seyfried TN 2001 Caloric restriction inhibits seizure susceptibility in epileptic EL mice by reducing blood glucose. *Epilepsia* 42:1371–1378

DOI: 10.1203/01.PDR.0000142217.72849.4A

To the Editor: We read with great interest the recent article by Timmons *et al.* (1) on stress/inflammatory responses to exercise in boys and men. While we applaud their efforts to directly compare subjects from these two age groups to research maturational mechanisms of immune responses, one aspect of their data caused us some concern. The authors report that the exercise bout had no effect on circulating interleukin-6 (IL-6) in the boys. This was troubling because of the many cytokines previously reported to be altered acutely by exercise, IL-6 has proved to be the most reproducibly elevated (2). Moreover, we

and others have shown IL-6 to be elevated with exercise in younger children and adolescents (3,4), and Perez Navero *et al.* (5) reported increased IL-6 following a soccer practice in 6- to 7-year-old boys.

One problem that might explain the unexpected negative results in boys lies, perhaps, with the sensitivity of their assay. This is a relevant concern with IL-6 because in healthy individuals at rest normal values in circulating blood are in the range of 1–20 or so pg/ml, whereas with infection or very strenuous exercise, these values can increase by several orders of magnitude.

Timmons *et al.* reported using the commercially available Human IL-6 ELISA system from Endogen (Woburn, MA). Although this assay is listed as having a range from 1–400 pg/ml, the lowest value of the standard curves provided by the company is 10.24 pg/ml. Extrapolating below this level without further dilution and creating an appropriate standard curve could lead to errors. Detecting changes in IL-6 at these low levels, even a doubling or tripling of circulating levels as may occur with light exercise, may not be feasible. Figure 3a in the Timmons article reveals that IL-6 levels were on average greater than 10 pg/ml only for the men post-exercise and in recovery, all of the samples obtained from the boys were, on average, below what appears to be the detectable range of the assay.

The problem of detecting low circulating levels of IL-6 has been observed by other investigators. For example, Ishiguro *et al.* (6) studying acute infections in children recently noted, “Concentrations of IL-6 were quantified by a commercially available ELISA system for infectious patients (Endogen, Woburn, MA, USA). IL-6 was not detectable in most control subjects by this system, and so we used a more sensitive ELISA system.”

Timmons *et al.* also noted that, “In our experience, some samples from the boys (9 of 54 for IL-6. . .) and from the men (7 of 54 for IL-6. . .) were below the detection level of the respective kit and were, therefore, set to the lowest positive number on the standard curve derived from the plate on which the sample was determined.”

In our own investigations of IL-6 responses to exercise in children in adolescents, we noted this sensitivity problem several years ago and when studying healthy subjects, we use an ELISA with a range of 0.56 to 10 pg/ml. The standard curves of this assay permit accurate measurements of IL-6 in a range likely to be encountered in healthy adults and children exposed to moderate exercise for relatively brief periods of time.

Dan M. Cooper
Dan Nemet
Frank Zaldivar, Jr.
Pietro Galassetti

Center for the Study of Health Effects of Exercise
 in Children
 Department of Pediatrics
 University of California, Irvine
 Irvine, CA

References

1. Timmons BW, Tarnopolsky MA, Bar-Or O 2004 Immune responses to strenuous exercise and carbohydrate intake in boys and men. *Pediatr Res* 56:227–234
2. Shephard RJ 2002 Cytokine responses to physical activity, with particular reference to IL-6: sources, actions, and clinical implications. *Crit Rev Immunol* 22:165–182
3. Scheett TP, Mills PJ, Ziegler MG, Stoppani J, Cooper DM 1999 Effect of exercise on cytokines and growth mediators in prepubertal children. *Pediatr Res* 46:429–434
4. Nemet D, Oh Y, Kim HS, Hill MA, Cooper DM 2002 The effect of intense exercise on inflammatory cytokines and growth mediators in adolescent boys. *Pediatrics* 110:681–689
5. Perez Navero JL, Jaraba Caballero S, Ibarra de la Rosa I, Jaraba Caballero MP, Guillen del Castillo M, Montilla Lopez P, Tunez Finana I, Romanos Lezczano A 1999 [Effects of competitive physical exercise on neuroendocrine response and interleukin-6 liberation in children]. *An Esp Pediatr* 51:267–272
6. Ishiguro A, Suzuki Y, Mito M, Shimbo T, Matsubara K, Kato T, Miyazaki H 2002 Elevation of serum thrombopoietin precedes thrombocytosis in acute infections. *Br J Haematol* 116:612–618

DOI: 10.1203/01.PDR.0000142214.88096.CE

Response

To the Editor: We would like to thank Cooper and colleagues for their letter regarding our recent article (1). In their letter, Cooper and colleagues raised a concern regarding our method of measuring plasma interleukin-6 (IL-6) and the sensitivity of the chosen assay. The authors are correct to point out that the ELISA kit from ENDOGEN suggests a standard curve between 10.24 and 400 pg/mL. We too were aware that plasma levels of IL-6 in humans tend to be low and, therefore, included a dilution to 5.12 pg/mL and excluded the highest 400 pg/mL concentration from each standard curve. With this approach our correlation coefficients for the standard curve regression equations ranged from 0.9954 to 0.9971 demonstrating excellent linearity and, therefore, providing confidence in extrapolating values between 0 and 5.12 pg/mL.

However, based on the concern of Cooper and colleagues and their assertion that exercise should increase circulating levels of IL-6 in children, we returned to our raw data and analyzed the absorbance units derived from the plate reader, before submitting to the standard curve equations. With this approach, our conclusions remained the same insofar as the absorbances after exercise ($p = 1.0$) and during recovery ($p = 0.9$) were not different than at rest for the boys, but were significantly higher after exercise ($p = 0.001$) and during recovery ($p = 0.0009$) than at rest in the men.

In their letter, Cooper and colleagues point out that IL-6 is reproducibly elevated with exercise in children and apparently dismiss our findings as “unexpected negative results.” We would like to respectively address the evidence as provided by the authors. First, the study by Perez Navero *et al.* (2) apparently measured IL-6 levels in saliva, which is not relevant to the discussion of circulating IL-6 levels. Second, Nemet *et al.* (3) investigated cytokine responses in boys ranging in age from 14 to 18.5 y (mean age of 16.5). It is very likely that most, if not all, of these boys were in advanced stages of puberty. If, as we hypothesized in our paper, maturity status plays a role in the magnitude of exercise-induced immune changes, one would expect this