Gender Differences in Cerebral Blood Flow Velocity and Autoregulation between the Anterior and Posterior Circulations in Healthy Children

MONICA S. VAVILALA, M. SEAN KINCAID, SAIPIN L. MUANGMAN, PILAR SUZ,
IRENE ROZET, AND ARTHUR M. LAM

*Department of Anesthesiology [M.S.V., M.S.K., S.L.M., P.S., I.R., A.M.L.], Pediatrics [M.S.V.] and
Neurological Surgery [M.S.V., A.M.L.], Harborview Medical Center, Seattle, WA 98104*

ABSTRACT

There is little information on gender differences in cerebral autoregulation. The purpose of this study was to compare autoregulation of the anterior and posterior circulations using the tilt test method in healthy boys and girls who were 10–16 y of age. Transcranial Doppler was used to measure middle cerebral artery and basilar artery flow velocities (Vmca and Vbas). Cerebral autoregulation (ARI) of the middle cerebral (ARImca) and basilar arteries (ARIBas) was examined using the tilt test method. An $ARI < 0.4$ indicates impaired autoregulation. Among the 13 boys and 13 girls, Vmca and Vbas were higher in girls. All children demonstrated intact autoregulation, but boys had higher ARImca than girls, whereas girls had higher ARIBas than boys. Girls demonstrated greater autoregulation in the basilar artery, whereas boys demonstrated greater autoregulation in the middle cerebral artery. Girls had higher flow velocities in both vessels. This study provides normative data on cerebral autoregulation of the poste-

rior circulation in healthy, awake boys and girls. (*Pediatr Res* 58: 574–578, 2005)

Abbreviations

ARImca, middle cerebral artery autoregulatory index

ARIBas, basilar artery autoregulatory index

BAS, basilar artery

CBF, cerebral blood flow

LLA, lower limit of autoregulation

MAP, mean arterial pressure

MAPE, mean arterial pressure at the external auditory meatus

MCA, middle cerebral artery

TCD, transcranial Doppler

Vmca, middle cerebral artery flow velocity

Vbas, basilar artery flow velocity

Autoregulation of cerebral blood flow (CBF) is a physiologic and homeostatic process that maintains nearly constant CBF over a range of mean arterial pressures (MAPs). Disease states, including traumatic brain injury, can impair cerebral autoregulation, rendering the brain susceptible to inadequate (cerebral ischemia) or excessive (cerebral hyperemia) CBF (1). Despite its critical role in maintaining CBF, there is limited information on cerebral autoregulation in healthy children.

Little has been published on pediatric cerebral autoregulation in children outside the clinical arena of neurogenic syncope (2–4). In one study that examined dynamic cerebral autoregulation in awake adolescent study participants, the time to return of middle cerebral artery flow velocity (Vmca) to normal after a transient hypotensive stimulus was reported to be more in healthy adolescents compared with their healthy adult counterparts (5). How-

ever, in a subsequent evaluation of cerebral autoregulation in children *versus* adults during general anesthesia using static autoregulation testing, the investigators reported no age-related differences in autoregulatory capacity and no difference in cerebral autoregulation compared with adults (6). Because both of these studies examined cerebral autoregulation of the anterior circulation only, differences in cerebral autoregulation between the anterior and posterior circulation in children could not be evaluated. In addition, to our knowledge, there is no information regarding cerebral autoregulation of the posterior circulation in children, no information on gender differences in cerebral autoregulation in children, and finally no normative data on cerebral autoregulation using the tilt test method. Therefore, the purpose of this study was to 1) provide normative data on cerebral autoregulation using the tilt test method, 2) describe cerebral autoregulation of the posterior cerebral circulation, and 3) examine gender-related differences in autoregulatory capacity in healthy, awake boys and girls.

METHODS

Study participants and setting. Healthy children who were 10 to 16 y of age and had no history of seizures, syncope, dysautonomia, or any neurologic/

Received November 29, 2004; accepted February 3, 2005.

Correspondence: Monica S. Vavilala, M.D., Department of Anesthesiology, Harborview Medical Center, 325 Ninth Avenue, Box 359724, Seattle, WA 98104; e-mail: vavilala@u.washington.edu.

This research was supported by the NIH K23 HD044632-02 (to M.S.V.).

DOI: 10.1203/01.PDR.0000179405.30737.0F

cardiac disorders were eligible for participation in this study. Eligible study participants were recruited from well-child visits in the general pediatrics clinic at Harborview Medical Center (Seattle, WA). Informed consent for participation was obtained from the parent or guardian, and age-appropriate assent was obtained from each child.

Physiologic testing was conducted in either the Harborview Cerebrovascular Laboratory or the Post-Anesthesia Care Unit at Harborview Medical Center. This study was approved by the University of Washington's institutional review board.

Study design and protocol. Each participant was positioned on a bed with the head and back adjustable for elevation. An appropriately sized noninvasive blood pressure cuff was placed on one arm. Transcranial Doppler (TCD) ultrasonography (Multidop X; DWL Corp., Sipplingen, Germany) was used to measure flow velocities in the middle cerebral (Vmca) and basilar (Vbas) arteries. A hand-held 2-MHz ultrasound probe was used to insonate the desired vessel and positioned for sufficient time to achieve steady-state measurement. Previously age appropriate depths were used to insonate both the middle cerebral artery (MCA) and basilar artery (BAS) (7). During steady-state conditions, Vmca, Vbas, MAP, heart rate, and respiratory rate were recorded in each position.

For examining cerebral autoregulation, change in position proceeded from supine to sitting upright (90 degrees). For the sitting upright position, the vertical distance between the noninvasive blood pressure cuff and the external auditory meatus was used to calculate the estimated MAP (MAPE) at the Circle of Willis. Because MAP decreases by 1 mm Hg for every 1.36-cm increase in vertical height, the change in height from supine to sitting upright was divided by 1.36 cm to calculate the MAPE in the sitting upright position.

Cerebral autoregulation was calculated off-line as previously described (8). Briefly, cerebral autoregulation was quantified using the autoregulatory index (ARI), where $ARI = \% \Delta eCVR / \% \Delta MAPe$; eCVR is the estimated cerebrovascular resistance calculated as the ratio of MAP to Vmca or Vbas as appropriate (8). Two ARIs were calculated for each participant: one for the middle cerebral artery (ARImca) and one for the basilar artery (ARibas).

Sample size calculations. Sample size calculation was based on previously published ARI data (6). A 30% difference in ARI of boys and girls between the MCA and BAS was considered clinically significant. Accepting an α of 0.05 and a β of 0.8, power analysis indicated the need for a minimum of 24 study participants. Two-factor ANOVA was used to compare 1) ARImca versus ARibas and 2) Vmca versus Vbas between boys and girls. Significance was set at $p < 0.05$. Age-related Vmca and Vbas in this study were compared with age-related normative data previously published by Bode (7). All data are presented as mean $\pm$ SD.

RESULTS

Twenty-six children (13 boys, 13 girls) between the ages of 10 and 16 y (boys: 12.9 ± 1.7 ; girls: 12 ± 1.4 ; $p = 0.14$) completed the study. Twenty-four of 26 children (12 of 13 boys and 12 of 13 girls) were pubertal (Tanner II–IV). Two boys and one girl were Tanner stage IV. Therefore, 21 of 26 participants were either Tanner II or III. There was no difference in the distribution of Tanner staging between boys and girls. There was no significant difference in MAPE, drop in MAPE with autoregulation testing, Vmca, Vbas, ARImca, ARibas, or the relationships examined when the two prepubertal children were excluded from the analyses. Therefore, the final analyses include data from all 26 participants.

At baseline (supine), boys had a higher MAPE than girls (79 ± 7 mm Hg for boys *versus* 75 ± 5 mm Hg for girls; $p = 0.004$). In the sitting upright position, MAPE decreased in both boys and girls (55 ± 8 mm Hg in boys *versus* 54 ± 8 mm Hg in girls; $p = 0.47$). The average drop in MAPE for the whole group was 22 ± 6 mm Hg, from 77 ± 6 to 55 ± 8 mm Hg ($p < 0.001$). There was no statistically significant difference in drop in MAPE between boys and girls (boys 23 ± 5 mm Hg *versus* girls 21 ± 6 mm Hg; $p = 0.12$). All study participants achieved at least a 10-mm Hg drop in MAPE with the position change. Respiratory rate was similar for both boys and girls in the supine (both 20 breaths/min; $p = 0.23$) and sitting upright positions (boys 20 breaths/min

versus girls 19 breaths/min; $p = 0.94$). Vmca, Vbas, and ARI data for individual participants are given in Tables 1 and 2. For the entire cohort, baseline (supine) Vmca and Vbas were 82 ± 17 and 55 ± 12 cm/s, and upright Vmca and Vbas were 81 ± 18 and 55 ± 12 cm/s (Vmca supine *versus* upright, $p = 0.26$; Vbas supine *versus* upright, $p = 0.89$). However, baseline Vmca (89 ± 16 cm/s) and Vbas (59 ± 11 cm/s) were higher in girls compared with boys (Vmca 75 ± 16 cm/s and Vbas 51 ± 12 cm/s; $p = 0.005$; Table 3, Fig. 1). In the sitting upright position, Vmca (87 ± 17 cm/s) and Vbas (60 ± 11 cm/s) were also higher in girls compared with boys (Vmca 75 ± 17 cm/s and Vbas 50 ± 12 cm/s; $p = 0.01$; Table 3).

All study participants had intact cerebral autoregulation of the MCA and BAS (Tables 1 and 2). For the whole cohort, there was no difference between ARImca and ARibas (ARImca: 0.95 ± 0.08 , ARibas: 0.94 ± 0.10 ; $p = 0.78$). There was also no overall difference in ARI for both MCA and BAS by gender (0.95 ± 0.09 for boys *versus* 0.95 ± 0.08 for girls; $p = 0.86$). However, boys had significantly higher ARImca than girls, whereas girls had higher ARibas than boys (ARImca: boys 0.98 ± 0.03 *versus* girls 0.92 ± 0.1 ; ARibas: boys 0.92 ± 0.12 *versus* girls: 0.97 ± 0.06 ; $p = 0.02$; Table 4, Fig. 2).

Table 1. Middle Cerebral Artery Flow Velocity (Vmca) and Autoregulatory Index (ARI) in Girls and Boys 10 to 16 Years old in the Supine and Sitting Upright Positions

Girls							
Age (y)	Supine MAPE	Supine Vmca	Supine CVRe	Upright MAPE	Upright Vmca	Upright CVRe	ARI
10	74	90	0.82	58	98	0.59	1
11	74	68	1.09	55	66	0.83	0.91
11	71	88	0.81	40	83	0.48	0.92
11	85	73	1.16	71	73	0.97	1
11	77	97	0.79	54	98	0.55	1
12	84	94	0.89	68	93	0.73	0.95
12	73	100	0.73	46	98	0.47	0.97
12	73	106	0.69	47	103	0.46	0.95
12	72	87	0.83	52	77	0.68	0.66
12	72	99	0.73	54	92	0.57	0.77
13	78	110	0.71	49	110	0.45	1
14	77	90	0.86	58	88	0.66	0.93
15	78	51	1.53	45	48	0.94	0.91

Boys							
Age (y)	Supine MAPE	Supine Vmca	Supine CVRe	Upright MAPE	Upright Vmca	Upright CVRe	ARI
10	86	83	1.04	69	83	0.83	1
10	69	100	0.69	46	100	0.46	1
11	80	87	0.92	62	86	0.72	0.96
12	84	84	1	63	85	0.74	1
13	81	62	1.31	43	62	0.69	1
13	89	100	0.89	56	104	0.54	1
13	79	76	1.04	62	82	0.76	1
14	76	53	1.43	50	51	0.98	0.92
14	76	76	1	51	77	0.66	1
14	75	48	1.56	47	47	1	0.96
14	84	69	1.22	59	67	0.88	0.93
15	83	64	1.30	58	64	0.90	1
15	75	73	1.03	50	72	0.69	0.97

MAPE = mean arterial pressure at the external auditory meatus, CVRe = estimated cerebrovascular resistance.

Table 2. Basilar Artery Flow Velocity (Vbas) and Autoregulatory Index (ARI) in Girls and Boys 10 to 16 Years old in the Supine and Sitting Upright Positions

Girls							
Age (y)	Supine MApe	Supine Vbas	Supine CVRe	Upright MApe	Upright Vbas	Upright CVRe	ARI
10	71	65	1.09	59	65	0.91	1
11	74	52	1.42	60	50	1.2	0.83
11	73	66	1.11	55	71	0.77	1
11	86	68	1.26	67	66	1.02	0.89
11	75	50	1.5	58	55	1.05	1
12	80	52	1.54	62	53	1.17	1
12	65	75	0.87	44	72	0.61	0.91
12	70	65	1.08	43	66	0.65	1
12	73	63	1.16	45	65	0.69	1
12	70	70	1	56	70	0.8	1
13	71	57	1.25	48	61	0.79	1
14	78	34	2.29	63	34	1.85	1
15	73	56	1.30	47	55	0.85	0.97

Boys							
Age (y)	Supine MApe	Supine Vbas	Supine CVRe	Upright MApe	Upright Vbas	Upright CVRe	ARI
10	86	61	1.40	69	60	1.15	0.93
10	67	70	0.96	43	70	0.61	1
11	69	50	1.38	53	57	0.93	1
12	84	58	1.45	63	59	1.07	1
13	74	46	1.61	57	42	1.36	0.68
13	96	63	1.52	68	63	1.08	1
13	76	50	1.52	54	48	1.13	0.9
14	75	40	1.88	58	41	1.41	1
14	76	64	1.19	51	64	0.80	1
14	70	31	2.26	44	37	1.19	1
14	81	40	2.03	52	39	1.33	0.95
15	83	41	2.02	58	38	1.53	0.82
15	71	45	1.58	49	39	1.26	0.66

MApe = mean arterial pressure at the external auditory meatus, CVRe = estimated cerebrovascular resistance.

Table 3. Supine and Upright Flow Velocities by Gender

Gender	Supine Flow Velocity (cm/s)		Upright Flow Velocity (cm/s)	
	Vmca	Vbas	Vmca	Vbas
Boys	75 ± 16	51 ± 12	75 ± 17	50 ± 12
Girls	89 ± 16	59 ± 11	87 ± 17	60 ± 11
Mean ± SD	82 ± 17	55 ± 12	81 ± 18	55 ± 12
	boys vs. girls	p = 0.005	boys vs. girls	p = 0.01

Vmca = middle cerebral artery flow velocity; Vbas = Basilar artery flow velocity. Girls have higher Vmca and Vbas than boys in both the supine and upright positions.

DISCUSSION

In this study of healthy children who were 10 to 16 y of age, boys demonstrated greater autoregulation of the MCA than girls, whereas girls had greater autoregulation of the BAS than boys. Girls also had higher flow velocities of both the MCA and the BAS than boys. The present study provides normative data on cerebral autoregulation of both the anterior and posterior circulations in healthy children using the tilt test method. To the best of our knowledge, this is the first description of cerebral autoregulation of the posterior circulation and the first description of gender-related differences in CBF velocity and cerebral autoregulation in children.

Figure 1. Vmca and Vbas for boys (shaded columns; n = 13) and girls (black columns; n = 13). Girls have higher Vmca and Vbas than boys (*p = 0.005). Vmca is higher than Vbas for both boys and girls. Data are presented as mean ± SD.**Table 4.** Cerebral Autoregulation by Gender

Gender	Autoregulatory Index		
	ARImca	ARibas	Mean ± SD
Boys	0.98 ± 0.03	0.92 ± 0.12	0.95 ± 0.09
Girls	0.92 ± 0.10	0.97 ± 0.06	0.95 ± 0.08
Mean ± SD	0.95 ± 0.08	0.94 ± 0.10	

p = 0.02

Girls have higher ARImca than boys whereas boys have higher ARibas than girls. ARI = autoregulatory index; MCA = middle cerebral artery; BAS = basilar artery.

Figure 2. ARImca and ARibas for boys (black columns; n = 13) and girls (shaded columns; n = 13). Girls had higher ARibas, whereas boys had better ARImca (p = 0.02). Data are presented as mean ± SD

We examined the effect of gender on cerebral autoregulation because there are known differences between the central nervous systems and the hormonal milieu of growing children of different genders (9). In addition, estrogen has been shown to correlate directly with CBF velocity (10). Clinically, the overall higher flow velocities in both the MCA and the BAS in pubertal girls between 10 and 16 y of age in this study are consistent with similar gender differences reported by Bakker *et al.* (11) in adults. We chose to study children who were 10 to 16 y of age because Bode (7) grouped Vmca and Vbas data for older children in this age range, thereby allowing us to compare our data with this referenced information. However, Bode did not analyze flow velocities according to gender. The higher flow velocities in girls in the MCA and BAS suggest that the gender-related differences in Vmca reported in adults are present in both the anterior and the posterior circulations before adulthood.

When examined without reference to gender, the present study is consistent with previous findings in adults of equivalent autoregulatory capacity in the anterior and posterior cir-

culations (12). Although we report that pubertal boys demonstrated greater autoregulation of the MCA, whereas pubertal girls demonstrated greater autoregulation of the BAS, the present differences are small and the clinical relevance is not yet known. However, differences in the hormonal milieu may account for the observed differences in autoregulation. For example, estradiol has been shown to exhibit tissue selectivity in its vasodilation in rabbit arteries (13). For example, one might expect, from a teleologic perspective, greater autoregulation in all humans in a vessel that supplies the brainstem. One might also expect a global change in autoregulation as a result of the cerebral vasodilatory properties of estrogen. In this study, gender-related differences in flow velocities (higher flow velocities and lower cerebrovascular resistance) do not seem to account solely for regional differences in autoregulatory capacity patterns between boys and girls. Like the reported gender-related differences in flow velocities, gender differences in autoregulatory patterns seem to exist before adulthood.

In the present study, we measured flow velocities and calculated ARI for tilt from supine to the sitting upright position, which resulted in a 22-mm Hg average drop in MAPE. We chose to use the sitting upright position (90 degrees) because we were concerned that the potentially large drop in MAPE with standing may approach the lower limit of autoregulation (LLA). The decrease in MAPE from supine to sitting upright in this age group corroborates our previous work that the LLA of the MCA in older children without neurologic disease is <55 mm Hg and that the autoregulatory reserve in children this age group is large (14,15).

Comparatively, little is known about cerebral autoregulation of the posterior circulation. In one study of 16 healthy, awake adults, Park *et al.* (12) reported no difference in ARI between the BAS and the MCA. Unlike Park's study, we used a static autoregulation testing method, in which the decrease in blood pressure is sustained. In addition, other studies compared the MCA with the posterior cerebral artery but not the BAS (16,17). We examined the BAS primarily because it supplies the posterior fossa contents as opposed to the posterior cerebral artery, which supplies the occipital cortex (18).

It is important to understand cerebral autoregulation of the posterior circulation because certain disease processes may affect the posterior circulation independent of the anterior circulation. Although strokes in children are uncommon, posterior circulation strokes are more common in previously healthy boys (19). The results from this study question whether regional differences in cerebral autoregulation (less well-developed cerebral autoregulatory mechanisms of the BAS) may play a role in this pattern. In addition, conditions such as traumatic dissection, sickle cell disease, and prothrombotic states may engender symptomatic lesions and impair cerebral autoregulation in one distribution and not the other (20–23).

Cerebral autoregulation can be evaluated quantitatively using several different techniques. Although static cerebral autoregulation testing, using TCD, has been validated and can be performed using pharmacologic intervention to increase MAPE, it may be too invasive for awake, healthy children (6). Tilt testing allows cerebral autoregulation to be studied at the bedside noninvasively, is well tolerated by children, and has

been used for many years in the evaluation of children and adults with syncope (24,25). It has also been combined with TCD ultrasonography to evaluate changes in cerebral vascular resistance in patients with dysautonomia (26).

There are some limitations to this study. First, we measured CBF velocity, not CBF. The study was conducted during steady state by investigators with experience in using TCD technology, and the flow velocities in this study correlate well with the normal values previously reported by Bode (7). The advantage of the present study over the previous study is the removal of potential confounders, such as the effect of volatile anesthetic on cerebrovascular tone (27–29). Although we could estimate the drop in MAPE with tilt testing, we could not control the blood pressure to ensure a specified drop in MAPE. However, the range of MAPE tested (77–55 mm Hg) was within physiologic norms and sufficiently broad to precipitate an autoregulatory response. We are unable to comment on potential differences in LLA between the anterior and the posterior circulations between boys and girls because we could not match the LLA. In addition, we evaluated only the MCA and the BAS, not the other cerebral arteries. Although we did not use capnography to measure end-tidal CO₂ and cannot exclude the influence of changes in tidal volume (surrogate for partial pressure of arterial CO₂) on change in V_{mca} and V_{bas}, we measured respiratory rate during steady state and found the rate to be similar whether supine or sitting upright. We measured respiratory rate and MAPE in each position to examine whether changes in minute ventilation were dependent on position, because partial pressure of arterial CO₂ and sympathetic tone both potentially influence V_{mca} and V_{bas}. Similarly, we could not control cerebral metabolic rate or flow-metabolism coupling in our awake study participants. To maintain steady-state conditions, we made attempts to minimize distractions during the study, and we ensured that flow velocities reached steady state before we recorded it. We did not use invasive arterial blood pressure monitoring. However, because the study participants were healthy children without vascular disease, we believe our results to be valid. Finally, we did not obtain hormone levels from our participants, precluding us from definitively attributing our observations of cerebrovascular physiology on hormonal rather than genetic influences.

CONCLUSION

In this study of children who were 10 to 16 y of age, girls demonstrated greater autoregulation of the BAS than boys, whereas boys demonstrated greater autoregulation of the MCA than girls. Like their adult counterparts, girls in this age group had higher CBF velocities of both the MCA and the BAS. It is possible that as we recognize age-related changes in pediatric CBF, CBF and cerebral autoregulation may now have to be considered with respect to both age and gender. This study provides new information on gender differences in CBF velocities and cerebral autoregulation of the anterior and posterior circulations, as well as normative data on autoregulatory indices using the tilt test method in healthy children. This information may be useful to researchers who study and clinicians who manage children with and without neurologic disease.

Acknowledgments. We thank the University of Washington's Department of Pediatrics, the nurses of the Harborview Medical Center Post-Anesthesia Care Unit, the staff of the Harborview Medical Center Children and Teens Clinic, Dominique Calhoun, and, most important, the children and their families who participated in this study for support of this research.

REFERENCES

- Obrist WD, Gennarelli TA, Segawa H, Dolinskas CA, Langfitt TW 1979 Relation of cerebral blood flow to neurological status and outcome in head-injured patients. *J Neurosurg* 51:292–300
- Albina G, Fernandez Cisneros L, Laino R, Nobo UL, Ortega D, Schwarz E, Barja L, Lagos R, Giniger A, Ameriso SF 2004 Transcranial Doppler monitoring during head upright tilt table testing in patients with suspected neurocardiogenic syncope. *Eurpace* 6:63–69
- Sung RY, Du ZD, Yu CW, Yam MC, Fok TF 2000 Cerebral blood flow during vasovagal syncope induced by active standing or head up tilt. *Arch Dis Child* 82:154–158
- Rodriguez-Nunez A, Couceiro J, Alonso C, Eiris J, Fuster M, Sanchez L, Martinon JM 1997 Cerebral oxygenation in children with syncope during head-upright tilt test. *Pediatr Cardiol* 18:406–409
- Vavilala MS, Newell DW, Junger E, Douville CM, Aaslid R, Rivara FP, Lam AM 2002 Dynamic cerebral autoregulation in healthy adolescents. *Acta Anaesthesiol Scand* 46:393–397
- Vavilala MS, Lee LA, Lee M, Graham A, Visco E, Lam AM 2003 Cerebral autoregulation in children during sevoflurane anaesthesia. *Br J Anaesth* 90:636–641
- Bode H 1988 *Pediatric Applications of Transcranial Doppler Sonography*. Springer-Verlag, New York, pp 114
- Tiecks FP, Lam AM, Aaslid R, Newell DW 1995 Comparison of static and dynamic autoregulation measurements. *Stroke* 26:1014–1019
- Giedd JN 2004 Structural magnetic resonance imaging of the adolescent brain. *Ann NY Acad Sci* 1021:77–85
- Shamma FN, Fayad P, Brass L, Sarrel P 1992 Middle cerebral artery blood velocity during controlled ovarian hyperstimulation. *Fertil Steril* 57:1022–1025
- Bakker SL, de Leeuw FE, den Heijer T, Koudstaal PJ, Hofman A, Breteler MM 2004 Cerebral haemodynamics in the elderly: the Rotterdam study. *Neuroepidemiology* 23:178–184
- Park CW, Sturzenegger M, Douville CM, Aaslid R, Newell DW 2003 Autoregulatory response and CO₂ reactivity of the basilar artery. *Stroke* 34:34–39
- Salom JB, Burguete MC, Perez-Asensio FJ, Torregrosa G, Alborch E 2001 Relaxant effects of 17-beta-estradiol in cerebral arteries through Ca(2+) entry inhibition. *J Cereb Blood Flow Metab* 21:422–429
- Vavilala MS, Lee LA, Lam AM 2003 The lower limit of cerebral autoregulation in children during sevoflurane anesthesia. *J Neurosurg Anesthesiol* 15:307–312
- Ramaekers VT, Casaer P, Daniels H, Marchal G 1990 Upper limits of brain blood flow autoregulation in stable infants of various conceptional age. *Early Hum Dev* 24:249–258
- Rosengarten B, Kaps M 2002 Cerebral autoregulation in middle cerebral artery territory precedes that of posterior cerebral artery in human cortex. *Cerebrovasc Dis* 13:21–25
- Haubrich C, Wendt A, Diehl RR, Klotzsch C 2004 Dynamic autoregulation testing in the posterior cerebral artery. *Stroke* 35:848–852
- Jatuzis D, Zachrisson H, Blomstrand C, Ekholm S, Holm J, Volkmann R 2000 Evaluation of posterior cerebral artery blood flow with transcranial Doppler sonography: value and risk of common carotid artery compression. *J Clin Ultrasound* 28:452–460
- Ganesan V, Chong WK, Cox TC, Chawda SJ, Prengler M, Kirkham FJ 2002 Posterior circulation stroke in childhood. *Neurology* 59:1552–1556
- Rommel O, Niedeggen A, Tegenthoff M, Kiwitt P, Botel U, Malin J 1999 Carotid and vertebral artery injury following severe head or cervical spine trauma. *Cerebrovasc Dis* 9:202–209
- tk;2Verdu A, Cazorla MR, Granados MA, Alonso JA, Casado LF 2001 Basilar artery thrombosis in a child heterozygous for factor V Leiden mutation. *Pediatr Neurol* 24:69–71
- Riebel T, Kebelmann-Betzing C, Gotze R, Overberg US 2003 Transcranial Doppler ultrasonography in neurologically asymptomatic children and young adults with sickle cell disease. *Eur Radiol* 13:563–570
- Paulson OB, Strandgaard S, Edvinsson L 1990 Cerebral autoregulation. *Cerebrovasc Brain Metab Rev* 2:161–192
- Lerman-Sagie T, Rechavia E, Strasberg B, Sagie A, Blieden L, Mimouni M 1991 Head-up tilt for the evaluation of syncope of unknown origin in children. *J Pediatr* 118:676–679
- Grossi D, Nozzoli C, Roca ME, Santostasi R, Simone F 1987 Head-up tilt for triggering and diagnosing syncope. *Funct Neurol* 2:457–464
- Hilz MJ, Axelrod FB, Haertl U, Brown CM, Stemper B 2002 Transcranial Doppler sonography during head up tilt suggests preserved central sympathetic activation in familial dysautonomia. *J Neurol Neurosurg Psychiatry* 72:657–660
- Cremer OL, Diephuis JC, van Soest H, Vaessen PH, Bruens MG, Hennis PJ, Kalkman CJ 2004 Cerebral oxygen extraction and autoregulation during extracorporeal whole body hyperthermia in humans. *Anesthesiology* 100:1101–1107
- Blaha M, Aaslid R, Douville CM, Correra R, Newell DW 2003 Cerebral blood flow and dynamic cerebral autoregulation during ethanol intoxication and hypercapnia. *J Clin Neurosci* 10:195–198
- Strebel S, Lam AM, Matta B, Mayberg TS, Aaslid R, Newell DW 1995 Dynamic and static autoregulation during isoflurane, desflurane, and propofol anesthesia. *Anesthesiology* 83:66–76