teeth using a combination of serial microCT scans and SEM have enabled us to reconstruct the chronology and extent of intrauterine and infant growth insults in DS infants, expressed in growth and differentiation of the developing teeth. Our results for the deciduous teeth show that in DS, there is accelerated dental development in the first trimester followed by progressive growth retardation associated with impaired differentiation of the dental tissues. These changes appear to primarily affect cells derived from ectoderm affecting cell division and differentiation resulting in a smaller modified tooth germ and impaired enamel matrix formation and bio-mineralization. The resulting enamel is thin and hypomineralized and this may contribute to the severe abrasion seen in the teeth of DS children despite the associated lack of muscle tone.

830

USE OF EMERGENCY O NEGATIVE BLOOD TRANSFUSION IN PAEDIATRIC INTENSIVE CARE UNIT

N. Shettihalli¹, A. Shefler¹, S. Stanworth²

¹Paediatric Intensive Care Unit, ²Hematology, John Radcliffe Hospital, Oxford, UK

Background: In emergency situations O negative blood (O-) transfusion may be required before standard compatibility tests can be completed. An increase in the use of emergency O- in our paediatric intensive care unit (PICU) triggered an audit to evaluate our approach and measure this against any identified guidelines.

Methods: PICU patients issued emergency O- between January 2004 and December 2008 were identified. Case notes were analysed to ascertain the indications for emergency transfusion, time from PICU admission to transfusion, volume and duration of transfusion and the presence of any transfusion reaction.

Results: Eight patients received a total of 10 units. Two patients were transfused twice. The patients median age was six months. Visible active bleeding with low cardiac output was recorded as an indication on three occasions. Five transfusions occurred in children with low cardiac output and suspected but not proven active bleeding. Two further patients were transfused for low Hb only. Up to 20 ml/kg were given as a bolus over 10 to 20 minutes in all but two cases where it was transfused over 90 and 150 minutes. All patients had been admitted to PICU

more than four hours previously. In three instances, crossmatched blood was already available. No transfusion reactions were documented.

Conclusions: Published data is scarce with regards to the use of emergency O-. Our audit demonstrates the variety and inconsistency of practise. Improved local and national guidance would lead to better understanding and safe transfusion practise in the PICU and wider acute paediatric setting.

831

PATTERNS OF MINOR BLEEDING IN NEONATES WITH SEVERE THROMBOCYTOPENIA: A PROSPECTIVE OBSERVATIONAL STUDY

V. Venkatesh¹, P. Muthukumar¹, S. Stanworth², P. Clarke³, L. Choo⁴, A. Curley¹, PlaNet 1 Study Group

¹NICU, Cambridge University Hospital NHS Foundation Trust, Cambridge, ²NHSBT, Oxford, ³NICU, Norfolk & Norwich University Hospitals NHS Trust, Norwich, ⁴Clinical Trials Unit, Medical Research Council, London, UK

Background and aim: A prospective cross sectional observational study to determine incidence and pattern of minor bleeding in neonates with severe neonatal thrombocytopenia (SNT) defined as platelet count < 60 x10⁹/L.

Methods: Neonates admitted to seven UK tertiary neonatal units (March 2005 -November 2006) documented to have SNT were enrolled for daily data collection.

Results: 169 babies with SNT were enrolled for 2055 study days. 123 (73%) babies showed minor bleeding, 15 (9%) minor and major bleeds, 31 (18%) no evidence of bleeding. Bleeding as a percentage of study days occurred with greatest frequency in extremely preterm infants: < 28 weeks, 44%; 28-34 weeks, 27%; >34 weeks, 28%. Neonates with early SNT (< 3 days) most commonly had hematuria (31%), oozing (20%) and/or nasogastric bleeding (17%). Neonates with late SNT (≥3 days) most commonly showed hematuria (42%), minor bleeding from the endotracheal tube (24%) and oozing (13%). Minor bleeding (bleeding days as a percentage of study days) with SNT was commoner in the first 10 days (50%) compared to >10 days (33%). Neonates with major bleeds did not show a higher incidence of minor bleeding events prior to the major haemorrhage. 41% of infants with nadir