Methods: Iranian CCHF probable children's sera were collected from 2000 to 2009. They were analyzed by specific IgM and IgG ELISA and RT-PCR (Gel-Based and Real-Time) assays.

Results: From 2000 to 2009, sera were collected of children with age range of 2 months to 14 years. The number of probable, confirmed, and death cases according to the year are respectively 2000 (10,1,0), 2001 (22,5,0), 2002 (18,5,0), 2003 (18,5,1), 2004 (7,1,0), 2005 (5,0,0), 2006(10, 2,0), 2007(8,3,0), 2008 (28, 9,2) and 2009 (24,2,1). Among the confirmed cases, 21 were boy and 12 were girl. This study demonstrated that the most infected province was Sistan-va-Balouchestan with 34.7% of confirmed cases.

Conclusions: Because of neighboring to Pakistan and Afghanistan, Sistan-va-Baluchistan is the most infected province of Iran. Therefore, the children resident in the infected provinces are in high risk to CCHF. Also, we observed that their main ways of contamination were contact with tissue or blood of infected livestock, so informing children and their parents about routes of transmission will be useful for control of CCHF in the infected regions.

902

ROUTE OF TRANSMISSION OF CRIMEAN-CONGO HEMORRHAGIC FEVER (CCHF) DISEASE TO IRANIAN CHILDREN

S. Chinikar¹, S.M. Ghiasi¹, M. Moradi¹, M.M. Goya², M.R. Shirzadi², M. Zeinali²

¹Arboviruses and Viral Hemorrhagic Fever Laboratory(National Reference Lab.), Pasteur Institute of Iran, ²Center for Disease Control, MOH, Tehran, Iran

Aims: Crimean-Congo Hemorrhagic Fever (CCHF) is a zoonosis and a viral tick-born disease which caused by CCHF virus, belonging to *Nairovirus* genus and *Bunyaviridae* family, and is transmitted to humans by infected tick bite, handling of infected blood or tissue of infected livestock or nosocomially. In this survey we studied the rout of contamination of Iranian children Patients to this virus.

Methods: Between Iranian children (2month-14years) who are contaminated in recent years to CCHF in different province we did investigations for finding rout of transmission of disease to these children.

Result: In studied children we found that rout of transmission of CCHF in 97% were contact with infected blood and/or exudates of infected livestock and only 3% were infected by tick bite and no nosocomial infection has been observed among them.

Conclusions: In contaminated area and provinces in IRAN such as neighborhood and near area to Pakistan and Afghanistan (Endemic area) we observed more children cases. As we mentioned in our results the main rout of transmission of CCHF in Iranian children is through out handling with contaminated blood and or tissue of infected livestock. So for control of this disease in Iranian children it seems that awareness and teaching program for children and their parent regarding rout of transmission of CCHF and prevention programs is very helpful and important for public health and also for children in endemic area.

903

CESAREAN SECTION "GOLDEN RULE "FOR OBSTETRICIAN BUT NOT FOR NEWBORN

D. Kerleshi, M. Batusha, J. Pruthi

Regional Hospital of Gjakova, Gjakova, Serbia

Aim: The aim of the study is to compare parameters as neonatal ophtalmia, hypothermia, breastfeeding problems and early neonatal deaths between newborns delivered vaginally and by section cesarean.

Materials and methods: In the study has been analyzed 1047 newborn delivery facting sheet for period of time January July 2006. The data have been analyzed by statistical methods as percentage and Hi quadrant test.

Results: The results of our study shows that for the newborn that have a chance to be delivered vaginally, delivering via Cesarean Section (SC) is not a good choice. They are more in risk to have problems with breastfeeding and hypothermia. The percentage of hypothermia, breastfeeding problems and neonatal ophthalmia and early neonatal deaths is 11.8%, 20.9% and 19.2 %,12.2 in newborns delivered by SC vica 7.7%, 16.6% and 6.15 % ,3.69 in newborns delivered vaginally. Results processed with HI quadrant test shows that there is a significance difference $p < 0.05$,for hypothermia , and breastfeeding problems but there is no significant difference for neonatal ophtalmia $p > 0.05$.

Conclusions: From this study there is clear evidence that delivering with SC is more risky for the newborns regarding hypothermia and breastfeeding problems. **Keywords** Hypothermia, Neonatal hypothermia, breastfeeding problems, SC delivery.

904

SPECTRUM PRE-NATAL INFECTIONS AT NEWBORNS

M.I. Appasova¹, S.A. Achmetova²

¹Department of Pediatrics, Kazakh National Medical University, ²Center for Perinatology, Almaty, Kazakhstan

Background and aims: Premature birth, asphyxia, pre-natal infections and congenital anomalies- principal causes of infantile death rate in Kazakhstan.

Methods: We had been analysed 1582 case records of newborn children from hospital № 1 (a city of Almaty) for 3 years.

Results: We have chosen stories of newborns which have been surveyed on pre-natal infections, had the different reasons for inspection. It has been revealed that on toxoplasmos 103 newborns have been surveyed, antibodies are revealed at 7 (6,8 %). On a clamidiosis 192 newborns have been surveyed, positive serologie is revealed at 56 (29,2 %). On cytomegalovirus 166 newborns are surveyed, antibodies are found out at 67 (40,1 %), On a virus of a simple herpes is surveyed 97 newborns, antibodies are found out at 52 (53,6 %). It is necessary to note, Jg M have been revealed in single instances. All surveyed newborns with a positive caption of antibodies, had the cerebral ischemia, prolonged jaundice, deficiency of weight, 20 (28,5 %) children had pneumonia, at 5 (7,1 %) children the syndrome abstinent is revealed. At 30 newborns died from generals infections blood on sterility has been investigated, have been revealed shtamms - St.aureus-9,37 %, St.epidermidis-9,37 %, Enterobacter-15,62 %. The analysis anamnesis has shown that many children are born in families with low social level, to 40 % of mothers have the a chronic infection, and are carriers of pre-natal infections.

Conclusions: Antibodies were revealed at newborn most often to virus simple herpes (53,6%) cytomegalovirus (40,1%), clamidiosis (29,2%).

905

THE ROLE IMMUNOMODULATIVE THERAPY IN ACUTE VIRAL MYOCARDITIS IN CHILDREN

L.S. Sulovic

Faculty of Medicine Priština, Kosovska Mitrovica, University Children's Hospital, Belgrade, Serbia

Summary: Myocarditis is defined, clinically and pathologically, as inflammation of the heart muscle. The autoimmune phase is belived to play a major role and the use immunosuppressive agents may be useful in containing myocyte destruction. Aims was to systematically review the impact of Immunomodulative therapy on the outcome of severe and acute myocarditis in children. We retrospectively studied all the children admitted to University Children s Hospital with an acute viral myocarditis (AVM) from 2004-2009 years. Patients were included if they had 1.severe and acute heart failure, 2 left ventricular dysfunction assessed by echocardiography, 3. a recent history of viral illness and 4. absens of personal or familial history of cardiomyopathy. Fourteen patient (37%) had severe acute myocaditis. Seven patients were treated with immunomodulative therapy and anticongestive therapy (group I). Other seven patients were treated with anticongestive therapy only (group II). We compared the use of antikongestiv therapy alone, with a combination of immunomodulative. Such studies have documented succesful outcome of AVM with immunomodulative treatment.

906

COMPARISON THE EFFECTIVENESS OF BOILED BLACK TEA WITH BOILED WATER ON SEVERITY OF INFANTS' CONJUNCTIVITIS

M. Attarha¹, K. Vakillian¹, P. Yosefi²

¹Midwifery, ²Pediatric, University of Arak Medical Sciences, Arak, Iran

Objective: To compare the efficacy of boiled black tea with boiled water on severity of infants' conjunctivitis

Design: Randomized Control Clinical Trial

Setting: Pediatric Clinic of Arak Medical Science University in Iran

Participants: 165 infants with acute infective conjunctivitis