Objective: To assess the correlation between PTHrP level in preterm HM and PTHrP in maternal and neonatal serum.

Design/methods: We collected samples of expressed HM obtained from 16 mothers of preterm infants (25-34 weeks' gestation). In addition, blood samples from both mothers and infants were drawn. All samples were collected at days 2 and 10 postpartum. PTHrP levels were measured by two-site immunoradiometric assay. Blood calcium (Ca), Phosphorus (P) and alkaline phosphatase (ALP) levels were also measured.

Results: Both maternal and neonatal PTHrP serum levels did not vary after 10 days of breastfeeding. No correlation was found between PTHrP level in HM and serum PTHrP, Ca, P and ALP of the breastfed PTI.

Conclusions: Despite high levels of PTHrP in preterm HM, serum levels of PTHrP of the breast fed PTI do not increase over time. We found no correlation between PTHrP in HM and neonatal Ca serum concentration. We speculate that PTHrP intestinal absorption in the PTI is not significant.

1028

GASTROESOPHAGEAL REFLUX IN CRITICALLY ILL CHILDREN

M.J. Solana, C. Sanchez, J. Lopez-Herce, M. Crespo, A. Sanchez, J. Urbano, M. Botran, J.M. Bellon, A. Carrillo, Gregorio Maran PICU

Gregorio Maran General University Hospital, Madrid, Spain

Introduction: Gastroesophageal reflux (GER) is common in children, but its incidence in critically ill pediatric patients is unknown. Our objective was to study the incidence and type of GER in critically ill children.

Methods: Prospective observational study in critically ill children aged 1 month to 14 years. A multichannel catheter (MII) was inserted at the admittance in the PICU. This catheter has 6 channels for esophageal multichannel intraluminal impedance (MII) and 2 channels for esophageal and gastric pH measurement. Patients were studied during 48 hours.

Results: Twenty-four patients (mean age, 16.4 months) were studied. MII detected the presence of GER in 80% of patients, whereas pH monitoring

only detected 20%, although the correlation between the two was good. MII detected 320 episodes of GER (58% acid, 42% alkaline; 143 were gas type) whereas pH monitoring detected 26 episodes (100% acid). Patients with most episodes of GER were children between 7 and 12 months. The median duration of the longest episodes of GER was 4.1 minutes by pH monitoring and 2.3 minutes by MII. There was no relationship between the PRISM or PIM scores and the number of episodes of GER. There were no differences in the number or types of reflux according to the administration of epinephrine, dopamine, midazolam, fentanyl, or vecuronium.

Conclusions: The incidence of GER in critically ill children is high. MII is more sensitive than pH monitoring because many GER episodes are alkaline or weak acid.

1029

POLYMORPHIC VARIANTS OF UGT1A1 IN NEONATAL JAUNDICE IN SOUTHERN BRAZIL

C.G. Carvalho¹, S.M. Castro², A.P. Santin², L.A. de Azevedo², M.L. Pereira³, R. Giugliani³

¹UFRGS - HCPA, ²UFRGS, ³Genetic Service, UFRGS - HCPA, Porto Alegre, Brazil

Background and aims: Neonatal jaundice seems a benign condition but babies must be monitored in order to identify those who are at higher risk of encephalopathy. Abnormalities in hepatic conjugation of bilirubin were proposed in those cases - polymorphisms of the hepatic enzyme UGT1A1. Herein, we estimated the frequency of alleles and of genotypes of the promoter region of UGT1A1 gene in newborns and evaluated its association with severe hyperbilirubinemia

Methods: Case-control study, including all neonates who were in phototherapy at NICU/HCPA from March to December 2007, born > 35w of gestational age and weight > 2000g. Controls were neonates not jaundiced. Informed consent form applied. PCR performed and analyzed in GeneMapper® program

Results (490 NBs included): There were 7 of the 10 possible UGT1A1 genotypes identified; those related to Gilbert's syndrome were in 16%. Prevalence of polymorphic genotypes in icteric patients was 13.5% and in normal patients was 18.2%, p=0.08. In 86.6% of white neonates - low-risk genotypes, 13.4%, Gilbert's genotypes, while