among blacks, 34% showed the latter genotypes, and 76%, low-risk ($p=0.001$).

Conclusions: In the present sample the presence of polymorphic variants of UGT1A1 could not be associated to severe hyperbilirubinemia. Variants showed allelic prevalence and frequencies similar to those observed in other populations, with a highlight to the genotypes of higher risk in blacks and mulattos. Possibly due to the high miscegenation found in our state, other factors and genic interactions should be sought in order to explain severe neonatal hyperbilirubinemia, including the study of other polymorphisms.

1030

VITAMIN D AND OTHER PARAMETERS OF CALCIUM AND PHOSPHATE METABOLISM IN HEALTHY TERM NEWBORNS AFTER BIRTH

J. Dort¹, M. Bayer², E. Dortova¹, V. Dubova¹,
A. Mockova¹

¹Dpt. of Neonatology, Univesity Hospital, Pilsen,

²Dpt. of Pediatrics, 1st Medical Faculty, Charles University, Prague, Czech Republic

Background: The literary information about vitamin D saturation in newborns is rare. Levels of vitamin D in healthy term newborns in the Czech Republic have not been examined yet. The vitamin D saturation in newborns depends mainly on maternal saturation during gravidity that may be decreased due to nutritional habits and climatic conditions in the Middle Europe.

Objective: To get comparative values for examining of bone metabolism in newborns with very-low-birth weight.

Design: A prospective study conducted in spring period 2006 in a cohort of 28 healthy term caucasian newborns with normal birth weight . Blood and urine samples were drawn at the age of 3-7 days for examination of 25-hydroxyvitamin D (25/OH/D), calcium (Ca), phosphate (P), alkaline phosphatase (ALP), osteocalcin (OC), intact parathormone (PTH) in serum and calcium and phosphate in urine.

Results: Though other parameters of Ca and P metabolism were in normal range, levels of 25/OH/D were considerably lower than reference values found in literature . A significant correlation was found between 25/OH/D level and OC level, and futher a negative correlation (but not significant) of 25/OH/D and PTH.

Conclusion: Term newborns in the Czech Republic are born with an insufficient supply of vitamin D in the spring period obviously. The significant correlation of 25/OH/D level with bone formation marker (OC) and the tendency to higher PTH concentrations in newborns with lower 25/OH/D levels support the premise of an adverse effect of low vitamin D supply on long-term bone development.

Keywords: Vitamin D, newborn, bone development

1031

A COMPARISON OF THE WHO CHILD GROWTH STANDARDS AND THE CZECH GROWTH REFERENCES: PREVALENCE OF WASTING AMONG INFANTS

J. Vignerova¹, J. Riedlova², M. Paulova¹,
L. Humenikova³

¹Dept. of Children and Adolescents, Institute of Public Health, ²Dept. of Anatomy, 3rd Faculty of Medicine, Charles University, Praha, Czech Republic, ³Dept. of Nutritional Sciences, Oklahoma State University, 311 Human Environmental Sciences, Stillwater, OK, USA

Subjects: The study (grant IGA Mof H grant No. NS 9974, 4/2008) utilized the 2006 WHO Child Growth Standards and the 1991 Czech growth references. The WHO standards were based on a longitudinal study of 882 children aged 0-24 months and on cross-sectional studies of 6,669 children aged 18-71 months. The 1991 Czech growth references were based on a cross-sectional survey including 90,910 children aged 0-18 years (4,164 were children < 5 years).

Methods: The overall ratio of wasting in the sample of Czech children from the 1991 survey was calculated using the WHO Anthro software (version 2) and the WHO criteria for wasting.

Results: The prevalence of wasting was significantly higher among Czech children when using the WHO growth standards compared to the Czech references. The prevalence of wasting among 0-5 month old children was 15.5 % among boys and 12.9 % among girls compared to the expected 2.3% of the WHO standards.

Conclusions: The application of the WHO growth standards may results in a significant increase in the prevalence of wasting among Czech children, especially among infants aged 0-5 months. The performance and potential impacts of the WHO

child growth standards should be evaluated further before their adoption in the Czech Republic and other countries with local growth references.

1032

DIETARY FATTY ACID COMPOSITION DURING PREGNANCY AND RISK OF ASTHMA IN THE OFFSPRING

M. Lumia^{1,2}, P. Luukkainen³, M. Kaila⁴, H. Tapanainen¹, M. Erkkola⁵, L. Uusitalo^{1,2}, S. Niinistö^{1,6}, M.G. Kenward⁷, J. Ilonen^{8,9}, O. Simell¹⁰, M. Knip^{3,11,12}, R. Veijola¹³, S.M. Virtanen^{1,2,14}

¹Nutrition Unit, Department of Lifestyle and Participation, National Institute of Health and Welfare, Helsinki, ²Tampere School of Public Health, University of Tampere, Tampere, ³Hospital for Children and Adolescents, Helsinki University Hospital, Helsinki, ⁴Paediatric Research Centre, Tampere University Hospital and University of Tampere, Tampere, ⁵Division of Nutrition, Department of Food and Environmental Sciences,

⁶Department of Public Health, University of Helsinki, Helsinki, Finland, ⁷Department of Epidemiology and Population Health, Medical Statistic Unit, London School of Hygiene & Tropical Medicine, London, UK, ⁸Immunogenetics Laboratory, University of Turku, Turku, ⁹Department of Clinical Microbiology, University of Eastern Finland, Kuopio, ¹⁰Department of Pediatrics, University of Turku, Turku, ¹¹Folkhälsan Research Institute, University of Helsinki, Helsinki, ¹²Department of Pediatrics, Tampere University Hospital, Tampere, ¹³Department of Pediatrics, University of Oulu, Oulu, ¹⁴Paediatric Research Centre and Research Unit, Tampere University Hospital and University of Tampere, Tampere, Finland

Background: Epidemiological studies suggest that the increasing frequency of asthma and allergic diseases are in part related to changes in dietary habits. More margarines and vegetable oils are consumed, while the reverse is true for oily fish and fish-products, leading to an increase in the intake of n-6 polyunsaturated fatty acids (PUFA) and decrease in the intake of n-3 PUFA. Fatty acids (FA) modulate the immune system and have been proposed to affect the incidence of IgE-mediated allergic diseases. We explored the association of dietary FA composition during pregnancy with the risk of asthma in the offspring.

Methods: Dietary intake was assessed by a validated 181-item food frequency questionnaire covering the 8th month of pregnancy. The occurrence of asthma was assessed at the age of 5 years with a questionnaire modified from International Study of Asthma and Allergies in Childhood (ISAAC). Logistic regression was used for statistical analyses.

Results: Lower maternal intake of α -linoleic acid (18:3n-3), linoleic acid (18:2n-6), n-6-PUFA, n-3-PUFA and total PUFA during pregnancy were associated with an increased risk of asthma in the offspring. The ratios of n-6 to n-3-PUFA and 18:2n-6 to 18:2n-3 and the maternal intake of oils, fish and fish products were not associated with the risk of asthma.

Conclusion: In the present study, higher intakes of PUFA, n-3-PUFA and 18:3n-3 during pregnancy were protective against asthma in the offspring. The finding that higher intake of total n-6-PUFA and 18:2n-6 also protects against asthma does not support the original lipid hypothesis.

1033

PECULIARITIES IN FATTY ACIDS PROFILE IN CHILDREN WITH FOOD ALLERGY

M. Giovannini¹, E. Riva¹, S. Ruotolo², E. D'Auria³, G. Terrin², M. Tardi², L. Cosenza², A. Coruzzo², L. Leone³, C. Agostoni⁴, R. Troncone², R. Berni Canani²

¹Department of Pediatrics, San Paolo Hospital, University of Milan, Milan, ²Department of Pediatrics /University of Naples Federico II, ³European Laboratory for the Investigation of Food Induced Diseases (ELFID), ⁴Department of Maternal and Pediatric Sciences /University of Milan/ Fondazione IRCCS Ospedale Maggiore, Naples, Italy

Elimination diet (ED) in children with food allergy (FA) could be responsible of inadequate caloric and essential fatty acids (EFA) intake. We investigated the plasmatic profile of EFA in children with FA on ED. **Methods.** Prospective multicenter study including children with FA (6-36 m) on ED for at least 30 days compared to healthy children. EFA were measured by capillary gas-chromatography at the enrolment (T0) and after 6 m (T1) of elimination diet established after specific dietary counseling. **Results.** We enrolled 86 children with FA, and 66 healthy controls. The length of ED diet was 10.8 m (95%CI 8-14 m). Table reports EFA profiles in